



Length-weight Relationship and Condition Factors of Seven Penaeus Shrimp Species along the Southeast Coast of Tamil Nadu (Southern India)

Hari Prasad Mohale¹, P. Jawahar², N. Jayakumar¹, G. Arul Oli³, T. Ravikumar⁴, R.V. Bhosle⁵, Swapnil Narsale Ananda¹, Rishikesh Venkatrao Kadam¹, M.C. Sonwal¹

10.18805/IJAR.B-5218

ABSTRACT

Background: The present study of the length-weight relationship and condition factors of seven shrimp species on the southeast coast of Tamil Nadu between July (2022) and February (2023) also estimated the coefficient of determination (r^2).

Methods: For shrimp species, the length-weight relationship shows some species have significant positive coefficients which have a determination by ($r^2 > 0.5$) like *Penaeus indicus*, *Penaeus merguensis*, *Penaeus canaliculatus*, *Penaeus latissulcatus*, *Penaeus monodon*, some species have shown negative as ($r^2 > 0.5$) value like *Penaeus japonicas* and *Penaeus semisulcatus*.

Result: All shrimp species have heterogeneity evident in intercept (a). The growth exponent (b) varied between 0.278 and 2.289 and it significantly ($p=0.05$) deviated from the value of <3 which denoted negative allometric growth for all *Penaeus* shrimp species.

Key words: Allometric growth, Condition factor, Length-Weight relationship, *Penaeus* shrimp.

INTRODUCTION

Around the world, coastal waters with warm temperatures and tropical climates gather the expensive seafood known as shrimp. In many parts of the world, shrimp fisheries provide economic assistance (Khademzadeh and Haghi, 2017; Olawusi *et al.*, 2014; Ajani *et al.*, 2013). The penaeid shrimp is found in a large area, from the west and east coasts of India to Hong Kong, to the waters of Malaysia and Indonesia, to tropical Australia and New Guinea. Most species are in the Indo-Pacific region, from the Persian Gulf through India, the east coast of Africa, Japan and Australia. These species are found in shallow inshore seas where spawning adults are frequently seen and they are also primarily fished in shallow coastal areas (Amani *et al.*, 2015). India has a sizable marine fishing industry contributes significantly to the national economy and has provided millions of people living in our nation with subsistence and employment opportunities in coastal communities (Rao *et al.*, 2023). In addition to fish, prawns play a significant ecological and economic role in capture fisheries. The foundation of the seafood export business is the penaeid prawn, which also generates most of the sector's foreign exchange earnings. Millions of people, directly and indirectly, depend on the fishery sector for their livelihood (Johnson, 2010; Joseph and Jayaprakash, 2003; Devaraj and Vivekanandan, 1999; George *et al.*, 1981). The crustacean fisheries resources around Palk Bay's shore are mostly harvested using four different types of gear: bottom-set gillnet, gillnet (a specialised gillnet for prawns) and thalluvalai (pushnet), which is used in the artisanal sector. Trawling is the mechanised sector. For the past few decades, Palk Bay's inshore waters have been used for thalluvalai operations

¹Department of Fisheries Biology and Resource Management, Fisheries College and Research Institute, Tamil Nadu Dr. J. Jayalalitha Fisheries University, Thoothukudi-628 008, Tamil Nadu, India.

²Tamil Nadu Dr. J. Jayalalitha Fisheries University, Nagapattinam-611 002, Tamil Nadu, India.

³Department of Fisheries Extension, Economic and Statistics, Fisheries College and Research Institute, Thoothukudi-628 008, Tamil Nadu, India.

⁴Department of Fishing Technology and Fisheries Engineering, Fisheries College and Research Institute, Thoothukudi-628 008, Tamil Nadu, India.

⁵Department of Aquaculture, College of Fisheries Science, Udgir-413 517, Maharashtra, India.

Corresponding Author: Hari Prasad Mohale, Department of Fisheries Biology and Resource Management, Fisheries College and Research Institute, Tamil Nadu Dr. J. Jayalalitha Fisheries University, Thoothukudi-628 008, Tamil Nadu, India. Email: haricof92@gmail.com

How to cite this article: Mohale, H.P., Jawahar, P., Jayakumar, N., Oli, G.A., Ravikumar, T., Bhosle, R.V., Ananda, S.N., Kadam, R.V. and Sonwal, M.C. (2024). Length-weight Relationship and Condition Factors of Seven *Penaeus* Shrimp Species along the Southeast Coast of Tamil Nadu (Southern India). Indian Journal of Animal Research. 58(8): 1383-1392. doi: 10.18805/IJAR.B-5218.

Submitted: 09-08-2023 **Accepted:** 21-01-2024 **Online:** 01-03-2024

aimed at prawn harvesting (Rajamani and Palanichamy, 2009). As a result, all species members of the Penaeidae family are referred to as "prawns" in this research. In Indian water, the Penaeidae family has 17 genera and 78 species. Most of the penaeid prawn species in Indian water belong

to the genus *Penaeus*, which is of commercial significance (Chanda, 2017). The main commercial shrimp species landed from the genera *Metapenaeus*, *Parapenaeopsis* and *Penaeus*. The procedure uses prawn trawl nets with a cod end mesh size of 10-20 mm in both the northern and southern orientations. With a crew of six, these trawlers can hold two to three tonnes of fish (Maheswarudu *et al.*, 2015). The primary tool used to exploit almost 90% of the penaeid prawn along the east coast is a trawl net. There are various varieties of trawlers based on the boat's size, engine capacity and gear size. These boats are Sona (13.1 m), Sorrah (11.4 m), Pablo (9.14 m) and Royya (9.75-10 m). The principal landing grounds and fishing harbours, which serve as the foundation for trawler operations, include Tuticorin, Mandapam, Kakinada, Digha, Paradeep, Visakhapatnam and Diamond Harbour. In addition to these trawlers, the coast of Tamil Nadu is home to Halluvalai (Sreeram *et al.*, 2014).

In fishery biology, ecology and fish stock assessment, the length-weight relationship (LWR) and length-length relationship (LLR) of shrimp are significant instruments (Karna, 2017). For comparing growth studies in fisheries management, length-weight connections are crucial. It aids in determining the two variables' mathematical relationship, allowing the conversion of one variable to characterise the growth in the wild to identify potential variations among several stocks of the same species, delineates the stocks and facilitate comparative growth research (Das *et al.*, 2021; Ahmadi, 2018). The length-weight connection can be a good indicator of environmental factors that may affect how energy is taken in and used by marine fish and crustaceans (Das *et al.*, 2021). An animal's physical health is frequently measured using Fulton's condition factor (K), which is also a helpful addition to a crustacean's growth estimate (Li *et al.*, 2016). Shrimp feeding intensity, age and growth rates may all be monitored using the condition factor, which is also beneficial. Both biotic and abiotic ecological circumstances heavily influence it and can be used as a gauge for the health of the aquatic habitat where fish reside (Prajapati and Ujjania, 2021; Solanki *et al.*, 2020; Uddin *et al.*, 2016). Fish and crustaceans' variations in the length-weight relationship are good predictors of energy absorption and allocation changes (Kop *et al.*, 2019; Qadri *et al.*, 2017). Further differences are influenced by various circumstances, including food, stress brought on by crowding, or the reproductive cycle (Oyebamiji *et al.*, 2008). The current study aims to add and complement already-existing knowledge about the shrimp growth pattern (Siddique *et al.*, 2022). To manage shrimp resources sustainably and also give relevant information on the length-length, length-weight connection (growth patterns) and condition factor on shrimps caught in coastal waters from the southeast coast of Tamil Nadu (Suryanti *et al.*, 2018; Olawusi *et al.*, 2014).

MATERIALS AND METHODS

The present study was conducted from July to February 2022-2023 at the sampling site from Thoothukudi (Station I),

Nagapattinam (Station II), Mandapam (Station III) and Palk Bay (Station IV) on the southeast coast of Tamil Nadu (Fig 1). Every month, shrimp samples were taken at depths of 100 to 150 meters in the Gulf of Mannar (Tamil Nadu). Samples were collected once or twice a month during the shrimp landing period from the sampling sites in Thoothukudi, Mandapam and Nagapattinam using a commercial bottom trawler and bottom set gillnet. The net's mesh size measured 20.2 mm at the cod end and 30.9 mm in the net body. Before being transferred to the Fisheries College and Research Institute, Laboratory of the Thoothukudi, the samples were promptly refrigerated in ice on board and treated with 5% formalin. They were identified using FAO Species Identification Sheets (Volume VI). Based on the presence of thelycum in females and petasma in males, all obtained specimens were sexed. The overall length (from the tip of the rostrum to the edge of the telson) was measured using a graded measuring board and the weight was determined using an electronic weighing balance.

Characters for diagnosis

Small to large, with a body length ranging from 2.5 to 35 cm. All five pairs of legs are fully formed, with the first three generating pincers that aren't huge abdomen with the front portion of the subsequent pleura covered by the posterior part (lateral plates). A huge copulatory apparatus is on the first pair of pleopods in males (petasma) and on the posterior thoracic sternites in females (thelycum). The females do not hold the eggs on their abdomens; instead, they release them straight into the water (Josileen, 2022; Abdurraheem *et al.*, 2021; Das *et al.*, 2021; Rao *et al.*, 2013; Rajkumar *et al.*, 2015). Among the many different sizes of crustaceans, shrimps range from tiny to approximately 35 cm in length. Shrimps often have a lengthy abdomen longer than their head or carapace, a compressed and toothed rostrum and an almost invariably laterally compressed body. Generally speaking, the antennae and antennules are big and plate-like. Typically, pereopods have slim legs, but some species have robust legs, either a single or a pair and some end in pincers or chelae. Except in a few species, all five anterior abdominal segments have well-developed pleopods or abdominal appendages utilised for swimming (Piratheepa *et al.*, 2016).

The length-weight relationship was calculated by the following formula, $W = a \cdot L^b$, i.e., $\log W = \log a + b \log L$ was estimated by linear regression analysis (Ricker, 1973) where a is intercept and b is the slope of linear regression on the log-transformed weight (g) and length data (cm) respectively (Froese, 2006; Froese *et al.*, 2011). The pattern of growth, i.e., negative showing allometry ($b < 3$), isometry ($b = 3$), or positive allometry ($b > 3$), was tested by using regression analyses (ANOVA) and setting a significance level of 0.05, p-value. The attributed data error and extreme outliers were excluded from the calculations. Using the least square residuals method, the optimized values of "a and b" were computed (Haddon, 2011). The statistical significance, 95% confidence intervals (CIs) of the parameters a, b and

coefficient of determination (r^2) were also estimated. The length-length relationships (LLRs) between TL and CL were also established using linear regression analysis:

$$CL = a + b \times TL$$

The Fultons condition factor (CF) of shrimp species was calculated as the body weight expressed as a percentage of the cube of the Total length following the formula (Pauly, 1983).

$$\text{Condition factor (K)} = \frac{100W}{L^3}$$

Where:

K = Condition factors.

L = Total length (cm).

W= Weight of shrimps (gm).

RESULTS AND DISCUSSION

Table 1 displays the length-weight relationship and the estimated coefficient of determination (r^2) for all seven shrimp species (Fig 2). The length-weight connection demonstrated a statistically significant positive coefficient of determination ($r^2 > 0.5$) like *Penaeus indicus*, *Penaeus merguensis*, *Penaeus canaliculatus*, *Penaeus latisulcatus*, *Penaeus monodon* and some species have shown negative as ($r^2 > 0.5$) value like *Penaeus japonicas* and *Penaeus semisulcatus*. All shrimp species have heterogeneity evident in intercept (a). The growth exponent (b) varied between 0.278 and 2.289 and it significantly ($p = 0.05$) deviated from the value of < 3 , which denotes negative allometric growth for all species, as mentioned in (Table 2), which is shown in Fig 3-9 the length-weight graphs




Fig 1: Study area of southeast coast of Tamil Nadu.

Table 1: The length-length relationships of seven *Penaeus* Shrimp species from the marine coast of Tamil Nadu.

Species	Length-length characters			
	Equation	a	b	r^2
<i>Penaeus indicus</i>	CL= a+b×TL	0.268	0.932	0.886
	BL= a+b×TL	3.878	0.282	0.459
<i>Penaeus merguensis</i>	CL= a+b×TL	0.678	0.433	0.53
	BL= a+b×TL	0.62	0.924	0.915
<i>Penaeus japonicus</i>	CL= a+b×TL	0.190	0.842	0.827
	BL= a+b×TL	0.698	0.802	0.912
<i>Penaeus canaliculatus</i>	CL= a+b×TL	0.663	0.437	0.609
	BL= a+b×TL	0.207	3.390	0.775
<i>Penaeus latisulcatus</i>	CL= a+b×TL	0.787	0.370	0.551
	BL= a+b×TL	2.271	0.381	0.721
<i>Penaeus monodon</i>	CL= a+b×TL	0.715	0.474	0.856
	BL= a+b×TL	2.677	0.333	0.572
<i>Penaeus semisulcatus</i>	CL= a+b×TL	0.268	0.932	0.886
	BL= a+b×TL	4.048	0.266	0.410

for each species of shrimp. The condition factor (K) analysis has demonstrated that female shrimp typically have higher K-values than male shrimp. The K-value varied from 0.38 (*P. semisulcatus*) to 1.79 (*P. japonicus*). The outcome suggested that heavier shrimps of a particular length are in better shape. Also, the study discovered that the condition factor varied between shrimp sexes and that female shrimp generally had higher K-values than male shrimp, as mentioned in (Table 3). The maximum condition factor of *P. japonicus* males (K=1.56) and the lowest value of

P. semisulcatus (K=0.38) were found in December 2022 and February 2023, respectively. In the case of Females, shrimp in *P. semisulcatus* had the lowest condition factor in January 2023 (K=0.58) and the highest condition factor in September (K=1.79) *P. japonicus*.

The results of the length-weight relationship (LWR) of shrimps in marine water revealed a strong and substantial connection between the overall length and weight of shrimp ($r^2= 0.015-0.953$) similarity observation found by Aye *et al.* (2019). In other words, shrimp grow in length and weight




Fig 2: Collected shrimp sample from the southeast coast of Tamil Nadu.

Table 2: The length-weight relationship of penaeus shrimps from the marine coast of Tamil Nadu.

Species	N	a	b	r ²	Growth pattern
<i>Penaeus indicus</i>	106 (F)	0.169	1.595	0.630	(-) Allometric
	70 (M)	0.143	1.705	0.773	(-) Allometric
	176 (C)	0.156	1.645	0.688	(-) Allometric
<i>Penaeus merguensis</i>	115 (F)	0.103	1.817	0.885	(-) Allometric
	26 (M)	0.139	1.676	0.940	(-) Allometric
	141 (C)	0.109	1.790	0.893	(-) Allometric
<i>Penaeus japonicus</i>	105 (F)	0.330	1.531	0.414	(-) Allometric
	36 (M)	4.19	0.278	0.015	(-) Allometric
	141 (C)	0.451	1.377	0.341	(-) Allometric
<i>Penaeus canaliculatus</i>	106 (F)	0.606	1.212	0.573	(-) Allometric
	46 (M)	0.586	1.234	0.563	(-) Allometric
	152 (C)	0.600	1.219	0.569	(-) Allometric
<i>Penaeus latisulcatus</i>	73 (F)	0.543	1.260	0.544	(-) Allometric
	28 (M)	0.533	1.268	0.556	(-) Allometric
	101 (C)	0.540	1.262	0.547	(-) Allometric
<i>Penaeus monodon</i>	52 (F)	0.037	2.289	0.953	(-) Allometric
	22 (M)	0.046	2.194	0.942	(-) Allometric
	74 (C)	0.040	2.259	0.949	(-) Allometric
<i>Penaeus semisulcatus</i>	111 (F)	0.470	1.296	0.161	(-) Allometric
	33 (M)	0.580	1.107	0.463	(-) Allometric
	144 (C)	0.482	1.261	0.179	(-) Allometric

similarly. The species-specific coefficient of determination $r^2 > 0.5$ results demonstrated the model's suitability for this investigation. The slenderness or stoutness of the shrimp could be attributed to dimensional discrepancy observed

in shrimps living in maritime environments. The shrimp gets skinny as it grows longer, showing negative allometric growth. As a result, an analysis of the length-weight relationship offers a crucial tool for assessing shrimp




Fig 3: The graph of the length-weight relationship for combined sex of *P. indicus*.




Fig 4: The graph of the length-weight relationship for combined sex of *P. merguensis*.




Fig 5: The graph of the length-weight relationship for combined sex of *P. japonicas*.

population stock and production potential. Positive allometric growth was revealed by the growth coefficient for both sexes across all species, demonstrating that the rise in body length rate is not inversely correlated with the

increase in body weight. The present study observed the negative allometric growth of all the Penaeus shrimp species, which means it grows longer than the stock population. Similar results were found by Torres *et al.* (2017)




Fig 6: The graph of the Length-Weight Relationship for combined sex of *P. canaliculatus*.




Fig 7: The graph of the length-weight relationship for combined sex of *P. latisulcatus*.




Fig 8: The graph of the length-weight relationship for combined sex of *P. monodon*.

Table 3: The mean length, weight and condition factor (K) for the Penaeus Shrimp species from the marine coast of Tamil Nadu.

Species	Female						Male						Combine sex				
	N	TL (cm)		TW (g)		K	N	TL (cm)		TW (g)		K	N	TL (cm)		TW (g)	
		Min - Max	95% CL	X±SE	Min - Max			95% CL	X±SE	Min - Max	95% CL			X±SE	Min - Max	95% CL	X±SE
<i>P. indicus</i>	106	7.83±0.15	4.70±0.16	1.08	70	7.87±0.31	5.04±0.23	1.03	176	7.84±0.12	4.84±0.13	1.05		7.84±0.12	4.84±0.13	1.05	
		5-12.5	1.5-9.5			5-12.5	1.5-9.5			5-12.5	1.5-9.5			5-12.5	1.5-9.5		
		0.298	0.330			0.420	0.465			0.242	0.272			0.242	0.272		
<i>P. merguensis</i>	115	7.91±0.16	4.66±0.19	0.99	26	7.81±0.35	4.35±0.38	0.96	141	7.89±0.14	4.64±0.17	0.97		7.89±0.14	4.64±0.17	0.97	
		5-12	2-11			5-12	2-10.5			5-12	2-11			5-12	2-11		
		0.321	0.387			0.731	0.797			0.290	0.344			0.290	0.344		
<i>P. japonicus</i>	105	8.91±0.24	11.00±0.82	1.79	36	8.11±0.25	8.00±0.45	1.56	141	8.71±0.19	10.23±0.63	1.71		8.71±0.19	10.23±0.63	1.71	
		6-17	3-36			6-11.5	3-12.5			6-17	3-36			6-17	3-36		
		0.493	1.640			0.516	0.914			0.391	1.256			0.391	1.256		
<i>P. canaliculatus</i>	106	11.71±0.16	2.18±0.27	0.86	46	11.69±0.26	12.40±0.44	0.82	152	11.70±0.13	12.25±0.23	0.81		11.70±0.13	12.25±0.23	0.81	
		7-14.5	5-20			7-14.5	5-20			7-14.5	5-20			7-14.5	5-20		
		0.330	0.546			0.52	0.904			0.276	0.463			0.276	0.463		
<i>P. latisulcatus</i>	73	11.75±0.19	12.36±0.35	0.82	28	11.76±0.30	12.37±0.55	0.80	101	11.75±0.16	12.36±0.29	0.80		11.75±0.16	12.36±0.29	0.80	
		7-14.5	5-20			7-14	5-20			7-14.5	5-20			7-14.5	5-20		
		0.394	0.703			0.634	1.143			0.329	0.588			0.329	0.588		
<i>P. monodon</i>	52	8.95±0.27	6.08±0.44	0.85	22	8.95±0.46	6.18±0.73	0.84	74	8.95±0.23	6.11±0.37	0.82		8.95±0.23	6.11±0.37	0.82	
		5-13.5	1.5-14			5-13.5	1.5-14			5-13.5	1.5-14			5-13.5	1.5-14		
		0.557	0.890			0.968	1.530			0.473	0.753			0.473	0.753		
<i>P. semisulcatus</i>	111	15.60±0.28	20.91±1.67	0.58	33	15.48±0.55	12.80±1.38	0.38	144	15.57±0.25	19.05±1.35	0.53		15.57±0.25	19.05±1.35	0.53	
		10-19.5	7-68			10-19.5	7-55			10-19.5	7-68			10-19.5	7-68		
		0.566	3.311			1.136	2.821			0.501	2.678			0.501	2.678		
Total	668				261				929								
	(71.90%)				(28.09%)												




Fig 9: The graph of the length-weight relationship for combined sex of *P. semisulcatus*.

from the Gulf of Cadiz (Spain) and Ahmadi (2018) studied negative allometric growth in female shrimp from the Barito River, Indonesia. One author from India also found that the negative allometric growth in both sexes of shrimp from the Digha coast of West Bengal has been studied by Uddin *et al.* (2016). Some other authors similarly study reported by (Das *et al.*, 2021; Ukpatu, 2021; Hasan *et al.*, 2020; Suryanti *et al.*, 2018; Khademzadeh and Haghi, 2017; Uddin *et al.* 2016; Amani *et al.*, 2015; Faye *et al.*, 2015; Fatima, 2001; Fontaine and Neal, 1971).

The variations may result from geographic conditions and regional shrimp growth patterns. The modest variance in b and r^2 values is acceptable given that a species' length-weight relationship may change depending on its environment and the time of year. The different physical traits shown in Table 3 clearly distinguish the sexes of shrimp in marine water. Many crustaceans have shown these alterations. Many species of commercially significant penaeid shrimp exhibit the same pattern of sexual dimorphism, in which the females are larger than the males. Compared to males of the same overall length, it was reported that females had longer carapaces and heavier bodies and this difference became more evident in larger-size groups. More weight and an increase in the number of moult cycles may contribute to larger female sizes. The condition factor (CF) can describe a shrimp's health or fatness. It varies based on sex, stage of development and season. The existence of gravid females or the larger size of the female gonads, which are absent in their male counterparts, may cause variances in the condition factor between males and females.

The greater K value for *P. japonicus* ($K= 1.71$) obtained in this study can be explained by the fact that this species grows bigger and quicker than other Penaeid shrimps, according to the usage of condition factor as an index of growth and feeding intensity. The lower k value for *P. semisulcatus* ($K= 0.80$) may be due to its high feeding rate, severe overexploitation and marine habitat. Ajani *et al.* (2013) conducted a similar investigation on two Penaeid

shrimps from the coastal state of Lagos in South West Nigeria. Likewise, observations were made by (Suryanti, 2021; Solanki *et al.*, 2020). According to Silva *et al.* (2021), the condition factor is a sign of changes in food stores. The individual condition component, as it relates to the health and fatness of the organisms, may cause weight variations for all species of shrimp. The combined sex condition factors were $K= 0.53$ for *P. semisulcatus*, $K= 0.80$ for *P. latisulcatus*, $K= 0.81$ for *P. canaliculatus*, $K= 0.82$ for *P. monodon*, $K=0.97$ for *P. merguensis*, $K= 1.05$ for *P. indicus* and $K= 1.71$ for *P. japonicus* differed from the record in the coastal waters of Tamil Nadu. According to a study present study, Olawusi-Peters *et al.* (2014) also reported the similarity condition factors value. These variations might result from oil reserves and the pollution level in Tamil Nadu maritime waters, which could impact the biotic integrity of the shrimp species. The condition parameters in Tamil Nadu thermal industries were also affected by pollution. The sex, maturity, season and environmental factors may all have a role in shrimp's length-weight relationship and condition factor.

CONCLUSION AND ACKNOWLEDGEMENT

The LWRs and condition factor (CF) of seven Penaeus shrimp from Tamil Nadu marine waters were assessed in the current study, exhibiting a negative allometric trend. The present research is suggested to be in good condition based on an analysis of growth-related features and condition factors. The condition factor (K) analysis showed that female shrimp typically have higher K -values of 1.79 (*P. japonicus*) than male shrimp of *P. semisulcatus* (K -value was 0.38; the outcome suggests that shrimps of a particular length are heavier in better shape.

The author appreciates the continued support provided by the Dr. J. Jayalalithaa Fisheries University in Nagapattinam, Tamil Nadu and the dean of the Fisheries College and Research Institute in Thoothukudi, for his wise advice. I am pleased to convey my sincere appreciation and gratitude to Dr. P. Jawahar, my mentor and advisor at

Tamil Nadu's Dr. J. Jayalalithaa Fisheries University in Nagapattinam. My work had only been finished because of his commitment to labour and assistance. His prompt counsel and careful inspection greatly aided me in completing this assignment.

Conflict of interest

All authors declared that there is no conflict of interest.

REFERENCES

- Abdulraheem, S., Ibrahim, G.D. and Ahmed, M.S. (2021). Identification and taxonomic study of shrimps in Bardawil Lagoon, North Sinai, Egypt. *Sinai Journal of Applied Sciences*. 10(1): 39-46.
- Ahmadi, A. (2018). Morphometric characteristic of *Macrobrachium* sp. from Barito River, Indonesia. *International Journal of Innovative Studies in Aquatic Biology and Fisheries*. 4: 19-31.
- Ajani, E.G., Bello, O.B. and Osowo, O. (2013). Comparative condition factor of two Penaeid shrimps, *Penaeus notialis* (Pink shrimp) and *Penaeus monodon* (Tiger shrimp) in a coastal state, Lagos, South West Nigeria. *Journal of Natural Sciences*. 11(4): 1-3.
- Amani, A.A., Arshad, A., Yusoff, F.M. and Amin, S.M.N. (2015). Length-weight relationship and relative condition factor of *Parapenaeopsis sculptilis* (Heller, 1862) from the coastal waters of Perak, Peninsular Malaysia. *Pertanika Journal of Tropical Agricultural Science*. 38(2): 211-217.
- Aye, M.M., Htay, S.S., Win, S., Win, S.S. and Lwin, Y.Y. (2019). Relationship between Weight and Length of Shrimp/Prawn Species Recorded from U-To Creek, Chaungtha, Patheingyi Township, Ayeyawady Region. 4(7): 1119-1122.
- Chanda, A. (2017). A check list of penaeid prawns found in indian water with their distribution. *IOSR Journal of Pharmacy and Biological Sciences*. 12(03): 26-31.
- Das, R.R., Panigrahi, A., Saravanan, A., Ambikanandham, K. and Arumugam, S. (2021). Length Weight Relationship and Condition Factor (K) of *Penaeus indicus* (H. Milne-Edwards) Based on Developmental Stages, Grow Out Stages, Brood Stock Stages and Sex. *Poultry, Fisheries and Wildlife Sciences*. 9(8): 1-6.
- Devaraj, M. and Vivekanandan, E. (1999). Marine capture fisheries of India: Challenges and opportunities. *Current Science*. 314-332.
- Fatima, M. (2001). Study on length-frequency and length-weight relationship of *Penaeus japonicus* and *Parapenaeopsis sculptilis*. *Journal of Biological Science*. 1: 171-172.
- Faye, A., Sarr, A., Diouf, M., Thiaw, M., Fall, J., Ndiaye, I., Ba, K. and Lazar, N. (2015). Contribution to the study of the size structure, the length-weight relationship, the condition factor and the sex-ratio of shrimp *Farfante penaeusnotialis* (Pérez Farfante, 1967) in the estuary of Sine-Saloum (Senegal). *American Scientific Research Journal for Engineering, Technology and Sciences (ASRJETS)*. 14(2): 97-112.
- Fontaine, C.T. and Neal, R.A. (1971). Length-weight relations for three commercially important penaeid shrimp of the Gulf of Mexico. *Transactions of the American Fisheries Society*. 100(3): 584-586.
- Froese, R. (2006). Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology*. 22: 241-253.
- Froese, R., Tsikliras, A.C., Stergiou, K.I. (2011). Editorial note on weight-length relations of fishes. *Acta Ichthyologica et Piscatoria*. 41: 261-263.
- George, M.J., Suseelan, C. and Balan, K. (1981). By-catch of the Shrimp Fishery in India. *Marine Fisheries Information Service, Technical and Extension Series*. 28: 1-13.
- Haddon, M., (2011). *Modelling and Quantitative Methods in Fisheries*, 2nd edn. CRC Press, Taylor and Francis Group, New York. pp: 449.
- Hasan, M., Al Mamun, A. and Hossain, M. (2020). Length-weight relationships of 12 indigenous fishes and 3 shellfishes from mangrove and floodplain ecosystems in Southwestern Bangladesh. *Egyptian Journal of Aquatic Biology and Fisheries*. 24(5): 69-76.
- Johnson, D.S. (2010). Institutional adaptation as a governability problem in fisheries: Patron-client relations in the Junagadh fishery, India. *Fish and Fisheries*. 11(3): 264-277.
- Joseph, M.M. and Jayaprakash, A.A. (2003). Status of exploited marine fishery resources of India. Kochi: Central Marine Fisheries Research Institute. 157.
- Josileen, J. (2022). Overview of Crustacean Fisheries and Crab Taxonomy in India.
- Karna, S.K. (2017). Length-weight and length-length relationship of *Thryssa purava* (Hamilton, 1822), *Thryssapoly branchialis* Wongratana, 1983 and *Thryssa mystax* (Bloch and Schneider, 1801) from Chilikka lagoon, India. *Journal of Applied Ichthyology*. 33(6): 1284-1286.
- Khademzadeh, O. and Haghi, M. (2017). Length-weight relationship and condition factor of white leg shrimp *Litopenaeus vannamei* (Boone, 1931) in culture systems of Choebdeh, West-South of Iran. *International Journal of Fisheries and Aquatic Studies*. 5(1): 298-301.
- Kop, A., Korkut, A.Y. and Gurkan, S. (2019). Length-weight relationship and condition factor as an indicator of growth and feeding intensity of Sea bream (*Sparus aurata* L., 1758) given feed with different protein contents. *Indian Journal of Animal Research*. 53(4): 510-514. doi: 10.18805/ijar.B-998.
- Li, Y., Zhou, F., Ma, Z., Huang, J., Jiang, S., Yang, Q., Li, T. and Qin, J.G. (2016). Length-weight relationship and condition factor of giant tiger shrimp, *Penaeus monodon* (Fabricius, 1798) from four breeding families. *Springer Plus*. 5: 1-5.
- Maheswarudu, G., Sreeram, M.P., Dhanwanthari, E., Varma, J.B., Sajeev, C.K. and Rao, S.S. (2015). Exploitation of penaeid shrimp resources by small mechanised trawlers off Visakhapatnam andhra Pradesh. *Indian Journal of Fisheries*. 62(2): 7-13.
- Olawusi-Peters Olamide, O., AjibareAdefemi, O. and Bello-Olusoji Oluayo, A. (2014). Length-weight relationship and condition factor of shrimps in coastal waters of Ondo state, South West, Nigeria. *International Journal of Fisheries and Aquatic Studies*. 1(3): 137-142.
- Oyebamiji, F.O., Asiru, R.A. and Oladunjoye, R.Y. (2008). Length-weight relationship, abundance and sex ratio of the giant river prawn *Macrobrachium vollehovenii* (herklotz, 1857) from river Osun, Southwestern Nigeria. *Journal of Aquatic Research and Development*. 9(10).

- Pauly, J.E. (1983). Chronobiology: Anatomy in time. *Developmental Dynamics*. 168(4): 365-388.
- Piratheepa, S., Chitravadivelu, K. and Edrininghe, U. (2016). Taxonomic study and Identification key to the Species of the shrimps, particularly the Family Penaeidae in Kakkaiyaru coastal waters, Jaffna, Sri Lanka. *International Journal of Scientific and Research Publications*. 6(5): 424-432.
- Prajapati, S.D. and Ujjania, N.C. (2021). Study on length-weight relationship and condition factor of whiteleg shrimp *Litopenaeus vannamei* (Boone, 1931) cultured in earthen pond, Khambhat (Gujarat). *International Journal of Fauna and Biological Studies*. 8(1): 67-70.
- Qadri, S., Shah, T.H., Balkhi, M.H., Bhat, B.A., Bhat, F.A., Najjar, A.M., Asmi, O.A., Farooq, I. and Alia, S. (2017). Morphometrics and length-weight relationship of *Schizothorax curvifrons* Heckel 1838 in River Jhelum, Kashmir, India. *Indian Journal of Animal Research*. 51(3): 453-458. doi: 10.18805/ija.r.v0i0f.3787.
- Rajamani, M. and Palanichamy, S. (2009). Need to regulate the thalluvalai fishery along Palk Bay, southeast coast of India. *Journal of the Marine Biological Association of India*. 51(2): 223-226.
- Rajkumar, G., Bhavan, P.S., Udayasuriyan, R. and Vadivalagan, C. (2015). Molecular identification of shrimp species, *Penaeus semisulcatus*, *Metapenaeus dobsoni*, *Metapenaeus brevicornis*, *Fenneropenaeus indicus*, *Parapenaeopsis stylifera* and *Solenocera crassicornis* inhabiting in the coromandel coast (Tamil Nadu, India) using MT-COI gene. *International Journal of Fisheries and Aquatic Studies*. 2(4): 96-106.
- Rao, G.S., Radhakrishnan, E.V. and Josileen, J. (2013). *Handbook of Marine Prawns of India*. ICAR-Central Marine Fisheries Research Institute.
- Rao, N.V., Jawahar, P., Jayakumar, N., Sujathkumar, N.V. and Rani, V., Mohale H.P., Karuppasamy, K., Kalavathi, N.R. (2023). Length-weight relationship and condition factor of siluriformes fish species, endangered *Eutropiichthys vacha*, hamilton (1922) and endemic deccan catfish *Rita kuturnee* (sykes, 1839) of godavari river Andhra Pradesh, South India. *Indian Journal of Animal Research*. 1: 6. doi: 10.18805/IJAR.B-5222.
- Ricker, W.E. (1973). Linear regressions in fishery research. *Journal of the Fisheries Research Board of Canada*. 30: 409-434.
- Siddique, Z., Saddozai, S., Achakzai, W.M., Khan, N. and Rahim, M. (2022). Length-weight relationship and fecundity of *Parastromateus niger* from Gwadar Port Balochistan province, Pakistan. *Indian Journal of Animal Research*. 56(10): 1258-1263. doi: 10.18805/IJAR.BF-1414.
- Silva, A.R., Lopes, A.E.B., Grabowski, R.C. and Castilho, A.L. (2021). Weight/carapace length relationship and condition factor of the roughneck shrimp, *Rimapenaeus constrictus* (Stimpson, 1874), on the southeastern Brazilian coast. *Anais da Academia Brasileira de Ciências*. 93.
- Solanki, H.G., Ujjania, N.C., Gopal, C. and Pillai, S.M. (2020). Length-weight relationship, condition factor and length-frequency analysis of tiger shrimp (*Penaeus monodon* Fabricius, 1798). *International Journal of Fauna and Biological Studies*. 7(4): 191-195.
- Sreeram, M.P., Dhanwanthari, E., Saleela, K.N., Nair, K.R., Thangaraj Subramanian, V., Rajamani, M., Rao, G.S., Maheswarudu, G. and Unnithan, K.A. (2014). Penaeid prawn resources along the east coast of India during 1991-2011. *Marine Fisheries Information Service and Technology*. 8-14.
- Suryanti, A., Riza, N. and Raza'i, T.S. (2018). Length-weight relationship and condition factor of white shrimp *Penaeus merguensis* captured in ecosystem mangrove of Bagan Asahan, Tanjungbalai, Asahan, North Sumatra, Indonesia. In *IOP Conference Series: Earth and Environmental Science*. 122(1): 012108.
- Suryanti, S. (2021). Osmoregulation pattern and condition factor of Indian white shrimp (*Penaeus indicus*) in the mangrove eco-ecotourism area of Tapak, Semarang. *AACL Bioflux*, 14(4).
- Torres, M.A., Vila, Y., Silva, L., Acosta, J.J., Ramos, F., Palomares, M.L.D. and Sobrino, I. (2017). Length-weight relationships for 22 crustaceans and cephalopods from the Gulf of Cadiz (SW Spain). *Aquatic Living Resources*. 30: 12.
- Uddin, N., Ghosh, S. and Maity, J. (2016). Length-weight relationship and condition factor of *Penaeus monodon* (Fabricius, 1798) from Digha coast, West Bengal, India. *International Journal of Fisheries and Aquatic Studies*. 4(3): 168-172.
- Ukpatu, J.E. (2021). Biometric Structure and Welfare of Tiger Shrimp (*Penaeus monodon* (Fabricius, 1798) in Okoro River Estuary, South Eastern Nigeria. *AKSU Journal of Agriculture and Food Sciences*. 5(1): 25-42.