# Potential of Vermiwash Prepared from Different Combinations of Organic Wastes to Improve the Growth, Yield and Quality of Organic Black Gram

Deepak Kumar<sup>1</sup>, S.K. Sharma<sup>2</sup>, Brijesh Kumar<sup>1</sup>, Sandeep Kumar<sup>1</sup>, Suryakanta Kashyap<sup>1</sup>, Rakesh Kumar<sup>1</sup>

### 10.18805/LR-4957

## ABSTRACT

**Background:** Excessive and unbalanced chemical fertilization in crop husbandry has attracted much attention because of soil health issues, reducing crop quality, increasing production cost and environmental risks. Organic farming is perceived as the best-known alternative method and has numerous environmental benefits. "Vermiwash" is liquid manure extracted from earthworms rich-vermicompost containing soluble nutrients, growth hormones, enzymes and many micro-organisms. As a result, the purpose of this study was to see how vermiwash affected the production and quality of black gram.

**Methods:** During the *kharif* 2018, a field experiment was set up in a randomized block design to assess the effect of nine vermiwash treatments prepared from different organic wastes like cow dung, buffalo dung, green and dry field wastes and their combinations on growth, yield attributes and yield, nutrients and protein content of organic black gram (var. PU-31) and biochemical properties of vermiwash.

**Result:** The neutral to slightly alkaline pH (7.72), E.C. (1.39 dSm<sup>-1</sup>) and maximum nutrients content, greatest average microbiological count (total number of bacteria, fungi and actinomycetes) and enzyme activity (acid phosphatase, alkaline phosphatase and dehydrogenase) were significantly found in treatment ( $T_1$ ) among the various vermiwash treatments at 35 days after collection. Moreover, the foliar application of vermiwash treatment ( $T_1$ ) significantly recorded highest growth, seed yield, nutrients and protein content. Based on-field results, it is possible to infer that treatment  $T_1$  (foliar spray in two equal split applications at a rate of 10% at the onset of flowering and 15 days following the first spray) is a better fertilizer practice for organic black gram cultivation.

Key words: Black gram, Organic, Quality, Vermiwash, Yield.

## INTRODUCTION

Black gram or Urdbean is the fourth most crucial pulse crop in India, being cultivated over about 4.6 million hectares and producing about 24.5 lakh tonnes annually, with average productivity of 533 Kg per hectare in 2020-21 (Anonymous, 2020-21). In Rajasthan, it is cultivated in 5.02 lakh hectares with 1.28 lakh tonnes of production (Anonymous, 2020-21). Being a leguminous crop, it can fix atmospheric nitrogen in its root nodules and can be used as a green manuring crop after picking the pods. Many intensive crop cycles, such as the rice-wheat cropping system, it will fit nicely due to its short duration and photo-insensitive varieties. The black gram plant possesses a deep root system that binds soil particles and thereby reduces soil erosion. The productivity of black gram is low compared to the world's average because of mainly cultivated under rainfed conditions with inadequate nutrient management on marginal lands that are unfitted for cereals/cash crops.

To improve the productivity and quality of black gram, better nutrient management practices that are economically viable and ecologically sound, such as cultivating black gram under organic conditions with potential organic sources of essential plant nutrients. Soil application of essential plant nutrients is often insufficient to meet the crop's nutrition demand, particularly in short-duration crops like black gram. <sup>1</sup>ICAR- National Dairy Research Institute, Karnal-132 001, Haryana, India.

<sup>2</sup>Rajasthan College of Agriculture, Maharana Pratap University of Agriculture and Technology, Udaipur-313 001, Rajasthan, India.

**Corresponding Author:** Deepak Kumar, ICAR-National Dairy Research Institute, Karnal-132 001, Haryana, India. Email: bhardwaj051238@gmail.com

How to cite this article: Kumar, D., Sharma, S.K., Kumar, B., Kumar, S., Kashyap, S. and Kumar, R. (2025). Potential of Vermiwash Prepared from Different Combinations of Organic Wastes to Improve the Growth, Yield and Quality of Organic Black Gram. Legume Research. 48(3): 430-436. doi: 10.18805/LR-4957. Submitted: 02-05-2022 Accepted: 22-07-2022 Online: 15-08-2022

Therefore, foliar application of organic liquid manures is the value option. Vermiwash is potential liquid manure extracted from earthworms rich vermicomposts. It contains essential plant nutrients and growth hormones like auxin and cytokinine, as well as other secretions, enzymes and a variety of beneficial microbes like heterotrophic bacteria and fungus (Gopal *et al.*, 2010). Furthermore, the composition of the raw materials used to prepare vermicompost affects the quality of vermiwash produced by earthworms (Rai and Bansiwal, 2008). Foliar spray of vermiwash significantly

increased the crops' growth, yield and quality (Nath and Singh, 2016; Senthilmuruga *et al.,* 2018; Kumar *et al.,* 2023).

Several vermiwash studies have been carried out in horticulture crops and only a few studies are available on food crops. The study's objectives were to analyze the nutrient status, beneficial microbial population and enzymatic activity of the liquid manure and evaluate the effect of liquid manure application on the crop's growth, yield and quality. The study will lend scientific credence to this low-cost approach because the materials for making liquid manure are readily available locally.

## **MATERIALS AND METHODS**

## Experiment layout and treatments details

The experiment was carried out at Organic Farm, Rajasthan College of Agriculture, Udaipur, with a latitude/longitude of 24°35'N/73°42'E at an altitude of 582.17 m above mean sea level. The mean weekly meteorological data collected at the Agro Meteorological Observatory in Udaipur during the cropping season shows in Fig 1. The soil of the experimental plot was clay loam having pH 8.1, organic carbon 0.55%, available N, P2O5 and K2O 241.2, 20.9 and 351.3 kg ha-1, respectively. The experiment was laid down in a randomized block design with three replication comprising nine treatments, *i.e.* (T<sub>1</sub>= vermiwash from 100% cow dung,  $T_2$  = Vermiwash from 100% buffalo dung,  $T_3$  = Vermiwash from 50% cow dung + 25% dry farm waste + 25% green farm waste,  $T_4$  = Vermiwash from 50% buffalo dung + 25% dry farm waste + 25% green farm waste,  $T_5$  = Vermiwash from 90% green waste incubated with 10% cow dung, T<sub>e</sub>= vermiwash from 90% green waste incubated with 10% buffalo dung, T<sub>2</sub>= Vermiwash from 90% dry waste incubated

with 10% cow dung and  $T_g$ = Vermiwash from 90% dry waste incubated with 10% buffalo dung and  $T_g$ = Control (water spray). The black gram crop (var. PU- 31) was sown in the plot (4.0 m × 3.0 m) by adopting a seed rate of 16 kg ha<sup>-1</sup> with a spacing of 30 cm × 10 cm and fertilization as a basal dose of 4 tonnes ha<sup>-1</sup> of vermicompost applied to all the treatments before 15 days of sowing and the rest of was applied through vermiwash liquid manure except control.

### Vermiwash unit

Vermicasts were prepared from different organic wastes, i.e., cow dung, buffalo dung, green and dry farm waste, using the earthen pot method. Take an earthen pot of 10liter capacity and a layer of bricks 2-3 cm was laid in the pot bottom. The pot is filled with organic wastes according to the treatments. Then, 1000 numbers (around 1 kg of earthworm's culture) of Eisenia foetida in the vermicompost pot. One more pot filled with water was hung above it so that water came out from the pot into the vermicompost pot drop by drop to keep the surface wet. The third pot was placed below it. The assembly of these three earthen pots was established in a shady place (Fig 2). After 30-35 days, the liquid vermiwash was collected in the third pot and taken to the lab for chemical analysis. The biochemical analysis of vermiwash was done as per the standard procedure shown in Table 1. Foliar spray of 35 days old vermiwash was applied into two splits- first, at the initiation of flowering and second, at the 15 days interval.

#### Microbial analysis of vermiwash

The total bacterial count (TBC), total fungal count (TFC) and total actinomycetes count (TAC) were calculated in triplicate using the standard spread-plate dilution method. T.B.C., T.F.C. and T.A.C. were calculated using nutrient



Fig 1: Weekly average of Meteorological data during experimental period during kharif, 2018-19.

Potential of Vermiwash Prepared from Different Combinations of Organic Wastes to Improve the Growth, Yield and Quality of ...

agar, potato dextrose and actinomycetes isolation agar media, respectively (Sharma *et al.*, 2021).

# Measurement of plant growth parameters, yield and quality analysis

Growth parameters *viz.* root length, nodulation, total chlorophyll, yield and yield attributes and quality content were recorded as per standard procedures/methods (Table 2). Five plants in each plot were selected randomly and labeled for recording observations in all three replications. The most extended root length from the plant's base to the root's tip was measured 45 and 60 days after



Fig 2: Vermiwash assembly.

Table	1:	Methods	used	for	vermiwash	bio-chemical	analysis
-------	----	---------	------	-----	-----------	--------------	----------

sowing and in centimeters. At the 50% flowering stage, five plants were uprooted randomly without disturbing their roots and nodules. Then dry and clean the plant's root, a number of the node were separated from the root with the help of forceps and their numbers were recorded. Yield components, namely number of pods plant<sup>-1</sup>, number of seeds pod<sup>-1</sup>, test weight and seed yield, were recorded at harvest.

## Statistical analysis

The data were subjected to statistical analysis of variance as outlined by Panse and Sukhatme (1985).

# **RESULTS AND DISCUSSION**

## Vermiwash biochemical analysis

Data indicate (Table 3) that the neutral to slightly alkaline pH (7.72), E.C. (1.39 dSm<sup>-1</sup>) and maximum content of nitrogen (0.464%), phosphorus (0.530%) and potassium (0.189%) nutrients were reported in vermiwash treatment  $T_1$  over other treatments. Moreover, high pH and E.C. were recorded in vermiwash treatment  $T_6$  and the lowest N, P and K nutrient content were found in vermiwash treatment  $T_8$ . Similar fundings were also reported by Zambare *et al.* (2008); Hatti *et al.* (2010); Gopal *et al.* (2010).

Significant differences in the microbial population and enzyme activity were observed in different vermiwash treatments (Table 4).

## **Total bacterial counts**

Total bacterial counts (T.B.C.) of various vermiwash treatments ranged from 0.54 to  $3.07 \times 10^8$  colony forming units (cfu) per milliliter of organic liquid. The highest counts

Determination	Reference/method of analysis
pH (1:10 w/v)	Garg <i>et al.</i> (2006)
Electrical conductivity	Garg <i>et al.</i> (2006)
Nitrogen	Bremmer and Mulvaney procedure (1982)
Phosphorus	Flame photometer method (Jackson, 1973)
Potassium	Flame photometer method (Bansal and Kapoor, 2000)
Phosphatase activity	Tabatabai and Bremner procedure (1969)
Dehydrogenase activity	TTC (2-3-5-Triphenyl tetrazolium chloride) reduction method (Casida et al., 1964)

Table 2: Methods adopted for plant analysis.

Determination	Methods	Reference
Total chlorophyll	Total Chlorophyll (mg g-1) = 20.2 (OD 645) + 8.02 (OD 663) $\times$ V/1000 $\times$ W	Arnon (1949)
	Where, OD = Optical density	
	V = Final vol. of 80% acetone (10 ml)	
	W = Wt. of sample taken (0.025 g)	
Nitrogen	Colorimetric method	Snell and Snell (1959)
Phosphorus	Vanadomolybdo phosphoric acid yellow colour method	Jackson (1973)
Potassium	Flame photometer method	Jackson (1973)
Protein content in seed	Nitrogen content in seed multiplied by factor 6.25	A.O.A.C. (1975)

were found in vermiwash treatment T<sub>1</sub> ( $3.07 \times 10^8$ ), while the lowest were found in treatment T<sub>8</sub> ( $0.54 \times 10^8$ ).

## **Total fungal counts**

Vermiwash treatments had total fungal counts (T.F.C.) ranging from 0.12 to  $0.79 \times 10^5$  cfu per millilitre of organic liquid. The highest counts were found in vermiwash treatment T<sub>1</sub> (0.79×10<sup>5</sup>), while the lowest concentrations were found in T<sub>7</sub> (0.12×10<sup>8</sup>).

## Total actinomycetes counts

The mean total actinomycetes counts (T.A.C.) of various vermiwash treatments ranged from 0.05-  $2.96 \times 10^5$  cfu per milliliter of organic liquid. The highest counts were observed in vermiwash treatment T<sub>1</sub> ( $2.96 \times 10^5$ ) and the lowest counts were observed in T<sub>8</sub> ( $0.05 \times 10^5$ ).

## **Enzyme activity**

The acid phosphatase activity of vermiwash treatments ranged from 2.93 to 7.55  $\mu$ g ml<sup>-1</sup>, alkaline phosphatase activity ranged from 0.74 to 2.54  $\mu$ g ml<sup>-1</sup> and dehydrogenase activity from 1.27 to 2.77  $\mu$ gml<sup>-1</sup> and these differences were statistically significant. The highest activity of these enzymes in treatment T<sub>1</sub> and the lowest was observed in treatment T<sub>8</sub>. In different vermiwash treatments, the kind of input

 Table 3: Chemical analysis of vermiwash prepared from different organic wastes.

Treatments		EC	Nitrogen	Phosphorus	Potassium
symbol	рн	(dSm <sup>-1</sup> )	(%)	(%)	(%)
T <sub>1</sub>	7.72	1.39	0.464	0.530	0.189
T <sub>2</sub>	7.79	1.47	0.445	0.522	0.178
T <sub>3</sub>	7.83	1.53	0.458	0.525	0.184
T <sub>4</sub>	7.88	1.59	0.439	0.518	0.173
T <sub>5</sub>	8.43	1.89	0.435	0.517	0.169
T <sub>6</sub>	8.46	1.92	0.430	0.511	0.167
T <sub>7</sub>	8.25	1.75	0.425	0.505	0.162
T <sub>8</sub>	8.29	1.79	0.419	0.502	0.157
S.Em±	0.009	0.011	0.001	0.001	0.001
C.D.	0.019	0.023	0.002	0.002	0.001
(P = 0.05)					

material used, resulting in diverse microbial populations and enzymatic activity that were significantly altered. Similar result was also reported by Sharma *et al.*, 2021. According to Sreenivasa *et al.* (2009), organic liquid manures developed using cow products contain many beneficial bacteria, as well as certain useful actinomycetes and fungus. The vermiwash contains an enzyme cocktail and nitrogen fixing bacteria and some phosphate solubilizing bacteria (Zambare *et al.*, 2008).

## **Growth parameters**

All the vermiwash treatments significantly influenced growth parameters (root length, nodulation and total chlorophyll content of black gram over the control. Significantly maximum root length of 17.00 and 22.00 cm was recorded at 45 and 60 days after sowing, with vermiwash treatment ( $T_1$ ), which was found to be at par with treatments  $T_3$ ,  $T_4$  and  $T_2$ , respectively and recorded an increase of 21.42 and 25.71 per cent root length at 45 and 60 days after sowing respectively, over control (Table 5). Enhanced root growth can be attributed to readily available essential plant nutrients and growth-promoting hormones in vermiwash, responsible for plants' rapid growth and development. Similar results were also obtained by Maya *et al.* (2015).

Similarly, like the growth of root length, no. of nodes per plant (65.50) at 50% flowering stage showed the maximum in number for vermiwash treatment  $(T_{\lambda})$  with an increase of 18.38 per cent over control and at par with treatment T3 while the other vermiwash treatments showed a consistent number of nodes. The vermiwash treatment (T<sub>4</sub>) showed the maximum chlorophyll content since nutrients available in vermiwash contributed to healthier plants (Table 5) and recorded an increase of 14.87 percent total chlorophyll of black gram over control. Since chlorophyll pigments synthesis in the plants is directly related to the availability of the physio-chemically active N, P, K, S and Fe nutrients. Hence, the availability of these nutrients to plants helps form chlorophyll in the leaves. Vermiwash application has been proved to promote the root formation, nodulation and chlorophyll content, as observed in many field crops (Varghese and Prabha (2014) and Senthilmuruga et al., 2018).

Table 4:	Microbial	populations	and enzyme	activities	in	vermiwash	prepared	from	different	organic	wastes
----------	-----------	-------------	------------	------------	----	-----------	----------	------	-----------	---------	--------

Treatments	Total bacterial	Total fungal	Total actinomycetes	Acid phosphatase	Alkaline phosphatase	Dehydrogenase
symbol	$count \times 10^{\scriptscriptstyle 8}$	count × 10⁵	$count \times 10^5$	(µg/ml)	(µg/ml)	(µg/ml)
T,	3.07	0.79	2.96	7.55	2.54	2.77
T <sub>2</sub>	2.77	0.45	2.76	5.63	1.95	1.87
T₃	2.96	0.75	2.86	6.45	2.11	1.96
T <sub>4</sub>	2.15	0.40	2.48	3.91	1.78	1.68
T <sub>5</sub>	1.27	0.17	0.45	3.55	1.51	1.66
T <sub>6</sub>	1.12	0.23	0.25	3.49	1.37	1.64
T <sub>7</sub>	1.03	0.12	0.08	3.34	0.77	1.31
T <sub>8</sub>	0.54	0.15	0.05	2.93	0.74	1.27
SEm±	0.043	0.032	0.046	0.097	0.037	0.040
CD (P=0.05)	0.131	0.098	0.140	0.295	0.112	0.120

## Yield and yield attributes

Foliar application of treatment ( $T_1$ ) recorded a significantly higher number of pods plant<sup>-1</sup> (24.67), seeds pod<sup>-1</sup>(6), test weight (35.88) and seed yield (800 kg ha<sup>-1</sup>), which was

 Table 5: Effect of vermiwash on root length, number of root nodules

 per plant and total chlorophyll content of organic black gram.

Treatments	Root lei	ngth (cm)	Number of root nodules	Total chlorophyll at 45 DAS
symbol	45 DAS	60 DAS	per plant at	(mg g⁻¹ fresh
			50% flowering	weight)
T <sub>1</sub>	17.0	22.0	65.50	2.78
T <sub>2</sub>	15.7	19.3	61.50	2.65
T <sub>3</sub>	16.0	19.5	62.07	2.70
<b>T</b> <sub>4</sub>	15.3	19.2	61.20	2.62
$T_5$	15.0	18.7	61.20	2.60
T <sub>6</sub>	14.7	18.5	61.17	2.60
T <sub>7</sub>	14.3	18.5	61.00	2.56
T <sub>8</sub>	14.0	18.3	60.83	2.56
T <sub>9</sub>	14.0	17.5	55.33	2.42
S.Em±	0.58	0.42	1.22	0.03
CD (P = 0.0	5) 1.8	1.27	3.65	0.10

found to be at par with the treatments  $T_3$ ,  $T_2$  and  $T_4$ , respectively (Fig 3 and 4). The higher yield and yield attributes under treatment ( $T_1$ ) may be attributed because of the positive impact of increased availability of plant nutrients, enzymes and beneficial microbes in a balanced form, which resulted in higher growth rate, chlorophyll synthesis, photosynthetic efficiency and assimilate partitioning from source to sink that increased yield attributes and help to produce healthy seed. The result was in agreement with the findings of Bhardwaj and Sharma (2016).

## Nutrients and protein content

Application of vermiwash treatment ( $T_1$ ) recorded the highest nitrogen (3.58%) and (2.55%), phosphorus (0.45%) and (0.223%) and potassium (1.30%) and (2.16%), content in seed and haulm respectively, with a significant increase of 15.48 and 62.42 per cent nitrogen, 40 and 15.47 per cent phosphorus and 9.2 and 3.8 per cent potassium in seed and haulm, respectively, over the control (Table 6). This might be because of the use of potential vermiwash, which helped in producing higher biomass and also in better recovery of N, P and K in the plant (Table 3 and 4). Similar effects have also been observed by Trivedi and Bhatt (2006).



Fig 3: Effect of vermiwash on yield attributes of organic black gram.





Potential of Vermiwash Prepared from Different Combinations of Organic Wastes to Improve the Growth, Yield and Quality of ...

Treatments	Nitrogen content (%)		Phosphorus c	ontent (%)	Potassium c	Protein content	
symbol	Seed	Haulm	Seed	Haulm	Seed	Haulm	in seed (%)
T <sub>1</sub>	3.58	2.55	0.45	0.223	1.30	2.16	22.38
T <sub>2</sub>	3.41	2.44	0.41	0.213	1.28	2.15	21.31
T <sub>3</sub>	3.46	2.53	0.43	0.216	1.30	2.15	21.65
T <sub>4</sub>	3.39	2.30	0.42	0.211	1.25	2.13	21.17
T <sub>5</sub>	3.36	2.26	0.37	0.209	1.26	2.13	20.98
T <sub>6</sub>	3.33	2.20	0.35	0.207	1.21	2.08	20.79
T <sub>7</sub>	3.34	1.78	0.36	0.208	1.20	2.12	20.88
T <sub>8</sub>	3.33	1.86	0.36	0.208	1.25	2.12	20.81
T <sub>9</sub>	3.10	1.57	0.32	0.193	1.19	2.08	19.38
S.Em±	0.029	0.115	0.009	0.004	0.009	0.009	0.179
CD (P = 0.05)	0.086	0.348	0.45	0.011	0.029	0.028	0.540

Table 6: Effect of vermiwash on nitrogen, phosphorus, potassium and protein content in seed and haulm of organic black gram.

The maximum protein content (22.38%) was recorded with vermiwash treatment T1 and a significant increase of 15.48 per cent over the control (Table 6). It can be due to enhanced absorption of nitrogen, which ultimately increases the protein content in seeds. Similarly, Hoffland *et al.* (2000) found that spraying vermiwash considerably changes tissue nitrogen content.

# CONCLUSION

Our study indicated that vermiwash is a potential source of plant nutrients, beneficial micro-organisms and enzymes. Vermiwash treatment  $(T_1)$  is the best liquid manure showing optimum pH, E.C. and highest nutrient content, microbial count and an increased number of enzymes. Foliar application of vermiwash significantly improved black gram's productivity and quality over the control (water spray). Among the vermiwash treatments, treatment  $(T_1)$  showed the highest growth, yield, nutrients and protein content in organic black gram.

## **Conflict of interest**

All authors declared that there is no conflict of interest.

## REFERENCES

- A.O.A.C. (1975). Official Method of Analysis. Edn 11, Association of Official Agricultural Chemists, Washington, DC, U.S.A. 174.
- Anonymous. (2020-21). Annual Report (2020-21) Issued by the Department of Agriculture, Cooperation and Farmers' Welfare. Ministry of Agriculture and Farmers' Welfare, Government of India, Krishi Bhawan. New Delhi-110 001. www.agricoop.nic.in
- Anonymous. (2020-21). Kharif Pulses Prospects. 2020-21. http://dpd.gov.in/Pulses-Kharif%20Propects%202020%20. pdf.
- Arnon, D.I. (1949). Copper enzymes in isolated chloroplasts, polyphenoxidase in *Beta vulgaris*. Plant Physiology. 24: 1-15.
- Bansal, S. and Kapoor, K.K. (2000). Vermicomposting of crop residues and cattle dung with *Eisenia foetida*. Bioresources Technology. 73: 95-98.

Bhardwaj, P. and Sharma, R.K. (2016). Effect of vermiwash and vermicompost on the growth and productivity of moong dal. Journal of Chemical, Biological and Physical Sciences. 6(4): 1381-1388.

- Bremmer, J.M. and Mulvaney, R.G. (1982). Nitrogen Total Method of Soil Analysis, In: [page, A.L., Millar, R.H. and Keeney, D.R. (eds.)], American Society of Agronomy, Madison. pp 575-624.
- Casida, L.E.J.R., Klein, D.A. and Santoro, T. (1964). Soil dehydrogenase activity. Soil Science. 98: 371-376.
- Chaudhari, P.S. (2005). Vermiculture and vermicomposting as biotechnology for conservation of organic waste into animal protect and organic fertilizer. Asian Journal of Microbiology Biotechnology and Environmental Science. 7: 359-370.
- Garg, V.K., Yadav, Y.K., Sheoran, A., Chand, S. and Kaushik, P. (2006). Live stocks excreta management through vermicomposting using an epigeic earthworm *Eisenia foetida*, Environmentalist. 26: 269-276.
- Gopal, M., Gupta, A., Palaniswami, R., Dhanapal, R. and Thomas, G.V. (2010). Coconut leaf vermiwash: A bio-liquid from coconut leaf vermicompost for improving the crop production capacities of soil. Current Science. 98(9): 1202-1210.
- Hatti, S.S., Londonkar, R.L., Patil, S.B., Gangawane, A.K. and Patil, C.S. (2010). Effect of *Eisenia foetida* vermiwash on the growth of plants. Journal of Crop Science. 1(1):6-10.
- Hoffland, E., Jeger, M.J. and Van Beusichem, M.L. (2000). Effect of nitrogen supply on disease resistance in tomato depends on the pathogen. Plant and Soil. 218: 239-247.
- Jackson, M.L. (1973). Soil: Chemical Analysis, Prentice Hall Inc. Engle Cliffs, New Jersey.
- Kumar, D., Massey, J.X., Sharma, S.K., Mundra, S.L. and Yadav, S.K. (2023). Vermiwash prepared from different combinations of organic sources to improve growth and yield of black gram [*Vigna mungo* (L.) hepper] for organic agriculture. Indian Journal of Agricultural Research. 57(5): 630-634. doi: 10.18805/IJARe.A-5630.
- Maya, M.J. and Satish, A.B. (2015). Influence of vermiwash on germination and growth parameters of seedlings of greengram (*Vigna radiata* L.) and blackgram (*Vigna mungo*). International Journal of Current Microbiology and Applied Sciences. 4(9): 635-643.

- Nath, G. and Singh, K. (2016). Vermiwash: A potent liquid biofertilizer. Research Journal of Science and Technology. 8(1): 21-30.
- Panse, V.G. and Sukhatme, V.P. (1985). Statistical Methods for Agricultural Workers. Indian Council of Agricultural Research. New Delhi.
- Rai, N. and Bansiwal. (2008). Vermiwash: An excellent source of nutrition for plant growth. Electronic Journal of Environmental Sciences. 1: 19-21.
- Senthilmurugan, S., Sattanathan, G., Vijayan, P., Pugazhendy, K. and Tamizhazhagan, V. (2018). Evaluation of different concentration of vermiwash on seed germination and biochemical response in okra (*Abelmoschus esculentus* L.). International Journal of Biology Research. 3(1): 228-231.
- Sharma, S.K., Jain, D., Choudhary, R., Jat, G., Jain, P., Bhojiya, A.A., Jain, R. and Yadav, S.K. (2021). Microbiological and enzymatic properties of diverse *Jaivik Krishi* inputs used in organic farming. Indian Journal of Traditional Knowledge. 20(1): 237-243.

- Snell, F.D. and Snell, C.P. (1959). Colorimetric Methods of Analysis 3<sup>rd</sup> Ed. Vol. 2<sup>nd</sup>. Van Nostrand Inc. New York. Soil Science. 50: 225-226.
- Sreenivasa, M.N., Nagaraj, M., Naik and Bhat, S.N. (2009). Beneficial Traits of Microbial Isolates of Organic Liquid Manures, In: First Asian PGPR Congress for Sustainable Agriculture, A.N.G.R.A.U., Hyderabad, India.
- Tabatabai, M.A. and Bremner, J.M. (1969). Use of p-nitrophenyl phosphate for assay of soil phosphatase activity. Soil Biology and Biochemistry. 1: 301-307.
- Trivedi, R. and Bhatt, S.A. (2006). Phosphate activity in semiarid soil of pathan. Asian Journal of Microbiology, Biotechnology and Environmental Science. 8: 303-305.
- Varghese, S.M. and Prabha, M.L. (2014). Biochemical characterization of vermiwash and its effect on growth of *Capsicum frutescens*. Malaya Journal of Biosciences. 1(2): 86-91.
- Zambare, V.P., Padul, M.V., Yadav, A.A. and Shete, T.B. (2008). Vermiwash: Biochemical and microbiological approach as ecofriendly soil conditioner. Journal of Agriculture and Biology Science. 3(4): 1-5.