



Effect dates of sowing and cutting time on seed yield and quality of alfalfa (*Medicago sativa* L.) CV. RL-88

P.R. Arvind Kumar*, B.C. Channakeshava and R. Siddaraju

University of Agriculture Sciences, College of Agriculture,
Gandhi Krishi Vignana Kendra, Bengaluru-560 065, Karnataka, India.

Received: 29-05-2017

Accepted: 04-07-2017

DOI:10.18805/IJARE.A-4804

ABSTRACT

An experiment was carried to find out the effect of sowing dates and cutting time on seed yield and quality of alfalfa cv. RL-88. The results of the experiment revealed that, the significantly higher seed yield per plant (1.491 g), seed yield per plot (161.07 g) and seed yield (355.00 kg ha⁻¹) was noticed in 15th August sowing and the cutting carried out at 60 days after sowing. Whereas, the seed quality parameters of the resultant seeds viz., highest seed germination (95.00 %), seedling length (13.90 cm), seedling dry weight (412.69 mg) and seedling vigour index (1320) found highest in 15th August sowing and the cutting followed at 60 days after sowing.

Key words: Alfalfa, Cutting time, Seed germination, Seedling vigour index.

INTRODUCTION

Alfalfa (*Medicago sativa* L.) is popularly known as lucerne and rightly called as “Queen of Forage” in India. It has been under cultivation under wide range of climatic and soil conditions throughout the world since ages. In Karnataka it is popularly called as Kudre Masale/ Kudare Menthe Seed is the most critical and important input to enhance the production potential of all agricultural crops, including fodder. The efficacy of other inputs is largely dependent on availability and timely sowing of quality seeds of improved genetics. Therefore, an assured supply of fodder seeds of improved varieties or hybrids to farmers at a reasonable price is crucial for enhancing fodder production. Production of quality seeds is one of the major hinderances in alfalfa due to many reasons viz., a) Physiological reasons: very poor seed-setting and low seed production ability of most of the varieties, many of forage species are of indeterminate growth habit, poor translocation mechanism of photosynthates from vegetative to reproductive structures, Non-synchronization of flowering, prolonged flower drop and uneven maturity, improper management of monetary and non-monetary inputs especially in forage crops, Preferential use of low capability marginal lands and rainfed conditions for raising most of the forage seed crops and low insect activity during hot summer months results in low seed productivity in entamophilous allogamous legume species. b) Others, blank seeds, seed dormancy, apomixes, seed shedding and harvesting (Channakeshava, 2014).

Tripping is a elite phenomenon observed in alfalfa which is the release of sexual column from the keel petal when the bees land on flower and probe for nectar. Upon

release, the column strikes the standard petal, sometimes striking the underside of the head of the bees first, at times with such a force that the bee can extricate its head. When the flower is tripped the pollen dust would spread upon the bee and is then carried to another flower. At the same time, pollen brought from another flower is accidentally rubbed upon the stigma and resulted in cross pollination as prescribed by Henslow (1867). The high plant density of flowers influence the activity of wild pollinators, particularly bumble bees and large solitary bees that are physically capable of tripping the flower. Among the different pollinators recommended in forage crops, alfalfa leaf cutter bee is world's most effectively used and intensely managed solitary bee because of their gregarious nature, presence of trichomes at lateral surface of the abdomen which facilitate the pollen dispersal and in turn results in effective pollination (Theresa *et al.*, 2011). Alfalfa has good production potential, but lack of suitable agro techniques (*i.e.* seed rate, time of sowing, cutting intervals and fertility level, different insect pollinator for tripping etc.) is responsible for reduction of quantity and quality of forage yield. In this regard a study to standardize the sowing time and cutting time on forage and seed yield of alfalfa was carried out. The time of first cut after sowing is important to obtain maximum number of cuts as well as green forage yield at each cut. Thus, cutting management not only provides information about the regeneration potential of the crop but also growth peak and yield too. Moreover, the cutting management may be responsible for quality and quantity for forage crops and particularly for alfalfa forage yield.

*Corresponding author's e-mail: arvindrathod09@gmail.com

MATERIALS AND METHODS

An experiment on effect of date of sowing and cutting time on growth, forage and seed yield of alfalfa (*Medicago sativa* L.) cv. RL-88 under Eastern Dry Zone of Bengaluru was carried out at Seed Science and Technology, Experimental E-block, GKVK, University of Agricultural Sciences, Bengaluru during 2015-16. Field experiment was laid out in split plot design with three replication with gross plot size: 2.1 m x 2.0 m=4.20 m² with spacing of 30 x 10 cm with treatment comprises of four different dates of sowing D₁: 1st July, D₂: 15th July, D₃: 1st August and D₄: 15th August with four cutting time viz., C₀: no cut, C₁: First cut @ 60 DAS, C₂: Second cut @ 85 DAS and C₃: Third cut @ 110 DAS carried out at different stage of crop growth.

The moisture content of seed was determined by the oven dry method Anonymous (2014). After drawing samples from different treatments, they were subjected to moisture determination immediately. At each stage of harvest, 10 pods were harvested and threshed separately. The seeds were dried at 103 °C for 17 hours and then samples were taken out from the drying chamber, cooled to room temperature and their dry weight was recorded. From the primary data, moisture percentage in the seed was computed as detailed below

$$\text{Seed moisture (\%)} = \frac{\text{Fresh weight of the seed} - \text{weight of the seed}}{\text{Fresh weight of the seed}} \times 100$$

The pods from each treatment after harvest were hand shelled and seeds obtained were dried in the morning hours of sunlight, to bring down the initial seed moisture content to around 9 per cent. These seeds were subjected to laboratory observations.

The germination test was conducted in the laboratory using between paper method Anonymous (2014). One hundred seeds each of four replications were placed on germination paper and rolled towels were incubated in germination chamber maintained at 25°C and 90 per cent relative humidity. The germinated seedlings were evaluated on 5th and 14th day as first and final count and percentage germination was expressed based on normal seedlings.

The seedling vigour index was calculated as per the formula given by Abdul Baki and Anderson (1973).
Seedling vigour index = Germination (%) x Mean seedling length (cm)

RESULTS AND DISCUSSION

The proper time for sowing determines forage yield of alfalfa crop. The optimum time of sowing of alfalfa depends upon the nature of variety and the temperature. Now-a-days, the high yielding varieties are most sensitive to time of sowing, hence, optimum time of sowing contributes more towards yield.

Effect of sowing dates and cutting time on seed yield of alfalfa cv. RL-88:

The seed yield of alfalfa was significantly influenced by dates of sowing (Table.1). The highest seed yield (322.00 kg ha⁻¹) was noticed in D₄: 15th August sown crop which was about 30.90 %, 26.46 % and 12.86 % higher than D₁: 1st July (222.52 kg), D₂: 2nd July (236.79 kg) and D₃: 1st August (280.59 kg), respectively. The higher seed yield obtained in D₄: 15th August might be due to favorable temperature during crop growth period resulting in increased number of pods per plant and seeds per pod and higher growth attributes, which may be responsible for better source sink relationship between environment and fertilization of lucerne. They indicated that numbers of filled pods/plant were positively correlated with the pollen tube length. Under the delayed sowing the crop was exposed to comparatively higher temperature (36 °C to 40 °C) during flowering period as compared to earlier sowing time. This might have increased the pollen tube length and “**autotripping of flowers**” consequently led to increase in all the yield attributes. Hence, the temperature prevailing during flowering period in terms of longer day length and better

Table 1: Effect of dates of sowing and cutting time on seed yield per plant, seeds yield per net plot and seed yield per hectare of alfalfa (*Medicago sativa* L.) Cv. RL-88

Treatments	Seed yield (g plant ⁻¹)	Seeds yield (g net plot ⁻¹)	Seed yield (kg ha ⁻¹)
D ₁ : 1 st July	0.953	102.35	222.52
D ₂ : 15 th July	0.997	107.44	236.79
D ₃ : 1 st August	1.185	127.32	280.59
D ₄ : 15 th August	1.351	146.13	322.00
S. Em±	0.004	0.02	2.07
CD (P=0.05)	0.014	0.06	7.18
C ₀ : No cut	0.971	104.54	230.44
C ₁ : First cut @ 60 DAS	1.265	136.40	297.51
C ₂ : Second cut @ 85 DAS	1.168	126.15	278.02
C ₃ : Third cut @ 110 DAS	1.082	116.16	255.93
S. Em±	0.003	0.02	1.86
CD (P=0.05)	0.010	0.05	5.42
D ₁ C ₀	0.717	77.08	170.08
D ₁ C ₁	1.137	122.87	258.39
D ₁ C ₂	1.009	108.87	239.94
D ₁ C ₃	0.950	100.59	221.67
D ₂ C ₀	0.845	90.22	198.83
D ₂ C ₁	1.041	112.47	247.86
D ₂ C ₂	1.061	114.63	252.62
D ₂ C ₃	1.041	112.47	247.86
D ₃ C ₀	1.041	112.47	247.86
D ₃ C ₁	1.391	149.19	328.81
D ₃ C ₂	1.231	132.99	293.09
D ₃ C ₃	1.078	114.63	252.62
D ₄ C ₀	1.281	138.39	305.00
D ₄ C ₁	1.491	161.07	355.00
D ₄ C ₂	1.371	148.11	326.43
D ₄ C ₃	1.261	136.97	301.59
S. Em±	0.007	0.03	3.71
CD (P=0.05)	0.020	0.10	10.84
CV	18.003	18.23	19.29

sun light under the treatments of later sowing might be optimum for better fertilization and seed setting diversification of more plant energy to the development of reproductive organ. These results were conformity with those results obtained by Kabir *et al.* (2009) in chickpea; Gawariya *et al.* (2015) in forage mustard.

Similarly, the study revealed the cutting time also exhibited statistically significant effect on various seed yield attributing and quality parameters.

The initial cutting at 60 days after sowing and then left for seed production recorded significantly higher seed yield (297.51 kg ha⁻¹) followed by C₀: no cut (230.44 kg ha⁻¹), C₂: cut taken at 85 (278.02 kg ha⁻¹) and C₃:110 DAS (255.93 kg ha⁻¹), respectively. Cutting operation carried out at initial stage of crop i.e., 60 days after sowing recorded better reactions. It is also obvious that cutting time at early growth stage might have disturbed the normal growth of cut plants causing slower re-growth, greater number of fertile tillers, as well the seed yield per plant, which might have ultimately resulted higher yield per hectares. These findings line in conformity with Bhatt *et al.* (2009) in forage grasses; Hadi *et al.* (2012) in barley.

The interaction effect between sowing dates and cutting time (D x C) revealed the marked variations on seed yield attributing parameters.

The highest seed yield (355.00 kg ha⁻¹) noticed in crop sown on 15th August with cutting imposed at 60 DAS followed by D₃C₁: 1st August sown crop with cutting interval imposed at 60 DAS (328.81 kg ha⁻¹), D₄C₂: 15th August sown crop with cutting interval imposed at 85 DAS (326.43 kg ha⁻¹), D₄C₀: 15th August sown crop with no cut (305.00 kg ha⁻¹) and D₁C₀: 1st July sown crop with no cut (170.08 kg ha⁻¹) as control.

The seed yield per plant (1.49 g) and seed yield per net plot (161.07 g) at harvest was recorded in 15th August sown crop with cutting imposed at 60 DAS. The consistent increase in seed yield components seen in D₄C₁ interaction may be attributed due to the synergetic effect of early sowing in June month and early crop growth cutting help in robust crown development and production of maximum number of seed yield attributing parameters in nut shell. These results are in conformity with the findings of Suneetha Devi and Satyanarayana Rao (2007) in forage cow pea and Asaad *et al.* (2014) in alfalfa cv. Bam.

Table 2: Effect of dates of sowing and cutting time on seed moisture content, seed germination, root length and shoot length of alfalfa (*Medicago sativa* L.) cv. RL-88

Treatments	Seed moisture content (%)	Seed germination (%)	Root length (cm)	Shoot length (cm)
D ₁ : 1 st July	7.71	89.00	5.50	6.53
D ₂ : 15 th July	7.61	90.42	5.68	6.78
D ₃ : 1 st August	7.44	91.75	5.90	6.97
D ₄ : 15 th August	7.38	93.33	6.18	7.13
S. Em±	0.05	0.65	0.01	0.03
CD (P=0.01)	0.16	2.25	0.03	0.12
C ₀ : No cut	7.86	89.83	5.65	6.54
C ₁ : First cut @ 60 DAS	7.27	92.92	6.15	7.14
C ₂ : Second cut @ 85 DAS	7.46	91.17	5.88	7.02
C ₃ : Third cut @ 110 DAS	7.56	90.58	5.58	6.72
S.Em±	0.04	0.37	0.01	0.02
CD (P=0.01)	0.12	1.09	0.03	0.05
D ₁ C ₀	8.02	87.00	5.40	6.03
D ₁ C ₁	7.48	90.67	5.80	6.93
D ₁ C ₂	7.67	89.33	5.50	6.70
D ₁ C ₃	7.69	89.00	5.30	6.47
D ₂ C ₀	8.00	89.00	5.50	6.53
D ₂ C ₁	7.35	92.67	6.10	7.10
D ₂ C ₂	7.46	90.33	5.70	6.93
D ₂ C ₃	7.65	89.67	5.40	6.53
D ₃ C ₀	7.61	90.67	5.60	6.73
D ₃ C ₁	7.25	93.33	6.20	7.12
D ₃ C ₂	7.37	92.00	6.10	7.20
D ₃ C ₃	7.55	91.00	5.70	6.83
D ₄ C ₀	7.83	92.67	6.10	6.87
D ₄ C ₁	6.99	95.00	6.50	7.40
D ₄ C ₂	7.33	93.00	6.20	7.23
D ₄ C ₃	7.35	92.67	5.93	7.03
S. Em±	0.08	0.75	0.01	0.04
CD (P=0.01)	0.25	2.18	0.03	0.10
CV (%)	4.64	2.65	7.36	8.92

Effect of sowing dates and cutting time on seed quality of alfalfa cv. RL-88: Significant variations in seed quality parameters were noticed due to different sowing dates (Table. 2 and 3). The crop sown during August 15th (D₄) recorded highest seed germination (93.33 %), mean seedling length (13.31 cm), seedling dry weight (410.726 mg) and seedling vigour index (1242) and lowest seed moisture content (7.38 %) as compared to other dates of sowing. This could be due to better source to sink relationship resulting better accumulation and assimilation of photosynthates. The early sown crop during 1st July gave lowest quality this could be due to stress conditions existed during growth, flowering (low abundance of pollinator activity in early sown crop, tripping per cent). The impact of flower abundance and pollinator movement on seed yield and quality is of economic importance. The hypothesis says that seed set is lower when standing crop of open flowers is high than when standing crop is low. This could occur if pollinators move more among flowers on the same plant, causing self-pollination, when flowers are abundant (Stanisavljevic *et al.*, 2012).

Similarly the cutting time imposed found significant effect on seed quality attributes. Among the three cutting treatments, C₁: cutting imposed at 60 DAS gave highest seed germination (92.92 %), mean seedling length (13.29 cm), seedling dry weight (411.308 mg), seedling vigour index (1235), and lowest seed moisture content (7.27 %), compared to other cutting time.

The interaction effect of sowing dates and cutting time shown marked variation in seed quality parameters of alfalfa.

The highest seed moisture content (8.02 %) was recorded in D₁C₀ (1st July sown crop with no cut) and the lowest (6.99 %) was recorded in D₄C₁ (15th August sown crop + cutting interval @ 60 DAS). Similarly, the higher seed germination (95.00 %), mean seedling length (11.90 cm), seedling dry weight (412.690 mg) and seedling vigour index (1320) recorded in 15th August sown crop + cutting interval @ 60 DAS was found minimum in D₁C₀: 1st July sown crop with no cut (87.00 %, 11.43 cm, 407.58 mg and 997, respectively). These findings were in conformity with Rade *et al.* (2012) and Azizza and Babo (2013) in alfalfa.

CONCLUSION

The non-availability of good quality seed is one of the major constraints in popularizing the seed production practice for most of the forage crop species. Hence, the

REFERENCES

- Abdul-Baki, A.A. and Anderson, J.D. (1973). Vigour determination of soybean seeds by multiple criteria. *Crop Sci.*, **13**:630-633.
- Anonymous, (2014). International Rules for Seed Testing by International Seed Testing Association. *Seed Sci. & Tech.*, **27**. 215.
- Asaadi, M., Bahareh D. and Bahman P. (2014). Determination the best time to harvest and seed rates in order to increase the yield production of alfalfa (*Medicago sativa* L.) seed. *Intl. J. Bio-sci.*, **5** (5): 100-106.
- Azizza, M. and Babo F. (2013). Effect of stage of cutting alfalfa on crude protein content and dry matter yield. *ARPJ. Sci. & Technol.*, **3** (10): 1-4.

Table 3: Effect of dates of sowing and cutting time on seedling length, seedling dry weight and seedling vigour index of alfalfa (*Medicago sativa* L.) cv. RL-88

Treatments	Seedling length (cm)	Seedling dry weight (mg)	Seedling vigour index
Dates of sowings (D)			
D ₁ : 1 st July	12.03	408.58	1073
D ₂ : 15 th July	12.45	409.59	1127
D ₃ : 1 st August	12.87	410.66	1181
D ₄ : 15 th August	13.31	410.73	1242
S. Em±	0.03	0.09	10
CD (P=0.01)	0.12	0.31	36
Cutting time (C)			
C ₀ : No cut	12.19	408.96	1098
C ₁ : First cut @ 60 DAS	13.29	411.31	1235
C ₂ : Second cut @ 85 DAS	12.89	409.87	1176
C ₃ : Third cut @ 110 DAS	12.29	409.41	1114
S. Em±	0.02	0.02	5
CD (P=0.01)	0.05	0.06	15
D ₁ C ₀	11.43	407.58	997
D ₁ C ₁	12.73	410.33	1155
D ₁ C ₂	12.20	408.28	1091
D ₁ C ₃	11.77	408.16	1048
D ₂ C ₀	12.04	408.02	1073
D ₂ C ₁	13.20	410.96	1224
D ₂ C ₂	12.63	409.88	1141
D ₂ C ₃	11.93	409.50	1070
D ₃ C ₀	12.33	410.07	1119
D ₃ C ₁	13.32	411.26	1243
D ₃ C ₂	13.30	410.79	1223
D ₃ C ₃	12.53	410.51	1140
D ₄ C ₀	12.97	410.19	1202
D ₄ C ₁	13.90	412.69	1320
D ₄ C ₂	13.43	410.54	1249
D ₄ C ₃	12.93	409.48	1198
S. Em±	0.04	0.04	10
CD (P=0.01)	0.10	0.13	29
CV (%)	7.87	0.33	8

production and supply of high quality seeds of these crops at reasonable price is very much necessary. The experiment results revealed that, sowing crop during 15th August sowing with cut at 60 days after sowing resulted in significant increase in seed yield and quality of seeds of alfalfa.

ACKNOWLEDGEMENT

The authors would like to thank Indian Grassland and Fodder Research Institute (IGFRI), Southern Regional Research Station, Dharwad (Karnataka) for providing certified seeds of alfalfa variety RL-88 for field experiments.

- Bhatt, R. K., Tripathi and Tiwari, (2009). Effect of cutting management on seed yield and its quality in grasses. *Seed Res.*, **37** (1&2): 26-29.
- Channakeshava, B. C.(2014). Problems and prospective of forage seed production in India. (*Edt.*, Quality Seed Production : Vasudevan *et al.*,) *Agro. Tech. Publ. Aca.*, Udaipur: pp. 265.
- Gawariya, S. C., Chopra, N. K., Chopra, N. and Harika, A. S.(2015). Effect of date of sowing and crop geometry on growth and yield parameters of forage mustard (Var. *Chinese Cabbage*). *African J. Agric. Res.*, **10**(33): 3292-3295.
- Hadi, F., Hussain, F. and Arif, M. (2012). Effect of different nitrogen levels and cutting on growth behavior of dual purpose barley (*Hordeum vulgare* L.). *Scholarly J. Agril. Sci.*, **2**(10):263-268.
- Henslow, C. (1867). Note on the structure of *Medicago sativa*, as apparently affording facilities for intercrossing of distinct flowers. *J. Linn. Soc.*, **9**:327-329.
- Kabir Fahm, Bari M. N., Karim M. A., Khaliq Q. A. and Ahmed J. U. (2009). Effect of sowing time and cultivars on the growth and yield of chickpea under rainfed condition. *Bangladesh J. Agric. Res.*, **34**(2):335-342.
- Rade S., Dragoljub, B., Dragan, D., Vladeta Stevovi., Dragan Terzic, Jasmina Milenkovi and Dragoslav Djokic, (2012). Influence of plant density on yield components, yield and quality of seed and forage yields of alfalfa varieties. *Romanian Agric. Res.*, **29**:245-254.
- Stanisavljevic, R., Bekovic, D., Djukic, D., Stevovic, V., Terzic, D., Milenkovic, J. and Djokic, D. (2012). Influence of plant density on yield components, yield and quality of seed and forage yields of alfalfa varieties. *Romanian Agril. Res.*, **29**: 245-253.
- Suneetha Devi, K. B. and Satyanarayana RAO V. (2007). Effect of spacing and varieties on seed yield of forage cowpea. *Forage Res.*, **33** (3) :161-163.
- Theresa, L., Pitts-Singer. and James, H. C. (2011). The alfalfa leaf cutting bee (*Megachile rotundata*). The World's most intensively managed solitary bee. *Ann. Rev. Entomol.*, **56**: 221-37.