



# Study of Diversity and Abundance Pattern of Natural Enemies Associated with the Mango Mealy Bug (*Drosicha mangiferae* G) at Malda of West Bengal, India

Subhamoy Roy<sup>1</sup>, Prasanta Saha<sup>2</sup>, Devanshu Gupta<sup>3</sup>,  
Kaushik Chakraborty<sup>1</sup>, Partha Sarathi Nandi<sup>1</sup>

10.18805/IJArE.A-6129

## ABSTRACT

**Background:** As a pest, the mango mealy bug (*Drosicha mangiferae* G) is very important as it ranks second in terms of damage inflicted to the fruit crop. Natural enemies such as predators, parasitoids are considered to be significant deterrent of mango mealy bug. These natural enemies can act as biological control agent which can bring about pest suppression as an important tool of integrated pest management. Therefore, studies of natural enemy complex of mango mealy bug become pertinent. In the current study the available predators, parasitoids of mango mealy bug was surveyed at different regions of Malda.

**Methods:** The predators of mango mealy bugs were studied by sampling mealy bug infested colonies present in the inflorescence and in 20 cm<sup>2</sup> area of trunk. Sampling was carried out by visualization of predatory behavior of the entomophagous insects. Parasitoid was studied by observing the emergence of adults from infested mealy bugs in a closed container.

**Result:** Ladybird beetles and Green lacewings were the most abundant predators; their relative abundance was 48.96% and 37.30%. Among the coccinellids, *Rodalia amabilis* and one unidentified coccinellid were found as dominant. The green lacewing larva, *Chrysopa* sp. acted as important predator. Besides Ladybird beetles and Green lacewings, the spiders and bugs were also noticed in the predatory guild. Only one dipteran parasitoid species, *Cryptochetum* sp. nr. *iceryae* (Williston) was found to parasitize the mealy bugs. The standardization of mass rearing techniques for the three predators and one parasitoid and their release in the mealy bug infested orchards can bring about efficacious control of this menacing pest.

**Key words:** Coccinellids, *Cryptochetum* sp. nr. *iceryae* (Williston), Green lacewing, Mango Mealybug, Parasitoids.

## INTRODUCTION

Mango (*Mangifera indica* L.) is deliberated to be the 'King of fruits', because the presence of its unique sweet taste and nutritional as well as medicinal values (Litz, 1997). On the basis of the production data, in India, the total production of mango is around 3.95 million tons, which is about 57.18% of the world fabrication (Sahoo and Jha, 2009). The Gangetic plain of West Bengal contributes immensely to the total production of West Bengal State with a variety of mango cultivars (Jha *et al.*, 2006). In terms of production Malda tops the list amongst different districts of West Bengal with an annual production of about 292.28 metric tons (Anonymous, 2018).

Besides the rampantly changing climatic conditions during the mango growing time, the insect pests inflict major damage to mango production (Ishaq *et al.*, 2004). There are about 260 insect species that wreak havoc on mango orchards (Pena *et al.*, 1998). If the insect pests of major importance is considered then mango mealy bug, *Drosicha mangiferae* (G.) is hugely important for the mango growers as it causes quantum fruit drop and loss (Karar, 2010). The coating of wax all over the body of mango mealy bug (MMB) makes skin impervious to water so it is very difficult to manage this insect pest with contact pesticide application. Farmers adopt array of different combinations of practices like cultural, mechanical, biological and chemical methods

<sup>1</sup>Laboratory of Insect Ecology and Pest Management, Department of Zoology, Raiganj University, Raiganj-733 134, West Bengal, India.

<sup>2</sup>Department of Zoology, Krishnath College, Berhampore, Murshidabad-742 101, West Bengal, India.

<sup>3</sup>Zoological Survey of India, Coleoptera Section, New Alipore, Kolkata-700 053, West Bengal, India.

**Corresponding Author:** Partha Sarathi Nandi, Laboratory of Insect Ecology and Pest Management, Department of Zoology, Raiganj University, Raiganj-733 134, West Bengal, India.  
Email: partharayma99@gmail.com

**How to cite this article:** Roy, S., Saha, P., Gupta, D., Chakraborty, K., Nandi, P.S. (2023). Study of Diversity Aand Abundance Pattern of Natural Enemies Associated with Tthe Mango Mealy Bug (*Drosicha mngiferae* G) at Malda of West Bengal, India. Indian Journal of Agricultural Research. DOI: 10.18805/IJArE.A-6129.

**Submitted:** 22-06-2023 **Accepted:** 06-10-2023 **Online:** 06-11-2023

of control to avoid the yield loss. Insecticides, though used widely, have environmental constrains. Therefore, natural enemies such as predators and parasitoids are very important for lowering the pest and for the reduction of pesticide application. Predatory coccinellids and Chalcid parasitoid wasps, encyrtids (Hymenoptera, Encyrtidae) act as the most important natural enemies for mango pests (Iperti, 1999). Mango mealy bugs in India are known to be

predated by four different genera of lady bird beetles belonging to Chilocorinae (Sub-Family) and six genera of Scymninae (Sub-Family); the Chilocorinae beetles are *Brumus*, *Stictobura*, *Orcus* and *Aspidimerus* and *Cryptolaemus*, *Nephus*, *Diomus*, *Sidis*, *Parasidis* and *Pseudoscymnus* belonging Scymninae subfamily also prey preferentially on mealy bugs population (Ipert, 1999). Beside Coccinellidae, brown lacewings (Neuroptera; Hemerobiidae) and predatory gall midges (Diptera; Cecidomyiidae) are important predator of scale insects (Tandon and Lal, 1978). A wide array of predators and parasitoids has been found to be present and they play important role to control the mango mealy bug populations in mango orchards (Jakhar *et al.*, 2016; Hayat, 2009). Nevertheless, there is no published literature on predator abundance or diversity against mango mealy bug of the species (*D. mangiferae*) which is one of the important pests of mango.

Hymenopteran and Dipteran parasitoids are considered to be very effective against scale insects and as found elsewhere (Larson, 2013). But work on dipteran parasitoid of mango mealy bug has not been reported from India till date. Information pertaining to parasitoid species of a pest is very important to formulate biological control strategies. In the backdrop of this context the current study was undertaken to measure the abundance and diversity of predators and parasitoids associated with mango mealy bug pest at Malda region of West Bengal.

## MATERIALS AND METHODS

### Site of study

Seasonal periodicity, incidence and abundance of mango mealy bug (MMB), *D. mangiferae* and occurrence of its natural enemies in the form of predators and parasitoids was noted from selected villages of the Malda District of West Bengal during 2020 and 2021. The villages had mostly langra, fazli, lakhanbhog, gopalbhog and himsagar cultivars. The villages were Manikchak (25°03'45"N-87°54'33"E), Ratua (25°20'01"N-87°92'74"E) and Chanchal (25°39'N-87°99'E) of the District Malda, West Bengal.

### In inflorescence

Ten mealy bug infested inflorescence from each of the two selected orchards from each village were primarily taken in to the laboratory after wrapping it in a plastic bag with proper aeration. For observation and identification of predators these MMB infested inflorescence were rinsed in 70% alcohol to kill and preserve predators. Side by side, the large and medium sized agile predators were captured on observational day using an aspirator and also preserved in 70% alcohol.

### In the tree trunk

Presence of natural enemy in 20 cm<sup>2</sup> area of tree trunk was located and visually counted from randomly chosen three trees from two different mango orchards. Individual sample of predator was kept as type specimen and also for further taxonomic identification.

### Study on the incidence of MMB parasitoids

During March-April 2018, 100 randomly collected mealy bugs from two different orchards were kept in a plastic jar, covered with organza material that was tied with rubber band. 50% honey soaked cotton was kept over the organza to ensure as food source for the emerging parasitoid, if any, that came out of the mealy bug. Fresh mango leaves as food was given to the mealy bugs. The leaf-petioles were tied with water soaked cotton to keep the moisture level. After 20-25 days flies emerged from mealy bugs. Fly was etherized and accordingly counted and the extent (%) of parasitization was further assessed (Chrysantus, 2012). The study was carried out at laboratory of insect ecology and pest management, Department of Zoology, Raiganj University.

### Taxonomic identification

- The adults of Coccinellids were identified using one stereo zoom microscope and available standard key (Jouveau *et al.*, 2018 and Angelfire, 2022).
- The larvae of lady beetles were identified after consulting available literature. (Bugguide, 2022).
- Spiders were identified using available pictorial keys (Yadav *et al.*, 2004).

Some important predators found during the study were identified by Scientist C of ZSI, Kolkata. Parasitoid was identified by Principal Scientist, National Research Centre for Banana (NRCB), Trichyrapally, Tamil Nadu, 620102.

## RESULTS AND DISCUSSION

### Study of the existing predators in the mango orchards

During 2020 and 2021 at specific mango orchards, grossly 20 species of predators are noticed. Ladybird beetles (Family: Coccinellidae), Green lacewings (Family: Chrysopidae), spiders (Family: Oxyopidae, Family: Lycosidae, Family: Salticidae and Family: Tetragnathidae), green stinkbug (Family: Pentatomidae, *Chivaria* sp.), brown marmorated stink bug (Family: Pentatomidae, *Halyomorpha* sp.) and Mediterranean red bug (Family: Pyrrhocoridae, *Scantius aegypticus*) were identified to feed on mealybug colony.

Ladybird beetles (Family: Coccinellidae) such as *Rodolia amabilis* (Kapur), *Harmonia axyridis* (Pallas), *Chelomenes sexmaculatus* Fabricius, *Micraspis discolor* Fabricius, *Coccinella transversalis* Fabricius, *Propylaea japonica* Thunberg, *Scymnus (pullus) posticalis* Motschulsky were found. Among the spiders, *Oxyopes tridens* Brady, *Oxyopes javonicus* Thorell, *Plexxipuspa kulli* Audouin, *Hyllus semicupreus* Simon, *Pardosa agricola* Thorell, *Leucauge decorata* Walckenaer are identified.

Apart from, green stinkbug (*Chivaria* sp.), brown marmorated stink bug (*Halyomorpha* sp.) and Pyrrhocorid bug (*Scantius* sp.) are identified. Considerable variation of the natural enemies is observed between the selected orchards in both insecticide treated and untreated fields.

*Rodalia amabilis* and Chrysopidae larva was found to be the most abundant followed by one unidentified Coccinellidae A larva and others (Fig 1, 2, and 3) (Table 1, 2).

#### Parasitoid in mango orchards

In the duration of the study only one species of parasitoid *Cryptochaetum* sp. nr. *iceryae* (Williston), had found to have emerged from mango mealy bug when reared in the laboratory for at least 3 weeks. The emerged parasitoid was identified as a dipteran fly, *Cryptochaetum* sp. nr. *iceryae* (Williston) (Fig 4) (NBAIR, 2022).

The identifying characteristics feature of the *Cryptochaetum* sp. nr. *iceryae* (Williston):

- i. Small sized body having dark bluish body color.
- ii. Length of the body 2-2.5 mm. with dark radish eyes.
- iii. Antenna with three segments along with terminal spine (small and heavy).
- iv) At the terminal segment of the antenna extended in the region of to inferior margin of the eyes.

#### Abundance and diversity of predators in three different places of study area at Malda, West Bengal

The numerical abundance is expressed as the exact cumulative number of individual predators found during the total period of study at three different areas i.e. at Manikchak, Ratua and Chanchal of Malda. The numerical abundance,





Fig 1: Coccinellidae predators of the MMB in mango orchards,at Malda, West Bengal.




diversity indices, as well as relative abundance of those predator species were calculated and depicted here. For counting the predators, ten inflorescences from three trees and 20 cm<sup>2</sup> areas of 10 trunks of three different trees were chosen at the particular orchards of those three different areas. The number given in Table 2 is the cumulative number of predators found during the total period of study from 7SMW-20 SMW, i.e., starting from the 3<sup>rd</sup> week of February (7 SMW) to the 2<sup>nd</sup> week of May (20 SMW) in 2020 and 2021.

During 2020 and 2021, predators of 20 different species belonging to the five different insect orders were identified from the mango inflorescence and trunk. Along with the increase of pest number the number of predators also started to increase from 12 SMW and attains its peak during 20

SMW. Among them, lady bird beetle was the most abundant group and a total of 448 individuals of lady bird beetles was obtained which included *R. amabilis* (202), unidentified Coccinellidae A (104), *H. axyridis* (39), *C. sexmaculatus* (34), *C. transversalis* (8), *M. discolor* (11), *P. japonica* (16), *S. (pullus) posticalis* (20), unidentified Coccinellidae B (11), *Coccinella* sp. (03). The total number of green lacewing (Chrysopidae) was 335. The total number of hemipteran insects was 74 which included Brown marmorated stink bug (12), *Scantius* sp. (45), *Chivaria* sp. (17). The total number of predatory spiders was 41 which included *O. tridens* (06), *P. agricola* (01), *O. javanicus* (06), *P. pakulli* (04), *L. decorata* (15), *H. semicuprius* (09). On the basis of the insects order, the most abundant order was the Coleoptera (448) followed



**Fig 2:** Some significant Hemipteran and Neuropteran insects act as predators of the MMB in mango orchards.



**Fig 3:** A few significant spiders act as predators of the MMB in mango orchards.

by Neuroptera (335). Then again on the basis of the individual taxa wise the most abundant was the Neuropteran green lacewings (335) followed by *R. amabilis* (202), unidentified Coccinellidae A (104), *Scantius* sp. (45), *H. axyridis* (39) and others.

The abundance of green lacewing larva is 37.30% followed by *Rodalia amabilis* (22.49%) and others. Coccinellids ladybird beetles, bugs and spiders were also found to be active predator but less abundant (Fig 5). Among the spiders *L. decorate* was most prevalent. The diversity indices of the predatory arthropods have been depicted in Table 3.

The results of the current study have clearly demonstrated that mango mealy bug serve as a most notorious sucker pest on the mango inflorescence by causing severe damage. During the observation in the year 2020 and 2021 at the selected places of Manikchak, Ratua and Chanchal of the district Malda, West Bengal significant MMB infestation was seen. In the year 2020, MMB started to appear either from 5 or 6 SMW and then the number gradually increased from 7 SMW to the 15 SMW, in inflorescence and then subsequently decreased from 19 SMW. Similarly, it was also noticed that the number of MMB gradually increased from 7 SMW to the 20 SMW, on the trunk of the mango tree.

In the year 2021, it was observed that the the number of MMB steadily increased in inflorescence from 7 SMW to the 18 SMW, in inflorescence. In case of the trunk, the

number of MMB gradually increased proportionally with the SMW, from 7 SMW to 20 SMW.

This current observation corroborated with the findings of Kumar *et al.* (2009), who had depicted that the incidence of MMB *i.e.* *D. mangiferae* started from January-February (4-7 SMW) and then peaked up in April (12-16 SMW) and then subsumed from 18 SMW *i.e.* May onwards. The pest dynamics result of current study also in line with Sarkar *et al.* (2017), where it has been observed that occurrence of MMB started from December remains active in the inflorescence and trunk till May.

According to Rahman and Khan (2009), the beneficial insect such predators and parasitoids play crucial role in the agricultural ecosystem by diminishing insect pest populations in the field. Rahman and Khan (2009) had reported a few natural enemies of mango insect pest but has found that even that low number was good enough to prevent the pest to go beyond economic thresholds (ET) for avoiding the yield loss inflicted by the pest. In this study the beneficial insects and spiders can also able to reduce the number of pest population successfully. The repeated use of huge amount of insecticide can be curbed by using beneficial insects in the orchard system (Kalita and Borah 1993). The predator diversity study of mango mealy bug in the current observation is corroborating with the findings of Pickett *et al.* (1946), Duale, (2005) and Chatterjee *et al.* (2009). According to their observations, coleopteran beetles

**Table 1:** The relative demonstration of the changeable species presence of predators in mango tree inflorescence and its trunk in relation to the three specified orchards.

Natural enemies	Occurrence of natural enemies*					
	Orchard 1		Orchard 2		Orchard 3.	
	INFLR	Trunk	INFLR	Trunk	INFLR	Trunk
<i>Rodolia amabilis</i>	●	●	●	●	●	●
Unidentified Coccinellidae A	○	○	●	●	●	●
Green lacewing (family chrysopidae)	○	●	○	●	●	●
<i>Harmonia axyridis</i>	●	●	○	●	●	●
<i>Chelomenes sexmaculatus</i>	●	○	○	●	○	●
<i>Coccinella transversalis</i>	○	○	○	●	○	○
Coccinellidae sp.	●	○	○	○	○	○
<i>Micraspis discolor</i>	○	○	○	○	●	●
<i>Propylaea japonica</i>	●	●	○	○	○	●
<i>Scymnus(pullus) posticalis</i>	●	●	○	○	○	●
Unidentified coccinellidae B	○	○	●	●	○	○
Brown marmorated stink bug	●	●	○	○	○	●
<i>Scantius</i> sp.	○	●	○	●	○	●
<i>Chivaria</i> sp.	○	○	○	●	○	●
<i>Oxyopes tridens</i>	○	○	●	○	●	○
<i>Pardosa agricola</i>	○	○	○	○	○	●
<i>Oxyopes javonicus</i>	○	○	○	●	○	●
<i>Plexxipuspa kulli</i>	○	○	○	○	○	●
<i>Leucauge decorate</i>	○	●	○	●	○	●
<i>Hyllus semicuprius</i>	○	●	○	●	○	○

Table 2: Comparative study of relative incidence of the predators found in the inflorescence of insecticide treated (INSTR) and untreated (INSUR) orchards.

Predators	Area-I Manikchak						Area-II Ratua						Area-III Chanchal					
	Insecticides treated			Insecticides untreated			Insecticides treated			Insecticides untreated			Insecticides treated			Insecticides untreated		
	2020	2021	2020	2021	2020	2021	2020	2021	2020	2021	2020	2021	2020	2021	2020	2021	2020	2021
<i>R. amabilis</i>	12 (3.54)	18 (4.30)	22 (4.74)	28 (5.34)	07 (2.74)	17 (4.18)	19 (4.42)	37 (6.12)	15 (3.94)	05 (2.35)	09 (3.08)	13 (3.67)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)
Unidentified Coccinellidae A	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	06 (2.55)	08 (2.92)	13 (3.67)	19 (4.42)	02 (1.58)	15 (3.94)	10 (3.24)	31 (5.61)	16 (4.06)	16 (4.06)	32 (5.70)	38 (6.20)	13 (3.67)	24 (4.95)
Green lacewing (family Chrysopidae)	02 (1.58)	00 (0.71)	04 (2.12)	04 (2.12)	04 (2.12)	02 (1.58)	07 (2.74)	05 (2.35)	04 (2.12)	02 (1.58)	02 (1.58)	03 (1.87)	02 (1.58)	02 (1.58)	04 (2.12)	01 (1.22)	00 (0.71)	00 (0.71)
<i>H. axyridis</i>	03 (1.87)	02 (1.22)	06 (2.55)	05 (2.35)	02 (1.58)	01 (1.22)	04 (2.12)	04 (2.12)	01 (1.22)	00 (0.71)	03 (1.87)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)
<i>C. sexmaculatus</i>	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	01 (1.22)	00 (0.71)	03 (1.87)	04 (2.12)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	02 (1.58)	02 (1.58)	04 (2.12)	01 (1.22)	00 (0.71)	00 (0.71)
<i>C. transversalis</i>	02 (1.58)	00 (0.71)	00 (0.71)	01 (1.22)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)
<i>Coccinella</i> sp.	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)
<i>M. discolor</i>	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)
<i>P. japonica</i>	02 (1.58)	00 (0.71)	04 (2.12)	03 (1.87)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	04 (2.12)	02 (1.58)	05 (2.35)	00 (0.71)	00 (0.71)	00 (0.71)
<i>S. (pullus) posticalis</i>	04 (2.12)	02 (1.58)	02 (1.58)	06 (2.55)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)
UnidentifiedCoccinellidae B	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	02 (1.58)	01 (1.22)	04 (2.12)	04 (2.12)	02 (1.58)	01 (1.22)	04 (2.12)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)
Brown marmorated stink bug	02 (1.58)	01 (1.22)	03 (1.87)	04 (2.12)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)
<i>Scantius</i> sp.	07 (2.74)*	05 (2.35)	11 (3.39)	08 (2.92)	00 (0.71)	02 (1.58)	03 (1.87)	04 (2.12)	00 (0.71)	02 (1.58)	03 (1.87)	02 (1.58)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)
<i>Chivaria</i> sp.	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	02 (1.58)	02 (1.58)	04 (2.12)	03 (1.87)	00 (0.71)	02 (1.58)	04 (2.12)	03 (1.87)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)
<i>O. tridens</i>	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	01 (1.22)	00 (0.71)	00 (0.71)	02 (1.58)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)
<i>P. agricola</i>	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)
<i>O. javonicus</i>	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	01 (1.22)	01 (1.22)	01 (1.22)	00 (0.71)	01 (1.22)	01 (1.22)	01 (1.22)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)
<i>P. pakulli</i>	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)
<i>L. decorata</i>	01 (1.22)	02 (1.58)	04 (2.12)	02 (1.58)	01 (1.22)	00 (0.71)	02 (1.58)	00 (0.71)	01 (1.22)	00 (0.71)	01 (1.22)	02 (1.58)	01 (1.22)	01 (1.22)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)
<i>H. semicuprius</i>	00 (0.71)	01 (1.22)	01 (1.22)	03 (1.87)	00 (0.71)	02 (1.58)	01 (1.22)	01 (1.22)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)	00 (0.71)

and spiders serve as two important groups of predators which feed on a diverse insect pest population. In the current study of predators of mango mealy bug, the existence of ten ladybird beetles belonging to family Coccinellidae and six different species of spiders were also noticed belonging to family Lycosidae, Oxyopidae, Tetragnathidae, Salticidae. The ladybird beetles which were found to act predators were *R. amabilis*, unidentified Coccinellidae A, *H. axyridis*, *C. sexmaculatus*, *C. transversalis*, *Coccinella* sp., *M. discolor*, *P. japonica*, *S. (pullus) posticalis* and Unidentified Coccinellidae B. Besides the ladybird beetles and spiders one larva of Green Lacewing was very much abundant as predator as per the present observation. It has been found elsewhere that these Green Lacewing larva perform a key role in the predatory guild of scale insects like mealy bug (Tanga *et al.*, 2013).

In the present finding there were only two kinds of effective hemipteran insects showing predatory effects on MMB, those were brown marmorated stink bug, *Scantius* sp. and *Chivaria* sp. This current observation on predator abundance and diversity study is very much similar to the

observation of Tanga *et al.* (2013) who had reported that the both larvae and adults of lacewings, a few species of spiders and various ladybird beetles and bugs act as the most important dominating predatory organisms of the *D. mangiferae*. He had also reported that in Tanzania and Kenya, the most abundant predatory spider families of mango mealy bug are Oxyopidae, Salticidae and Thomisidae. Six different species of spiders belonging to different families were identified during the current study and the most abundant were spiders of the families Oxyopidae, Salticidae, Lycosidae. So, it very much similar to the observation of Tanga (2013). According to the (Tanga *et al.*, 2013; Moore 1988; Iperti 1999) the beetles belonging from the Coccinellidae act as most abundant natural enemies in the each and every mango orchards. Just similar to these earlier observations in current study also a huge numbers of lady bird beetles were found devouring the mealy bug colonies found in the inflorescence as well as trunk. Instead of the diverse predators the mealy bugs were still persisting in high numbers. The number of the predators/inflorescence or in the trunk was not enough to curb the




Fig 4: Stereo-microscopic view of the *Cryptochaetum* sp. nr. *iceryae* (Williston) of (a) and (b) Dorsal view (c) Lateral view.




Fig 5: Relative abundance of different predators of mango mealy bug at Malda.



**Table 3:** Diversity indices of the predators in the study area.

Ecological Index	Value
Dominance (D)	0.21
Simpson (1-D)	0.78
Shannon(H)	2.03
Evenness- $e^H/S$	0.38
Margalef	2.79
Equitability (J)	0.67
Fisher (alpha)	3.62
Berger-parker	0.37

mealy bug growth. So, mass rearing of these predators and release in natural adobe should be done if possible to check mango mealy bug.

According to the (Heong and Escalada, 1997) in Philippine, there were 14 hymenopteran insect species and 46 bugs species can successfully reduce insect pests by parasitizing on those particular insects. According to the reports by (Tanwar *et al.*, 2008) there was only one parasitoid species, *Aenasius bambawalei* belonging to the Family Encyrtidae (Order - Hymenoptera) had emerged out from body of the MMB. Hayat (2009) reported that, in India *A. bambawalei* serve as one of the major copious parasitoid in the mealy bug affected orchard system. According to the reports of Noyes and Hayat (1994) in India a few parasitoids genus of MMB were *R. iceryoides* namely *Leptomastrix* sp., *Anagyrus* sp. and *Praleurocerus* sp and *Cryptochaetum* sp.nr. *iceryae* serve as the parasitoid insect of MMB. In the current study also only one parasitoid was found to emerge from mango mealy bug and that was *Cryptochaetum* sp. nr. *iceryae* (Williston) which is an established effective parasitoid dipteran insect against the MMB, *D. mangiferae*. In present study average parasitization percentage in insecticide untreated orchard was 14.83% while emergence from the two different dates was considered. The application of insecticide reduced the parasitization percentage. On the contrary average parasitoid emergence percentage was in insecticide tread field was found to be 7.66% much lesser compared to insecticide untreated field. Therefore, it can be concluded with lesser use of chemical pesticide parasitization percentage increases which can bring about natural control if these parasitoids can be reared and released in sufficient quantity in the field. There are so many case studies where parasitoid release had brought about significant reduction of pest number.

## CONCLUSION

The conservation of the natural enemies can act as one of the main strategies to combat mango mealy bug. Moreover, augmentation of the most abundant predators and parasitoids through mass rearing and then releasing them in the infested orchard can bring down the pest number drastically. Therefore this natural enemy study of mango mealy bug can open new vistas in biological pest control in near future.

## ACKNOWLEDGEMENT

The authors will remain indebted to Dr. J. Poorani, the Principal Scientist, National Research Centre for Banana (NRCB), Trichirapalli, Tamil Nadu, for the identification of the parasitoid of mango mealy bug.

## Conflict of interest

The authors declare that there is no conflict of interest among the authors.

## REFERENCES

- Angelfire, (2022). Common species of Coccinellidae of the Indian Region. Retrieved from [https://www.angelfire.com/bug2/j\\_poorani/morphology.htm](https://www.angelfire.com/bug2/j_poorani/morphology.htm).
- Anonymous, (2018). Horticulture statistics at a glance. Horticulture Statistics Division, Department of Agriculture, Government of India.
- Bugguide, (2022). Identification of Hepimteran insects. Retrived from <http://bugguide.net/node>.
- Chatterjee, S., Isaia, M., Venthurino, E. (2009). Spider as biological controllers in the agro-ecosystem. *Journal of Theoretical Biology*. 258: 352-362.
- Chrysantus, M.C. (2012). Bioecology of the Mango Mealybug, *Rastrococcus iceryoides* Green (*Hemiptera: Pseudococcidae*) and its Associated Natural Enemies in Kenya and Tanzania. University of Pretoria, South Africa.
- Duale, A.H. (2005). Effect of temperature and relative humidity on biology of stem borer parasitoid *Pediobius fuvvus* (Gahan) (*Hymenoptera: Eulophidae*) for management of stem borers. *Environmental Entomology*. 34(1): 1-3.
- Hayat, M. (2009). Description of a new species of *Aenasius* Walker (*Hymenoptera: Encyrtidae*), parasitoid of the mealybug, *Phenacoccus solenopsis* Tinsley (*Homoptera: Pseudococcidae*) in India. *Biosystematica*. 3: 21-26.
- Heong, K.L. and Escalada, M.M. (1997). Pest Management of Rice Farmers in Asia (Manila: International Rice Research Institute). Phillipines.
- Iperti, G. (1999). Biodiversity of predaceous coccinellidae in relation to bio-indication and economic importance. *Agricultural Ecosystem and Environment*. 74: 323-342.
- Ishaq, M., Usman, M., Asif, M., Khan, I.A. (2004). Integrated pest management of mango against mealy bug and fruit fly. *International Journal of Agricultural Biology*. 6: 452-454.
- Jakhar, B.L., Singh, N., Venilla, S., Patel, M.H., Vekaria, M.V., Patel, D.B., Panickar B. (2016). Influence of climate change on *Helicoverpa armigera* (Hubner) in pigeon pea. *Journal of Agricultural Ecology*. 2: 25-31.
- Jha, S.N., Kingsly, A.R.P. and Chopra, S. (2006). Physical and mechanical properties of mango during growth and storage for determination of maturity. *Journal of Food Engineering*. 72(1): 73-76.
- Jouveau, S., Delaunay, M., RégineVignes, L. and Nattier, R. (2018). A multi-access identification key based on colour patterns in ladybirds (*Coleoptera, Coccinellidae*). *Zookeys*. 758: 55-73.
- Kalita, D.N. and Borah, D.C. (1993). Parasitoids and predators of jute pests in certain localities of Central Brahmaputra valley zone of Assam. *Indian Journal of Agricultural Science Society*. 6(1): 19-23.



- Karar, H. (2010). Bioecology and management of mango mealy bug, *Drosicha mangiferae* Green in mango orchards of Punjab, Pakistan. Ph.D. thesis, University of Agriculture, Agriculture University Road, Faisalabad, Pakistan. Zip Code.
- Kumar, A., Pandey, S.K., Kumar, R. (2009). Population dynamics of mango mealy bug, *Drosicha mangiferae* Green from Jhansi, Uttar Pradesh Biological forum. An International Journal. 1(2): 77-79.
- Larson, D.J. (2013). Key to lady beetles (*Coleoptera: Coccinellidae*) of Saskatchewan.
- Litz, R.E. (1997). The Mango: Botany, Production and Uses. CAB International, University Press, Cambridge.
- Moore, D. (1988). Agents used for biological control of mealybugs (*Pseudococcidae*). Biocontrol News and Information. 9: 209-225.
- NBAIR (2022). Identification characters of *Cryptochaetum* sp. nr. *iceryae*. [https://databases.nbair.res.in//Featured\\_insects/Cryptochaetum-iceryae.php](https://databases.nbair.res.in//Featured_insects/Cryptochaetum-iceryae.php).
- Noyes, J.S. and Hayat, M. (1994). Oriental mealybug parasitoids of the Anagyrini (*Hymenoptera: Encyrtidae*). Wallingford, United Kingdom, CAB International.
- Pena, J.E., Mohyuddin, A.I., Wyoski, M. (1998). A review of the pest management situation in Mango agroecosystem. *Phytoparasitica*. 26(2): 1-20.
- Pickett, A.D., Patterson, N.A., Stultz, H.T., Lord, F.T. (1946). The influence of spay program on the fauna of apple orchards in Nova Scotia: An appraisal of the problem and a method of approach. *Scientific Agriculture*. 26: 590-600.
- Rahman, S. and Khan, M.R. (2009). Natural enemies of insect and mite pests of jute ecosystem. *Annals Plant Protection Science*. 17(2): 466-467.
- Sahoo, S.K. and Jha, S. (2009). Bioecology of Mango Fruit borer, *Autocharis* (=Noorda) *Albizonalis* Hampson (*Pyrallidae, Lepidoptera*) - A Recent Threat to Mango Growers in West Bengal, India. *Proceedings of VIII<sup>th</sup> International Mango Symposium* (Ed: Oosthuysen, S.A.). *Acta Horticulture*. 1345-1425.
- Sarkar, A., Nandi, P.S. and Chakraborty, K. (2017). Description, Bionomics and Bio-Rational Control of Mango Mealy Bug, *Drosicha mangiferae* Glover (*Coccidae: Hemiptera*) at Malda, West Bengal, *International Journal of Applied Environmental Sciences*. 12(4): 661-672.
- Tandon, P.L. and Lal, B. (1978). The mango coccid, *Rastrococcus iceryoides* Green (*Homoptera: Coccidae*) and its natural enemies. *Current Science*. 13: 46-48.
- Tanga, C.M., Ekesi, S., Govender, P., Mohamed, S.A. (2013). Effect of six host plant species on the life history and population growth parameters of *Rastrococcus iceryoides* (Hemiptera: *Pseudococcidae*). *Floral Entomology*. 96(3): 1030-1041.
- Tanwar, R.K., Bhamare, V.K., Ramamurthy, V.V., Hayat, M., Jeyakumar, P., Singh, A., Bambawale, O.M. (2008). Record of new parasitoid on mealybug, *Phenacoccus solenopsis*. *Indian Journal of Entomology*. 70: 404-405.
- Yadav, J.L., Singh, S.P. and Kumar, R. (2004). The population density of the mango mealy bug (*Drosicha mangiferae* G.) in mango. *Progressive Agriculture*. 4(1): 35-37.