



Transcriptome Analysis of the Regulation of Retinoic Acid on the Growth and Development of Brandt's Vole

L. Batdorj[#], C.M. Xu[#], S. Li, Y. Deng, H.C. Babar¹, X.B. Tu, Z.H. Zhang

10.18805/IJArE.A-667

ABSTRACT

Background: Studying the growth and developmental mechanisms of steppe (Brandt's) voles is important for biological control and protection of steppe ecology. In this paper, preliminary biological experiments' data showed that the mortality of Brandt's vole increased after retinoic acid treatment.

Methods: LPH inhibitor, retinoic acid, quercetin and DMSO (Standard treatment concentration 10mg/ml) were given them through oral with the help of Animal oral gavage feeding needles. The hind legs muscles of surviving Brandt's voles were collected for transcriptomic analysis.

Result: A total of 4047 and 2251 up- and down-regulated transcripts, respectively, were observed upon retinoic acid treatment compared to the DMSO (control) treatment. Transcriptomic changes were enriched significantly in 31 KEGG pathways, most of which were involved in sugar metabolism, particularly for fructose and mannose (ko00051), protein digestion and absorption (ko04974), phenylalanine metabolism (ko00360), tyrosine metabolism (ko00350) and the PI3K-Akt signaling pathway (ko04151), which is related to growth and development. A total of 114 differentially expressed genes were screened as candidate genes according to the enrichment results, including c-Jun N-terminal kinase and tubulin epsilon chain verified by RT-PCR. Ultimately, we proposed a preliminary regulatory molecular mechanism of retinoic acid relevant to mouse development and death.

Key words: Brandt's voles, Molecular mechanism, Retinoic acid, Transcriptomic.

INTRODUCTION

Brandt's vole is distributed throughout Russia, Mongolia, Inner Mongolia and China. And is one of the most harmful rats to grassland ecology, particularly in the temperate steppe of China. Brandt's voles not only overgraze the forage grass, resulting in a substantial decrease in the livestock carrying capacity of the region with high population levels, but also accelerate vegetative degradation and desertification. For example, after vole caves are abandoned, the ecological environment needs 3-5 years to recover, which seriously affects the development of animal husbandry (Wang *et al.*, 2010). Integrated control of Brandt's voles is a complex animal husbandry ecological project. At present, the primary control measures that are practiced include artificial capture and stocking of natural enemies, such as snakes and cats and the use of chemicals such as anticoagulant rodenticide (Hou *et al.*, 1991). In order to improve the prevention and control measures, we aimed to uncover different strategies that might help assist in the management of Brandt's voles from the perspective of limiting biological growth and development. Toward this goal, we explored the mechanisms that regulate the growth and development of Brandt's voles.

Retinoids are important biological molecules that act in cell proliferation, epithelial cell growth and maturation, apoptosis, immunological functions and are vital in establishing organ development during embryogenesis (Niederreither *et al.*, 2008). Lactase (lactase-phlorizin hydrolase, or LPH) is a member of the β -galactosidase family of enzymes and is involved in the hydrolysis of the disaccharide lactose into

The Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing, 100193, P.R. China.

¹Department of Entomology, Sindh Agriculture University Tandojam 70060, Sindh, Pakistan.

These authors contributed equally to this work.

Corresponding Author: Z.H. Zhang, The Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing, 100193, P.R. China. Email: zhangzehua@caas.cn

How to cite this article: Batdorj, L., Xu, C.M., Li, S., Deng, Y., Babar, H.C., Tu, X.B. and Zhang, Z.H. (2022). Transcriptome Analysis of the Regulation of Retinoic Acid on the Growth and Development of Brandt's Vole. Indian Journal of Agricultural Research. DOI: 10.18805/IJArE.A-667.

Submitted: 07-06-2021 **Accepted:** 01-09-2022 **Online:** 11-11-2022

constituent galactose and glucose monomers (Day *et al.*, 2000). Quercetin has antioxidant and anti-inflammatory effects which might help reduce inflammation, kill cancer cells, control blood sugar and help prevent heart disease (Ali *et al.*, 2018; Kim *et al.*, 2018; Kim *et al.*, 2019). These three chemical molecules all play a regulatory role in apoptosis in the growth and development of mammals.

The purpose of this study was to find the most significant treatment for controlling Brandt's vole populations by comparing the mortality and body weight changes of Brandt's voles under various chemical treatments. Furthermore, we screened differentially expressed genes related to the growth, development and death of mice under this treatment to explore a new mechanism for regulating the growth and development of Brandt's vole.

MATERIALS AND METHODS

Sample preparation

To study the function of retinoic acid in female *B. vole* development, we selected female mice for test (Nowickij *et al.*, 2008). 270 *B. voles*, 108 of them were female voles, were captured in the Naranbulag Soum, Abagage, Xilingol League, Inner Mongolia, China, from where Brandt's vole populations began expanding, from June 12, 2018 to July 08, 2018.

Female *B. Voles* divided into 35 cages, each cage for 3 to 4 voles as a group. *B. Voles* are stored at 23±1°C temperature room. The size of a small iron tank containing *B. Voles* is (27 × 28 × 22 cm) with sawdust for comfortable environment. Every day prepare food (carrot) and water twice (morning and evening).

LPH inhibitor, retinoic acid, quercetin and DMSO (Standard treatment concentration 10 mg/ml) were given them through oral with the help of Animal oral gavage feeding needles and 1 ml syringe was used. Depending on bodyweight, each vole treated with a standard of 0.5 ml/30 g. The doses were given them in the interval of 24 hours, 72 hours and 120 hours. Each treatment has five repeats, each repeat with 5 voles. After each treatment, the hind legs muscles of surviving Brandt's voles were randomly sampled and mixed. All the samples were frozen at -80°C for transcriptomic analysis.

RNA extraction, library construction and sequencing

Total RNA was isolated from all four treatments using the RNA plant mini kit with on-column DNase digestion (Qiagen, Hilden, Germany) following the manufacturer's instructions. RNA degradation and contamination were detected on 1% agarose gels. RNA concentration was measured using a Qubit RNA Assay Kit in Qubit 2.0 Fluorometer (Life Technologies, Carlsbad, CA, USA). Additionally, RNA integrity was assessed using the RNA Nano 6000 Assay Kit of the Bioanalyzer 2100 system (Agilent Technologies, Santa Clara, CA, USA).

A total amount of 3 µg RNA per sample was used as input material for cDNA preparations. Finally, three samples with RNA integrity number (RIN) values >8 were used for construction of cDNA libraries. Sequencing libraries were generated using the NEBNext Ultra™ RNA Library Prep Kit for Illumina (NEB, USA) following manufacturer's instructions.

The clustering of the index-coded samples was performed on a cBot Cluster Generation System using TruSeq PE Cluster Kit v3-cBot-HS (Illumina) according to the manufacturer's instructions. After cluster generation, the library preparations were sequenced on an Illumina HiSeq 2500 platform and 125 bp paired-end reads were generated.

Sequence reads, mapping, assembly and annotation

Raw data (raw reads) of fastq format were firstly processed through in-house Perl scripts. In this step, clean data (clean reads) were obtained by removing reads containing adapters, reads containing ploy-N and low-quality reads from

raw data. At the same time, Q20, Q30, GC-content and sequence duplication level of the clean data were calculated. All the downstream analyses were based on clean data with high quality. Transcriptome assembly was accomplished based on left.fq and right.fq using Trinity with min-kmer-cov set to 2 by default and all other parameters set default (Hao *et al.*, 2017).

Gene function was annotated based on following seven databases: Nr (NCBI non-redundant Protein sequences), Nt (NCBI non-redundant nucleotide sequences), Pfam (Protein family), KOG/COG (Clusters of Orthologous Groups of proteins), Swiss-Prot (A manually annotated and reviewed protein sequence database), KO (KEGG Ortholog database) and GO (Gene Ontology), using Diamond software (Benjamin *et al.*, 2015).

DEGs analysis

Prior to DEG analysis, for each sequenced library the read counts were adjusted using the edge R program package through one scaling normalized factor. DEG analysis of two samples was performed using the DEGseq R package (Anders *et al.* 2010). P value was adjusted using Padj value (Storey *et al.* 2003). Padj value ≤ 0.05 and |log₂ (fold change) ≥ 1 was set as the threshold for significantly differential expression.

KEGG enrichment analysis of DEGs

KEGG is a database resource for understanding high-level functions and utilities of the biological systems (Kanehisa *et al.* 2008) at the cell, the organism, or the ecosystem level. We used KOBAS software to test the statistical enrichment of differential expression genes in KEGG pathways (Mao *et al.* 2005).

RESULTS AND DISCUSSION

Regulation of all-trans retinoic acid in various animals

High concentrations of all-trans retinoic acid (atRA) inhibit adipogenic differentiation of rat bone marrow mesenchymal stem cells. The mechanism of retinoic acid nuclear receptor β (PPAR β) regulates peroxisome proliferation-activated receptor β (PPAR β) and CCAAT enhancer binding protein α (C/EBP α) during adipogenic differentiation of rat bone marrow mesenchymal stem cells (Zhang *et al.*, 2013). Retinoic acid, as a transcriptional regulatory factor, has a

Table 1: Statistics of mortality and vole body weight change under four treatments. We used the average of surviving vole post-treatment body weight minus pre-treatment body weight as body weight change.

Treatment	Death number	Mortality	Body weight change	STDEV
CK	8	32%	0.79	3.84
LPH inhibitor	9	36%	-0.59	5.50
Quercetin	14	56%	1.45	5.88
Retinoic acid	21	84%	2.30	0.67

wide range of physiological functions, it regulates cell proliferation and differentiation at the physiological level, which is closely related to normal growth and development; as a teratogen, retinoic acid can induce a variety of tissue and organ deformities in a variety of organisms (Feng *et al.*, 2018). There is a feedback regulation cycle of migratory locust rhythm genes *clk*, *cry* and *per*, which are all regulated by all-trans retinoic acid and promote migratory locust diapause. Upstream all-trans retinoic acid, *rarp* and rhythm genes promote the expression of downstream *dh*, which in turn regulates the insulin transactivation pathway (Babar *et al.*, 2019; Hao *et al.*, 2019).

Mortality statistics

Retinoic acid, the first morphogen described so far in vertebrates, is a vitamin A derivative which exerts striking effects on development and differentiation. The identification of three retinoic acid receptors as members of the nuclear receptor super-family provides an explanation for the molecular action of morphogens on gene expression (de Thé *et al.*, 1990). RA levels require precise regulation by controlled synthesis and catabolism and when RA concentrations deviate from normal levels, in either direction, abnormal growth and development occurs (McCaffery *et al.*, 2003). In rodents, the control of neural patterning and differentiation are disrupted when RA concentrations are lowered, whereas inappropriately high concentrations of RA result in abnormal development of the cerebellum and hindbrain nuclei. The latter parallels the malformations seen in the human embryo exposed to RA due to treatment of the

mother with the acne drug Accutane (13-cis RA). In cases where the child survives beyond birth, a particular set of behavioral have been described. Even adult brains might be susceptible to an imbalance of RA, particularly the hippocampus (McCaffery *et al.*, 2003). According to the mortality, it could be seen that the mortality rate of *B. voles* after treatment with retinoic acid was 81%, which was significantly higher than that of DMSO, which was only 32%. Thus, we took retinoic acid-treated and DMSO-treated Brandt voles' hind leg muscles for transcriptome sequencing.

Illumina HiSeq mRNA sequence

In total, 494.6 million short reads were generated from the six libraries (RA and DMSO), with 481.4 million classified as clean and at least 94% high-quality (Q30) reads selected for further analysis (Table 1). After *de novo* assembly was completed, we aligned unigene sequences to protein databases, including NR, Swiss-Prot, KEGG, KOG, eggNOG, GO and Pfam (e-value<1e-5) via Diamond software (Benjamin *et al.*, 2015). We then retrieved proteins with the highest sequence similarity to the given transcripts along with their functional annotations. Of the 48844 unigenes, we found 21513 that (44%) that were annotated (Table 2).

The UniGene database obtained by splicing was used as a database and the expression abundance of each Unigene in each sample was identified by sequence similarity alignment. 86.79% ~ 87.89% of clean reads from six samples were compared to the Unigene database by using the bowtie2 software (Langmead *et al.*, 2012) (Table 2) and the eXpress software (Roberts *et al.*, 2013) to calculate

Table 2: Summary statistics of clean reads in the muscle transcriptomes of voles.

Sample	DMSO_1	DMSO_2	DMSO_3	Retinoic_Acid_1	Retinoic_Acid_2	Retinoic_Acid_3
Raw-reads	82256562	82035790	82513878	82919344	81743044	83148232
Clean-reads	79958142	79995526	79899552	80773666	79705930	81122228
Q30	94.64%	95.02%	94.18%	94.83%	94.96%	95.03%
Reads mapped	88.23%	87.61%	87.89%	87.10%	86.83%	86.79%
Total unigene	48844					
Annotated	21513					




Fig 1: Correlation between the transcriptomes of two treatment samples. (a) Correlation coefficient analysis of RNA-seq data from six samples. (b) PCA plot showing clustering of transcriptomes of DMSO-fed and retinoic acid-fed Brandt's voles.

the Unigene expression FPKM value. The correlation coefficient were above 0.99, indicating high quality of the experimental replicates (Fig 1).

Differentially expressed genes (DEGs) between DMSO and Retinoic acid treatments

A total of 4047 and 2251 up- and down-regulated transcripts were observed in the retinoic acid treatment compared to the DMSO (control) treatment ($FDR \leq 0.001$ and $|\log_2 \text{ratio}| \geq 1$), respectively (Fig 2).

GO and KEGG pathway enrichment analysis

We analyzed the KEGG enrichment of the 6298 DEGs and found that the DEGs were enriched in 225 different signaling pathways including: retinol metabolism (ko00830), the Foxo signaling pathway (ko 04068) and Apoptosis-multiple species (ko 04215), along with 31 metabolic pathways were

significantly enriched ($P\text{-value} < 0.05$) including: ECM-receptor interaction (ko 04512), protein digestion and absorption (ko 04974) and the PI3K-Akt signaling pathway (ko 04151) (Table 3). At the same time, we analyzed 1231 genes from 6289 DEGs annotated to the GO database by GO enrichment. A total of 4861 assigned terms were obtained, of which 1845 terms were significantly enriched. We found 847 GO terms with the corresponding number of DEGs greater than two among the three GO classifications (Molecular function, cellular component and biological process) (Fig 3.).

Combining the analyses of functional enrichment from GO, KEGG and phenotypic data, we selected 114 genes related to cell proliferation, cell division, regulation of biological growth and development and regulation of cell death related metabolic pathways, such as cyclic AMP-




Fig 2: Differential gene expression in the DMSO treatment as compared to the retinoic acid treatment. (a) Number of up-regulated (upper bars) and down-regulated (lower bars) genes examined in the comparison. (b) Clustering Heatmap of the total 6298 DEGs. Red bands indicate up-regulated genes while blue bands indicate down-regulated genes.




Fig 3: GO enrichment analysis top 30. screening GO entries with a number of different DEGs greater than 2 in the three classifications, according to the $-\log_{10} P$ value of each entry, 10 items in descending order. Caption: The X coordinate in the figure is the name of the GO entry and the Y coordinate is $-\log_{10} P$ value.

responsive element-binding protein 3-like protein 1, gamma-aminobutyric acid receptor-associated protein-like 1, phosphorylase b kinase gamma catalytic chain, skeletal muscle/heart isoform and SHC-transforming protein 4 isoform X1 (Fig 4.)

Validation by RT-PCR

DEGs of c-Jun N-terminal kinase and tubulin epsilon chain, were used to verify the RNA sequencing results. Our results showed that the relative expression of c-Jun N-terminal




Fig 4: 114 candidate DEGs selected based on GO and KEGG enrichment results. Red in the picture indicates that the expression of the corresponding gene is up-regulated and blue indicates that the expression of the corresponding gene is down-regulated.




Fig 5: The relative quantitative expression trends of two specific DEGs.

Table 3: Significantly enriched KEGG pathways comparing retinoic acid treatment with DMSO treatment.

Group	Num.	ID	Term	Nos.differentially expressed genes		P-Value
				Up-regulated	Down-regulated	
Retinoic acid	1	ko02020	Two-component system	4	1	0.000194
	2	ko04964	Proximal tubule bicarbonate reclamation	3	3	0.001868
VS	3	ko04512	ECM-receptor interaction	16	2	0.002765
DMSO	4	ko04140	Regulation of autophagy	3	3	0.003576
	5	ko00901	Indole alkaloid biosynthesis		1	0.004742
	6	ko00643	Styrene degradation	1		0.004742
	7	ko00780	Biotin metabolism		2	0.005561
	8	ko00565	Ether lipid metabolism	5	3	0.005603
	9	ko00051	Fructose and mannose metabolism	2	6	0.005603
	10	ko04974	Protein digestion and absorption	10	2	0.006862
	11	ko03440	Homologous recombination	5	2	0.007056
	12	ko00790	Folate biosynthesis	3		0.00966
	13	ko00531	Glycosaminoglycan degradation	4		0.009932
	14	ko00965	Betalain biosynthesis		1	0.013576
	15	ko00360	Phenylalanine metabolism	2	1	0.020859
	16	ko00564	Glycerophospholipid metabolism	8	6	0.021362
	17	ko04666	Fc gamma R-mediated phagocytosis	9	6	0.021369
	18	ko04139	Regulation of mitophagy - yeast	1	4	0.023375
	19	ko00340	Histidine metabolism	2	1	0.025829
	20	ko00300	Lysine biosynthesis	1		0.025916
	21	ko00130	Ubiquinone and other terpenoid-quinone biosynthesis	2		0.027115
	22	ko00350	Tyrosine metabolism	3	1	0.034798
	23	ko00460	Cyanoamino acid metabolism	1	1	0.035416
	24	ko04977	Vitamin digestion and absorption		3	0.037735
	25	ko04978	Mineral absorption	2	3	0.038956
	26	ko04672	Intestinal immune network for IgA production	3	2	0.043679
	27	ko00627	Aminobenzoate degradation	2		0.044866
	28	ko00630	Glyoxylate and dicarboxylate metabolism	3	1	0.04576
	29	ko04640	Hematopoietic cell lineage	3	5	0.046062
	30	ko04151	PI3K-Akt signaling pathway	31	5	0.048086
	31	ko00500	Starch and sucrose metabolism	4	1	0.048746

kinase and tubulin epsilon chain was upregulated, which were similarly to what was found in the RNA sequencing results (Fig 5).

CONCLUSION

DEGs of Brandt's voles treated with retinoic acid were enriched in multiple KEGG signaling pathways including: ko04512:ECM-receptor interaction, ko04140:Regulation of autophagy, ko00051:Fructose and mannose metabolism, ko04974:Protein digestion and absorption, ko04977:Vitamin digestion and absorption, ko04151:PI3K-Akt signaling pathway, ko04911:Insulin secretion, ko04713:Circadian entrainment, ko04215:Apoptosis-multiple species, ko04068:FoxO signaling pathway, ko04210:Apoptosis, ko03320:PPAR signaling pathway, ko04910:Insulin signaling pathway, related to birth, growth and rearing. Enrichment of the ko04620: Toll-like

receptor signaling pathway revealed that retinoic acid can be involved in the regulation of growth and development of Brandt's vole by regulating these networks.

Conflict of interest: None.

REFERENCES

- Ali, A., Kim, M.J., Kim, M.Y., Lee, H.J., Roh, G.S., Kim, H.J., Cho, G.J., Choi, W.S. (2018). Quercetin induces cell death in cervical cancer by reducing O-GlcNAcylation of adenosine monophosphate-activated protein kinase. *Anatomy and Cell Biology*. 51: 274-283.
- Day, A.J., Cañada, F.J., Díaz, J.C., Kroon, P.A., Mclauchlan, R., Faulds, C.B., Plumb, G.W. *et al.* (2000). Dietary flavonoid and isoflavone glycosides are hydrolysed by the lactase site of lactase phlorizin hydrolase. *FEBS Letters*. 468: 166-170.

- Babar, H.C., Cui, B.Y., Hidayat, U., Li, S., Hao, K., Tu, X.B., Wang, G.J. *et al.* (2019). Role of PTP/PTK trans activated insulin-like signalling pathway in regulation of grasshopper (*Oedaleus asiaticus*) development. *Environmental Science and Pollution Research*. 26: 8312-8324.
- Benjamin, B., Chao, X., Daniel, H.H. (2015). Fast and sensitive protein alignment using diamond. *Nature Methods*. 12: 59-60.
- De, T.H., Vivanco, R.M.M., Tiollais, P., Stunnenberg, H., Dejean, A. (1990). Identification of a retinoic acid responsive element in the retinoic acid receptor beta gene. *Nature*. 343: 177-108.
- Feng, H., Xu, M., Zhang, Y., Han, B., Wang, J., Sun, P. (2018). Identification of differentially expressed micrnas involved in the pathogenesis of colorectal cancer. *Clinical Laboratory*. 64: 797-804.
- Hou, X.X., Dong, W.H., Zhou, Y.L., Zhang, P.L., Yang, Y.P., Zhang, Y.X., Xue, X.P. (1991). Research on integrated control of Brandt's Voles. *Chinese Journal of Vector Biology and Control*. 5: 301-304.
- Hao, K., Aftab, R.J., Hidayat, U., Tu, X.B., Nong, X.Q., Zhang, Z.H. (2019). Transcriptome sequencing reveals potential mechanisms of the maternal effect on egg diapause induction of *Locusta migratoria*. *International Journal of Molecular Sciences*. 20: 1-19.
- Hao, K., Wang, F., Nong, X.Q., McNeill, M.R., Liu, S.F., Wang, G.J., Cao, G.C., Zhang, Z.H. (2017). Response of peanut *Arachis hypogaea* roots to the presence of beneficial and pathogenic fungi by transcriptome analysis. *Scientific Reports*. 7: 964-970.
- Kanehisa, M., Araki, M., Goto, S., Hattori, M., Hirakawa, M., Itoh, M., Katayama, T., Kawashima, S. *et al.* (2008). KEGG for linking genomes to life and the environment. *Nucleic Acids Res*. 36: 480-484.
- Kim, H.R., Kim, B.M., Won, J.Y., Lee, K.A., Ko, H.M., Kang, Y.S., Lee, S.H., Kim, K.W. (2018). Quercetin, a plant polyphenol, has potential for the prevention of bone destruction in rheumatoid arthritis. *Journal of Medicinal Food*. 22: 152-161.
- Kim, S.S., Jang, H.J., Oh, M.Y. (2019). Quercetin enhances the function and reduces apoptosis of mouse islets. *Transplantation Proceedings*. 51: 1451-1457.
- Langmead, B. and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature methods*. 9: 357-359.
- Mao, X., Cai, T., Olyarchuk, J.G., Wei, L. (2005). Automated genome annotation and pathway identification using the KEGG Orthology (KO) as a controlled vocabulary. *Bioinformatics*. 21: 3787-3793.
- McCaffery, P.J., Adams, J., Maden, M., Rosa, M.E. (2003). Too much of a good thing: Retinoic acid as an endogenous regulator of neural differentiation and exogenous teratogen. *The European Journal of Neuroscience*. 18: 457-472.
- Niederreither, K. and Dollé, P. (2008). Retinoic acid in development: Towards an integrated view. *Nature Reviews. Genetics*. 9: 541-553.
- Nowickij, S.M., Chithalen, J.V., Cameron, D., Tyshenko, M.G., Petkovich, M., Wyatt, G.R., Jones, G., Walker, V.K. (2008). Locust retinoid X receptors: 9-Cis-retinoic acid in embryos from a primitive insect. *Proceedings of the National Academy of Sciences*. 105(28): 9540-9545.
- Roberts, A. (2013). Ambiguous fragment assignment for high-throughput sequencing experiments. University of California, Berkeley.
- Storey, J.D. and Tibshirani, R. (2003). Statistical Significance for Genome Wide Studies. *Proceedings of the National Academy of Sciences*. 100: 9440-9445.
- Wang, D.W., Cong, L., Wang, Y., Liu, X.H. (2010). Differences in population parameters and physiological characteristics of *Brandt's voles* during breeding and non-breeding seasons. *Acta Ecologica Sinica*. 30: 3562-3568.
- Zhang, M., Shu, X.L., Li, W.D., Chen, J., Liu, Y.X. (2013). AtRA inhibits adipogenic differentiation of rat bone marrow mesenchymal stem cells. *Journal of the Third Military Medical University*. 35: 42-46.