



# Optimization of 17 $\alpha$ -Methyl Testosterone Dose in *Artemia* Nauplii and Rotifer using Enrichment Technique

R.V. Bhosle<sup>1</sup>, J.S. Sampath Kumar<sup>2</sup>, Cheryl Antony<sup>3</sup>,  
S. Aanand<sup>4</sup>, V. Senthilkumar<sup>4</sup>, R.S.S. Lingam<sup>4</sup>

10.18805/IJAR.B-4800

## ABSTRACT

**Background:** The present study explored the possibility of using live feed, as vector for supplementing the androgenic hormone in aquaculture seed production.

**Methods:** The study consisted of five treatment doses of 17  $\alpha$ -methyl testosterone namely; control (without hormone), T1 (20 mg L<sup>-1</sup>), T2 (40 mg L<sup>-1</sup>), T3 (60 mg L<sup>-1</sup>) and T4 (80 mg L<sup>-1</sup>) and each were triplicated. The doses were tested in the live feed of *Artemia* nauplii and rotifer by following completely randomized design. The *Artemia* nauplii (*Artemia salina*), collected by hatching the *Artemia* cyst and 5-8 days old rotifers, collected from mass culture unit of rotifer, were released in their respective treatment doses at 100 and 800 individuals/ml, respectively.

**Result:** The study found significant difference in accumulation of hormone in the tissues of live feed at different time intervals. In *Artemia*, significantly higher amount of 17 $\alpha$ -MT (15.33 $\pm$ 0.09 mg L<sup>-1</sup>) was observed in 80 mg L<sup>-1</sup> group at 24 h. In rotifer, significantly higher value of 17  $\alpha$ -MT (17.42 $\pm$ 0.06 mg L<sup>-1</sup>) was recorded in 60 mg L<sup>-1</sup> group at 6 h. The level of 17 $\alpha$ -MT in *Artemia* nauplii and rotifer started showing a decreasing trend in 24 h and 6 h, respectively. Overall the findings of the present study support that supplementation of hormone through live feed is possible as live feed accumulated 17  $\alpha$ -MT in its tissue after enrichment which can be further explored for producing all-male population in a more sustainable way.

**Key words:** *Artemia*, Androgenic hormone, Bio-enrichment, 17  $\alpha$ -methyl testosterone, Live feed, Mono-sex seed production, Rotifer.

## INTRODUCTION

The recent fish production trend shows a significant contribution by aquaculture to the world fish production especially by freshwater aquaculture. As per the latest FAO report, freshwater aquaculture alone contributes around 88% of the world's farmed fish (FAO, 2020). In order to improve the quality seed production and better fish production the sector has started using various chemical derivatives like antibiotics and hormones in the recent times (Emeka *et al.*, 2014). In fish many of the phenotypic traits such as growth rate, colour pattern, *etc.* were sex based which lead to the development of mono-sex seeds, either all male or all female population. Further, the development of captive seed production systems - hatcheries - paved the way for egg or uniform yolk-sac larvae collection which opened the gate way for development of mono-sex seed production (male) using 17 $\alpha$ -methyltestosterone through oral administration (Gale *et al.*, 1996). In the past decades, many authors have reported that hormone is a naturally-occurring or synthetic androgens and estrogens which shown growth-promoting effect in many cultured fishes (Pelissero and Sumpter, 1992; James and Sampath, 2006). In the recent years, monosex seed production technology has become a common practice among the various animal production sectors (Bardhan *et al.*, 2021). However, the hormone residue is creating various negative impacts to the natural ecosystem which making the hormone based sex reversal as a much challenging task in the recent times (Azizi-lalabadi and Pirsahab, 2021). Therefore, delivering of potent hormones to the fish larvae through

<sup>1</sup>Department of Aquaculture, Fisheries College and Research Institute, Thoothukudi-628 008, Tamil Nadu, India.

<sup>2</sup>Directorate of Sustainable Aquaculture, Tamil Nadu Dr. J. Jayalalithaa Fisheries University, Nagapattinam-611 002, Tamil Nadu, India.

<sup>3</sup>Department of Aquaculture, Dr. M.G.R. Fisheries College and Research Institute, Ponneri-601 204, Tamil Nadu, India.

**Corresponding Author:** R.V. Bhosle, Department of Aquaculture, Fisheries College and Research Institute, Thoothukudi-628 008, Tamil Nadu, India. Email: bhosleramu330@gmail.com

**How to cite this article:** Bhosle, R.V., Kumar, J.S.S., Antony, C., Aanand, S., Kumar, V.S. and Lingam, R.S.S. (2022). Optimization of 17 $\alpha$ -methyl Testosterone Dose in *Artemia* Nauplii and Rotifer using Enrichment Technique. Indian Journal of Animal Research. DOI: 10.18805/IJAR.B-4800.

**Submitted:** 06-10-2021 **Accepted:** 02-02-2022 **Online:** 12-03-2022

suitable vectors -without leaching to the environment- may reduce the negative impacts of the hormone-based mono-sex seed production.

Alternative to commercial diet, live feed - a natural diet rich of nutrients - has gained an importance in hatcheries and started playing a pivotal role in larviculture. Among the various live feed, rotifer and *Artemia* are widely used due to their availability, low cost, ease of culture and nutritional biochemical composition (Kolkovski *et al.*, 1997; Sorgeloos and Lavens, 1996). Additionally, these live feed act as a very good carrier of supplying essential nutrients (PUFA) and hormones (androgens or estrogens) to the target animal through enrichment process.

Therefore, the best alternative option for supplementing the hormone for monosex seed production is live feed enrichment process. In commercial feed, a particular dose is selected as optimal dose for all male production, however, it is very difficult to achieve that particular dose in live feed. Therefore, it is essential to standardize the dose of hormone in live feed through enrichment process, before recommending them as potential carrier for hormone-based mono-sex seed production in future. Keeping this in mind, the study was aimed to optimize the 17 $\alpha$ -methyl testosterone hormone dose in *Artemia* nauplii and rotifer using enrichment process.

## MATERIALS AND METHODS

### Experimental site and design

The present experiment was conducted at the Krishnagiri Barur-Center for sustainable Aquaculture, Tamil Nadu Dr. J. Jayalalithaa Fisheries University, Tamil Nadu and India and research was carried out in summer 2021. The study consisted of five treatment doses of 17 $\alpha$  methyl testosterone namely; control (without hormone), T1 (20 mg L<sup>-1</sup>), T2 (40 mg L<sup>-1</sup>), T3 (60 mg L<sup>-1</sup>) and T4 (80 mg L<sup>-1</sup>) and each were triplicated. The doses were tested in *Artemia* nauplii and rotifer by following completely randomized design.

### Decapsulation of *Artemia* cyst

The cyst of *Artemia* strain (*Artemia salina*) was procured from Tuty Enterprises, Tamil Nadu, India. The procured cysts (Dynasty TM, Salt Lake City, UT, USA) were hatched out by following standard method (Sorgeloos *et al.* 1986) on daily basis. First, 3 g cyst were decapsulated using sodium hypochlorite solution (at 15 mL/g cyst) for 15 min, filtered through nitex screen (50  $\mu$ m) and washed with fresh water. To ensure complete removal of chlorine, the cysts were treated with 0.1% sodium thiosulphate. The washed cysts were then transferred to a conical *Artemia* hatching jar containing 4 L of seawater (30 ppt), with vigorous aeration, and maintained at 28°C for 24 h under 1000 lux light for hatching. After 24 h, the hatched-out larvae were harvested using 50-micron nitex screen and used for enrichment study.

### Rotifer culture

Pure culture of rotifer (*Brachionu s calyciflorus*), procured from live feed culture unit, Bharathidasan University, Tamil Nadu, India, was maintained at Krishnagiri-Barur Centre for Sustainable Aquaculture. First it was cultured in a 1000 ml flask and was periodically up-scaled to 500 litre Fibre-reinforced plastic (FRP) tanks for mass culture. The culture was maintained by feeding them with yeast @ 0.15 g/l and *Chlorella vulgaris* (1.5-2.5 $\times$ 10<sup>6</sup> cells/ml) during morning (08.00 AM) and evening (17.00 PM) times, respectively (Aker *et al.* 2013). Once, the culture density reached 250-450 individuals/ml, the rotifer were collected and used for enrichment process.

### Enrichment of *Artemia* nauplii and rotifer

The enrichment process was carried out in plastic buckets, each of 3 L capacity and provided with aeration and each

treatment were triplicated. The enrichment process of androgenic hormone, 17  $\alpha$ -methyl testosterone (MT), (Sigma Chemical Company, St Louis, MO, USA) in the *Artemia* and rotifer was carried out by following the standard method (Sorgeloos *et al.* 1986). The freshly hatched out *Artemia* nauplii and 6-8 days old rotifer were used for enrichment process. The hormone (17  $\alpha$ - MT) and ethanol mixture was prepared by following the method of (Stewart *et al.*, 2001) with slight modifications. Firstly, a stock solution of hormone-ethanol mixture (20000 ppm) was prepared. This stock solution of 20000 ppm was prepared by dissolving 1000 mg of 17  $\alpha$ -MT in 50 ml of 70% ethanol. 1 ml of this stock solution contains 20 mg of 17  $\alpha$ -MT which was further used for treatment doses preparation. A pre-determined volumes of stock solution such as 1 ml, 2 ml, 3 ml and 4 ml were added to individual replicate buckets to obtain the treatment doses of T1 (20mg L<sup>-1</sup>), T2 (40 mg L<sup>-1</sup>), T3 (60 mg L<sup>-1</sup>) and T4 (80 mg L<sup>-1</sup>), respectively. In control group, 2 ml of 70% ethanol without hormone was added. After the preparation of enrichment medium, rotifer (800 individuals/ml) and *Artemia* nauplii (100 individuals/ml) were introduced into the respective treatment medium and kept for different durations of enrichment.

### Sample collection, preparation and hormone analysis

After enrichment, samples were collected at different time intervals. The hormone-enriched *Artemia* nauplii and rotifer were harvested processed by following the standard method (Sandifer and Smith 1985). Samples of 10000 *Artemia* nauplii and rotifer taken from each replicate on 6, 12, 18, 24, 30, 36 and 42 hrs (for rotifer up to 24 hrs only collected). Testosterone levels were measured using enzyme-linked immunosorbent assay kit (ELISA) method. Testosterone was extracted using CALBIOTECH company kits following the manufacturer's instruction manual and ELISA reader (Model No is ER-180) was used (Jester *et al.*, 2014; Liu *et al.*, 2013; Jiang *et al.*, 2011).

### Statistical analysis

The collected data were statistically analysed using a statistical program SPSS version 28.0. Significant difference between the mean values were tested using repeated one-way analysis of variance by keeping dose and time as independent variables (each dose vs different time intervals; each time interval vs different doses). Tukey's test was used to rank the mean values and the statistical significance for the test was set as p>0.05.

## RESULTS AND DISCUSSION

In both live feed, the different durations enrichment found significantly higher 17  $\alpha$ -MT content at different doses. Similarly, *Daphnia magna* enriched with canola oil for different durations displayed significant difference in the lipid content between enriched and non-enriched daphnia (Fereidouni *et al.*, 2013). In *Artemia* nauplii, 17 $\alpha$ -MT was in the range of 0.35-0.50 mg L<sup>-1</sup> before the commencement of experiment (0<sup>th</sup> h). The study did not find any significant

variation in 17 $\alpha$ -MT hormone of *Artemia* kept in control group at different time intervals and it was in the range of 0.35-0.49 mg L<sup>-1</sup>. Similarly, no significant difference was found at 0<sup>th</sup> hour in the *Artemia* kept in different dose groups, however, as the time increased, *Artemia* started accumulating 17  $\alpha$ -MT in its tissues at different rates (Table 1). In general, a normal *Artemia* has lower quantity of steroids, for example 17  $\alpha$ -estradiol, than the enriched *Artemia* (Martin-Robichaud *et al.*, 1994) and it may be the reason for lower levels of 17  $\alpha$ -MT detection in the control group of the present study.

The increasing trend of 17 $\alpha$ -MT augmentation in *Artemia* was noticed up to 24 h. Significantly higher amount of 17 $\alpha$ -MT (15.33 $\pm$ 0.09 mg L<sup>-1</sup>) was observed in 80 mg L<sup>-1</sup> group at 24 hours (Table 1). Similarly, Martin-Robichaud *et al.*, (1994) reported that *Artemia* nauplii enriched in different concentrations of 17  $\alpha$ -estradiol was displayed significantly higher accumulation of 17 $\alpha$ -estradiol after 24 hours of enrichment. The study examined the assimilation of 17  $\alpha$ -MT at different time points, up to 42 hours, to know highest point of accumulation and starting point of withdrawal period. Interestingly, the study found maximum accumulation at 24 hours and, thereafter, a reduction in 17  $\alpha$ -MT assimilation has been noticed which may be due to higher metabolic activities and excretion activities (Navarro *et al.*, 1991; Dhert

*et al.*, 2001). Similarly, *Artemia* enriched with estradiol (E2) showed increasing trend and stable up to 24 hours (Contreras-Sánchez *et al.*, 2004). In an another study, when *Artemia* enriched with ascorbic acid (AA) at different durations found maximum absorption at 18 hrs and the AA content was started declining after 18 hrs of enrichment (Noshirvani *et al.*, 2006).

In rotifer, different doses and time intervals investigation found a significant difference in hormone accumulation. 17 $\alpha$ -MT levels in rotifer ranged from 0.37 to 0.47 mg L<sup>-1</sup> at 0<sup>th</sup> hr. The study did not find any significant differences in the 17  $\alpha$ -MT hormones of rotifer kept in the control group at various time intervals and it was in the range of 0.36-0.49 mg L<sup>-1</sup>. Similarly, there was no significant difference in the 17  $\alpha$ -MT level among the rotifers kept in different treatment doses at 0<sup>th</sup> hour.

At 6 hours, the level of 17-MT was significantly higher in all treatment groups. Further, the dose vs time analysis revealed that rotifer accumulated statistically higher quantum of 17 MT at 6 hrs than the 12 hrs (Table 2). After 6 hrs, a decreasing trend was noticed in 17 MT accumulations and significantly higher value of 17  $\alpha$ -MT (17.42 $\pm$ 0.06 mg L<sup>-1</sup>) was recorded at 6 hrs in 60 mg L<sup>-1</sup>. Previous studies on rotifer reported maximum accumulation of n-3 HUFA within 12 hrs of enrichment (Imada *et al.*, 1979; Watanabe *et al.*, 1983)

**Table 1:** *Artemia* enriched with different doses of 17  $\alpha$ -MT for different durations.

	0 h	6 h	12 h	18 h	24 h	30 h	36 h	42 h	P value (Dose *Time)
0 mg L <sup>-1</sup>	0.45 $\pm$ 0.06	0.41 $\pm$ 0.09 <sup>1</sup>	0.44 $\pm$ 0.02 <sup>1</sup>	0.49 $\pm$ 0.17 <sup>1</sup>	0.41 $\pm$ 0.015 <sup>1</sup>	0.43 $\pm$ 0.09 <sup>1</sup>	0.35 $\pm$ 0.05 <sup>1</sup>	0.38 $\pm$ 0.04 <sup>1</sup>	0.420
20 mg L <sup>-1</sup>	0.43 $\pm$ 0.05 <sup>a</sup>	1.35 $\pm$ 0.0.15 <sup>ab1</sup>	2.55 $\pm$ 0.15 <sup>b12</sup>	5.47 $\pm$ 0.22 <sup>cd2</sup>	7.48 $\pm$ 0.14 <sup>d2</sup>	7.42 $\pm$ 0.08 <sup>d2</sup>	4.25 $\pm$ 0.11 <sup>c2</sup>	3.60 $\pm$ 0.15 <sup>bc2</sup>	0.000
40 mg L <sup>-1</sup>	0.45 $\pm$ 0.01 <sup>a</sup>	1.97 $\pm$ 0.04 <sup>ab1</sup>	6.15 $\pm$ 0.65 <sup>bc2</sup>	8.10 $\pm$ 0.09 <sup>cd23</sup>	9.56 $\pm$ 0.02 <sup>d23</sup>	8.41 $\pm$ 0.03 <sup>cd2</sup>	7.54 $\pm$ 0.02 <sup>c3</sup>	4.65 $\pm$ 0.00 <sup>bc2</sup>	0.001
60 mg L <sup>-1</sup>	0.50 $\pm$ 0.01 <sup>a</sup>	8.13 $\pm$ 0.01 <sup>bc2</sup>	10.35 $\pm$ 0.08 <sup>cde3</sup>	11.06 $\pm$ 0.08 <sup>es34</sup>	12.19 $\pm$ 0.02 <sup>e3</sup>	11.30 $\pm$ 0.01 <sup>e3</sup>	9.69 $\pm$ 0.01 <sup>cd4</sup>	7.67 $\pm$ 0.15 <sup>b3</sup>	0.000
80 mg L <sup>-1</sup>	0.35 $\pm$ 0.01 <sup>a</sup>	9.53 $\pm$ 0.18 <sup>c2</sup>	12.60 $\pm$ 0.01 <sup>cd4</sup>	12.85 $\pm$ 0.04 <sup>d4</sup>	15.33 $\pm$ 0.09 <sup>e4</sup>	12.68 $\pm$ 0.05 <sup>d3</sup>	10.63 $\pm$ 0.01 <sup>cd4</sup>	8.20 $\pm$ 0.22 <sup>b3</sup>	0.000
P value	0.382	0.000	0.005	0.000	0.000	0.000	0.000	0.000	

In each row, mean values with different alphabet superscripts differ significantly at p<0.05, where each hormone dose was tested at different time intervals (dose\*time).

In each column, mean values with different numerical superscripts differ significantly at p<0.05, where different hormone doses were tested against each time interval (time\*dose).

**Table 2:** Rotifer enriched with different doses of 17  $\alpha$ -MT for different durations.

	0 h	6 h	12 h	18 h	24 h	P value (Dose*Time)
0 mg L <sup>-1</sup>	0.45 $\pm$ 0.08	0.49 $\pm$ 0.06 <sup>a1</sup>	0.36 $\pm$ 0.05 <sup>1</sup>	0.39 $\pm$ 0.09 <sup>1</sup>	0.38 $\pm$ 0.05 <sup>1</sup>	0.096
20 mg L <sup>-1</sup>	0.37 $\pm$ 0.04 <sup>a</sup>	14.82 $\pm$ 0.10 <sup>c2</sup>	14.04 $\pm$ 0.03 <sup>c23</sup>	11.04 $\pm$ 0.03 <sup>b2</sup>	10.30 $\pm$ 0.01 <sup>b2</sup>	0.000
40 mg L <sup>-1</sup>	0.47 $\pm$ 0.06 <sup>a</sup>	15.96 $\pm$ 0.07 <sup>c2</sup>	14.68 $\pm$ 0.02 <sup>c3</sup>	11.57 $\pm$ 0.09 <sup>b23</sup>	10.49 $\pm$ 0.11 <sup>b2</sup>	0.000
60 mg L <sup>-1</sup>	0.43 $\pm$ 0.05 <sup>a</sup>	17.42 $\pm$ 0.06 <sup>e3</sup>	15.68 $\pm$ 0.02 <sup>d3</sup>	12.59 $\pm$ 0.02 <sup>c3</sup>	11.13 $\pm$ 0.06 <sup>b2</sup>	0.000
80 mg L <sup>-1</sup>	0.40 $\pm$ 0.03 <sup>a</sup>	15.79 $\pm$ 0.03 <sup>d2</sup>	12.59 $\pm$ 0.06 <sup>c2</sup>	11.97 $\pm$ 0.06 <sup>bc23</sup>	10.16 $\pm$ 0.06 <sup>b2</sup>	0.000
P value	0.741	0.000	0.000	0.000	0.000	

In each row, mean values with different alphabet superscripts differ significantly at p<0.05, where each hormone dose was tested at different time intervals (dose\*time).

In each column, mean values with different numerical superscripts differ significantly at p<0.05, where different hormone doses were tested against each time interval (time\*dose).

and did not find any significant difference in fatty acid content when they enriched for longer duration, up to 24 hrs (Kotani *et al.*, 2010). Similarly, rotifer kept for 8 hrs in higher concentration of enrichment medium did not produce significantly higher quantity of fatty acid (Kotani *et al.*, 2010). At 6 hrs, there is no statistical difference in assimilation of 17  $\alpha$ -MT in the tissues of rotifers kept in 20 mg L<sup>-1</sup>, 40 mg L<sup>-1</sup> and 80 mg L<sup>-1</sup> treatment groups 9 (Table 2). Generally, in rotifer, it has been suggested to increase the duration of enrichment rather than quantity of enrichment medium to obtain more efficient enrichment (Rodriguez *et al.*, 1996; Yahyavi and Farzane, 2009). Interestingly, the present study found significantly higher assimilation of 17  $\alpha$ -MT in shorter enrichment period (in 6 hrs) which in turn may help to reduce cost associated with live feed production (Southgate and Lou, 1995).

## CONCLUSION

The harmful residual effect of hormones, released from the hormone incorporated diets, to the natural environment is alarming. As an alternate, supplementation of hormone via live feed, using enrichment techniques, could be a viable option which improves the larval ingestion rate, survival rate, growth rate and reduces hormone residual effects. The study found higher assimilation of 17  $\alpha$ -MT by *Artemia* (at 24 hours in 80 mg L<sup>-1</sup>) and rotifer (at 6 hours in 60 mg L<sup>-1</sup>) which could be further explored for mono-sex seed production in a more sustainable way.

## ACKNOWLEDGEMENT

The funding support by Tamil Nadu Dr. J. Jayalalithaa Fisheries University, Nagapattinam, and Tamil Nadu is gratefully acknowledged. We would like to express our sincere thanks to The Vice-Chancellor, Tamil Nadu Dr. J. Jayalalithaa Fisheries University, Nagapattinam, for his support and encouragement.

**Conflict of interest:** None.

## REFERENCES

- Akter, S., Shahjahan, M., Rahman, M.S., Das, P.S. (2013). Suitability of *Chlorella Ellipsoidea* as Food for Production of the Rotifer *Brachionus calyciflorus*. *International Journal of Agricultural Research, Innovation and Technology (IJARIT)*. 3: 41-48.
- Azizi-lalabadi, M., Pirsahab, M. (2021). Investigation of steroid hormone residues in fish: A systematic review. *Process Safety and Environmental Protection*. 152: 14-24.
- Bardhan, A., Sau, S.K., Khatua, S., Bera, M., Paul, B.N. (2021). A review on the production and culture techniques of *Monosex tilapia*. *International Journal of Current Microbiology and Applied Sciences*. 10: 565-577.
- Contreras-Sánchez, W.M., Hernández-Vidal, U., Álvarez González, C.A., Indy, J.R., Hernández-Franyutti, A., McDonal-Vera, A. (2009). Development of snook (*Centropomus Spp*) seed production technology for application in aquaculture and restocking of Over-Fished Populations. 1-61.
- Dhert, P., Rombaut, G., Suantika, G., Sorgeloos, P. (2001). Advancement of rotifer culture and manipulation Techniques in Europe. *Aquaculture*. 200: 129-146.
- Emeka, U., Iloegbunam, N.G., Gbekele-Oluwa, A.R. (2014). Natural products and aquaculture development. *Journal of Pharmacy and Biological Sciences*. 9: 70-82.
- Food and Agriculture Organization. (2020). The State of the World Fisheries and Aquaculture. Rome. FAO.1-224.
- Fereidouni, A.E., Fathi, N. Kholesi, M.K. (2013). Enrichment of *Daphnia magna* with canola oil and its effects on the growth, survival and stress resistance of the Caspian Kutum (*Rutilus frisii kutum*) larvae. *Turkish Journal of Fisheries and Aquatic Sciences*. 13: 1-15
- Gale, W.L., Fitzpatrick, M.S., Schreck, C.B. (1996). Masculinization of Nile Tilapia (*Oreochromis niloticus*) through immersion in 17 $\alpha$ -methyl testosterone. Thirteenth Annual Technical Report Oregon Cooperative Fishery Research Unit, Department of Fisheries and Wildlife, Oregon State University.
- Imada, O., Kageyama, Y., Watanabe, T., Kitajima, C., Fujita, S., Yone, Y. (1979). Development of a New Yeast as a Culture Medium for Living Feeds used in the Production of Fish Seed. *Bulletin of the Japanese Society of Scientific Fisheries*. 154-159.
- James, R., Sampath, K. (2006). Effect of dietary administration of methyltestosterone on the growth and sex reversal of two ornamental fish species. *Indian Journal of Fisheries*. 53: 283-290.
- Jester, E. L., Abraham, A., Wang, Y., Said, K.R.E., Plakas, S.M. (2014). Performance evaluation of commercial ELISA kits for screening of furazolidone and furaltadone residues in fish. *Food Chemistry*. 145: 593-598.
- Jiang J., Wang Z., Zhang H., Zhang X., Liu X., Wang, S. (2011). Monoclonal antibody-based ELISA and colloidal gold immunoassay for detecting 19-Nortestosterone residue in animal tissues. *Agricultural and Food Chemistry*. 59: 9763-9769.
- Kolkovski, S., Arieli, A., Tandler, A. (1997). Visual and chemical cues stimulate microdiet ingestion in sea bream larvae. *Aquaculture International*. 5: 527-536.
- Kotani, T., Genka, T., Tanabe, M., Miyashima, A., Fushimi, H., Hayashi, M. (2010). Effect of nutritional enrichment method on fatty acid contents of rotifer *Brachionus plicatilis*. *Journal of the World Aquaculture Society*. 46: 884-892.
- Lavens, P., Sorgeloos, P., (1996). Manual on the Production and Use of Live Food For Aquaculture. FAO Fisheries Technical Paper.
- Liu, Y.C., Jiang, W., Chen, Y.J., Xiao, Y., Shi, J.L., Qiao, Y.B., Zhang, H.J., Li, T., Wang, Q. (2013). A Novel Chemiluminescent ELISA for detecting furaltadone metabolite, 3-amino-5-Morpholinomethyl-2-Oxazolidone (AMOZ) in fish, egg, honey and shrimp samples. *Journal of Immunological Methods*. 395: 29-36.
- Martin-Robichaud, D.J., Peterson, R.H., Benfey, T.J., Crim, L.W. (1994). Direct feminization of lumpfish (*Cyclopterus lumpus*) Using 17 $\alpha$ -Oestradiol-Enriched Artemia as food. *Aquaculture*. 123: 137-151.

- Navarro, I., Canals, P., Sánchez, J., Gutiérrez, J., and Planas, J. (1991). Some Plasma Hormones and Metabolites in the Pyrenean Brown Trout (*Salmo trutta fario*). Comparative Biochemistry and physiology. A, Comparative Physiology. 100(4): 919-923.
- Navarro, J.F. (1996). The Structure of Cold Dark Matter Halos. In: Symposium-International Astronomical Union. Cambridge University Press. 171: 255-258.
- Noshirvani, M., Takami, A.G., Rassouli, A., Bokaei, S. (2006). The stability of ascorbic acid in artemia urmiana following enrichment and subsequent starvation. Journal of Applied Ichthyology. 22: 85-88.
- Pelissero, C., Sumpter, J. (1992). Steroids and "Steroid-Like" substances in fish diets. Aquaculture. 107: 283-301.
- Rodriguez, C., Pérez, J.A., Izquierdo, M.S., Cejas, J.R., Bolanos, A., Lorenzo, A. (1996). Improvement of the Nutritional Value of Rotifers by Varying the Type and Concentration of Oil and the Enrichment Period. Aquaculture. 147: 93-105.
- Sandifer, P.A. Smith, T.I.J. (1985). Freshwater Prawns. In: Crustacean and Mollusk Aquaculture in United States. 63-12.
- Sorgeloos P., Levens P., Leger, P., Tackaert, W., Versichele, D. (1986). Manual for the Culture and Use of Brine Shrimp Artemia in Aquaculture. State University of Ghent, Belgium.310.
- Southgate, P.C., Lou, D.C. (1995). Improving the n -3 HUFA composition of *Artemia* using microcapsules containing Marine Oils. Aquaculture. 134: 91-99.
- Stewart, A., Spicer, A., Inskeep, E., Dailey, R. (2001). Steroid Hormone Enrichment of *Artemia* Nauplii. Aquaculture. 202: 177-181.
- Watanabe, T., Kitajima, C., Fujita, S. (1983). Nutritional values of live organisms used in Japan for mass propagation of fish: A review. Aquaculture. 34: 115-143.
- Yahyavi, M.A. Z.I.Y. A.R., Farzane, M.R. (2009). Effects of concentration and duration of enrichment of rotifer *Brachionus plicatilis* with cod oil emulsion with emphasis on fatty acids. Iranian Scientific Fisheries Journal. 18: 153-164.