



# Heat Stress-induced Changes in Milk Composition and Physiological Responses of Lactating Kankrej Cow during Summer Season as Compared to Thermo-neutral Period

Nikita Bhalakiya<sup>1</sup>, Nilufar Haque<sup>2</sup>, Pankaj Patel<sup>3</sup>, Gaurav Kumar<sup>1</sup>, Deepesh Bharat Mishra<sup>4</sup>

10.18805/IJAR.B-4821

## ABSTRACT

**Background:** The Kankrej cattle is the heaviest and most priced of the Indian breeds of cattle. It is adapted to the geo-climatic conditions of North Gujarat and Kutch, which is normally hot and dry. It is reared in the district of Banaskantha in the state of Gujarat and also distributed in other parts of India and abroad. The experiment was carried out to study the effect of the summer season on the milk composition and physiological responses in lactating Kankrej cow compared to the thermoneutral period.

**Methods:** For the experiment, total number of 16 clinically healthy Kankrej cows were randomly selected from Livestock Research Station (LRS), Sardar Krushinagar Dantiwada Agricultural University, Banaskantha, Gujarat and categorized into two different groups according to seasons. Physiological responses *i.e.* respiration rate and rectal temperature were recorded and milk samples were collected from each animal during the summer and thermoneutral period at fortnight intervals. The milk samples were analyzed by using an automated milk analyzer.

**Result:** The results demonstrate that fat, SNF and density were significantly higher in the summer season. The physiological responses include respiration rate and rectal temperature were found significantly higher during summer than thermoneutral period. Milk protein was significantly higher during the thermo-neutral period as compared to summer.

**Key words:** Kankrej, Milk composition, Rectal temperature, Respiration rate, Summer stress, Thermo-neutral period.

## INTRODUCTION

Kankrej cow is a good milk producer and bullocks are very powerful, so it is suitable for draft purposes and well known for their "SAWAI CHAL means a particular walking style where the hind limbs of cow reach faster to the ground than the front limb which helps in faster plowing the fields. Due to this advantage, Kankrej bulls are used for road transportation. The population of Kankrej cattle was reviewed and found that its population was 5.0 lakh during 1951 and it has been increased to 30.28 lakh during 2013. It has been estimated that the average standard lactation milk yield of Kankrej cattle is 2523.33 litre, the average lactation length is 301.58 days and the calving interval is 443.72 days at the organized farm of Livestock Research Station, Sardar Krushinagar Dantiwada Agricultural University, Sardar Krushinagar, Gujarat.

Livestock keeping is an integral part of Indian society, which helps to provide nutritional security to rural people and opens the employment opportunities. However, most of the world's human and domestic animal population lies in regions where seasonal stressors adversely influence productivity. Stress is the condition where there is undue demand for physical and mental energy due to excessive and aversive environmental factors (stressors) and causes deformations identifiable through physiological disequilibrium. Seasonal stress imparts physical and economic losses to livestock production in temperate, subtropical and tropical regions. Each species, breed or animal with its physiological state has a comfort zone, in

<sup>1</sup>Division of Animal Physiology, ICAR-National Dairy of Research Institute, Karnal-132 001, Haryana, India.

<sup>2</sup>West Bengal University of Animal and Fishery Sciences, Kolkata -700 037, West Bengal, India.

<sup>3</sup>Department of Physiology and Biochemistry, College of Veterinary Science and Animal Husbandry, Sardarkrushinagar Dantiwada Agricultural University, Banaskantha-385 506, Gujarat, India.

<sup>4</sup>Division of Animal Nutrition, ICAR-National Dairy of Research Institute, Karnal-132 001, Haryana, India.

**Corresponding Author:** Gaurav Kumar, Animal Physiology Division, ICAR-National Dairy of Research Institute, Karnal-132 001, Haryana, India. Email: gauravyadav1725@gmail.com

**How to cite this article:** Bhalakiya, N., Haque, N., Patel, P., Kumar, G. and Mishra, D.B. (2022). Heat Stress-induced Changes in Milk Composition and Physiological Responses of Lactating Kankrej Cow during Summer Season as Compared to Thermo-neutral Period. Indian Journal of Animal Research. DOI: 10.18805/IJAR.B-4821.

**Submitted:** 01-11-2021 **Accepted:** 23-03-2022 **Online:** 31-05-2022

which the energy expenditure of the animal is minimal, constant and independent of environmental temperature. When environmental temperatures move out of the thermo-neutral zone (or comfort zone) during a different season, dairy cattle begin to experience heat stress in summer or cold stress in winter. These seasonal stresses negatively impact a variety of dairy parameters including milk yield and reproduction which created a significant financial burden in many dairy-producing areas of the world.

Under seasonal stress, many physiological and behavioral responses vary in intensity and duration in relation to the animal genetic makeup and environmental factors by integrating many organs and systems *viz.* behavioural, endocrine, cardio-respiratory and immune system Altan *et al.* (2003). Nonetheless, with the ever-increasing interest in indigenous livestock breeds as a possible solution to increase the efficiency of production in harsh conditions. There is an urgent need to study the seasonal stress-induced physiological and production changes in such indigenous breeds to plan and implement health and disease monitoring programs to ensure their sustainable development and economic viability. In addition, these parameters can be used to evaluate animal stress and welfare levels Anderson *et al.* (1999).

## MATERIALS AND METHODS

### Selection of animals

The experiment was conducted at Livestock Research Station, Sardar Krushinagar Dantiwada Agricultural University, Banaskantha, Gujarat. A total number of 16 clinically healthy animals were selected which having second to fifth parity with average body weight around 430 kg. All the cows were clinically healthy, maintained under standard management conditions and allowed to take ad-libitum food. Prior approval of the Institutional Animal Ethics Committee (IAEC) was taken before conducting an experiment.

### Experimental design

An experiment was carried out in two periods according to the temperature and relative humidity recorded: From May 2018 to June 2018 (maximum temperature-42.1 and maximum relative humidity-77, minimum temperature-23.9 and minimum relative humidity-50) for the summer season and from February 2019 to March 2019 (maximum temperature-36.5 and maximum relative humidity-68, minimum temperature-11.8 and minimum relative humidity-38) for a thermo-neutral period. During each season, eight animals were selected (n=8).

### Milk sampling

Milk samples were collected during each season at fortnight intervals from animals to measure milk fat, milk protein, lactose, SNF and density. These milk parameters were measured by Automated Milk Analyzer (Ekomilk, Milk Analyzer Milkana KAM98-2A, Ultra pro). Physiological parameters *viz* rectal temperature (RT, °F) and respiration rate (RR, breaths/min) were recorded by Digital thermometer (N and B Medical Products Co., New Delhi) and by observation of flank movement respectively at fortnight intervals during the experimental periods. Respiration rate and rectal temperature were taken in the morning (9 a.m.) and milk samples were collected in the afternoon (4 p.m.) on the same day of physiological parameters recorded.

### Statistical analysis

The data were analyzed by applying one way ANOVA test

using the Sigma stat 32 software package.  $P < 0.05$  was considered to be statistically significant.

## RESULTS AND DISCUSSION

### Rectal temperature (°F)

The recorded mean and standard error (mean $\pm$ S.E.) values of rectal temperature (°F) during different seasons in lactating Kankrej cow were presented in the Table 1 and Fig 1. It was observed that rectal temperature was significantly ( $p < 0.05$ ) higher in the summer season than thermoneutral period. The mean rectal temperature determined in the present study corroborated the values of Haque *et al.* (2012) and Yadav *et al.* (2016); who reported higher RT during summer months in buffaloes and cattles respectively. The core body temperature of cows in thermo-neutral conditions is maintained by the thermoregulation system of the animal's body with a range of about 1°C (between 38 to 39.2°C) Ammer *et al.* (2016). Under these ambient conditions, the exchange of animal heat within the animal (across cellular and vascular membranes) and between the animal and its environment are kept in balance; however, it is always a dynamic equilibrium Taylor *et al.* (2014). During periods of excess heat (heat stress), homeostatic mechanisms of the body are activated to re-establish the internal environment's thermal status or regulate it within acceptable physiological limits Werner *et al.* (2008). During the study, the higher rectal temperature in summer indicates that the animals could not dissipate the excess heat by usual means (conduction, convection, evaporation), which increased body temperature. A similar increase in the RT at higher temperatures/heat stress have been reported in cattle by Marai *et al.* (1995); and Nessim (2004) in buffalo, Korde *et al.* (2007) and Krishnan *et al.* (2009). Bewley *et al.* (2008) mentioned that the rectal temperature is directly affected by the ambient temperature and increasing during warmer weather. Rectal temperature is an indicator of thermal balance and may be used to assess the adversity of the thermal environment, which can affect the growth, lactation and reproduction of dairy cows Hansen and Arechiga (1999) and West (1999).

### Respiration rate (Breaths/min)



The recorded mean and standard error of respiration rate (Breaths/min) in lactating Kankrej cow was shown in Table 1 and Fig 2. It was observed that the respiration rate was significantly ( $p < 0.05$ ) higher during the summer season in compared to the thermoneutral period. The mean respiration rate determined in the present study corroborates the values previously reported by Haque *et al.* (2012). They reported a non-significant difference at 42°C and 45°C in the respiration rate in adult animals, except in the case of 45°C, a significant higher ( $P < 0.05$ ) respiration rate was observed in young animals than adults. The increase in respiration rate with the increasing temperature may be due to the more demand for oxygen by the tissues in stressful conditions. A similar increase in the respiration rate at higher temperatures/heat

**Table 1:** Physiological parameters in Lactating Kankrej cow during Summer and Thermoneutral periode.

Seasons	Season-I Summer (May-June)	Season-III Thermoneutral (February-March)
Parameters	Temp-Min-23.9, Max-42.1 RH-Min-50, Max-77	Temp-Min-11.8, Max-36.5 RH-Min-38, Max-68
Rectal temperature (°F)	100.91±0.21 <sup>b</sup>	99.90±0.06 <sup>a</sup>
Respiration rate (breaths/min )	29.00±0.62 <sup>b</sup>	20.12±0.36 <sup>a</sup>

\*Means with different superscripts (a, b) within a row differ significantly from each other.

\*(Source of Temperature (Temp) and Relative Humidity (RH) data- Department of Meteorology, Chimanbhai Patel College of Agriculture, Sardar Krushinagar Dantiwada Agriculture University, Gujarat).

**Fig 1:** Rectal temperature during different seasons in lactating Kankrej cow.**Fig 2:** Respiration rate during different seasons in lactating Kankrej cow.

stress have been reported by Gaughan *et al.* (2000), Nonaka *et al.* (2008) and Pereira *et al.* (2008) in cattle and Korde *et al.* (2007) and Zhang *et al.* (2010) in buffaloes.

The respiratory rate (RR) was highly affected by ambient air temperature changes because the respiratory rate is considered one of the major physiological reactions that can keep body temperature within the normal range Frangiadaki *et al.* (2003). McLean (1963) observed the significance of the increase in respiration rate under heat stress and

enables the animal to dissipate the excess of body heat by vaporizing more moisture in the expired air, which accounts for about 30% of the total dissipation. On the other hand, McDowell (1972) and Gaughan *et al.* (1999) observed contradictory results; they reported a low respiratory rate under hot weather and identified animals with lesser discomfort. This fact is evident when comparing respiration rates of *Bos taurus* versus *Bos indicus* under hot summer weather conditions where *Bos indicus* (Zebu) cattle maintain

lower respiration rates. There was a significant ( $p<0.05$ ) increase in respiration rate to maintain body temperature in the present study. Still, animals could not maintain their body temperature and there was a significant ( $p<0.05$ ) increase in rectal temperature, which indicates heat stress felt by animals during the extreme summer season.

### Milk composition

#### Fat (%)

Hot and humid environment not only affects milk yield but also affects milk quality. It was observed that the values of the fat content (%) in milk during the summer season and thermoneutral period were depicted in Table 2 and Fig 3. Results showed that the fat content of milk during the summer season was significantly ( $p<0.05$ ) higher than the thermoneutral period. An increase in fat% might be due to an increase in roughage intake during the summer season, leading to higher acetate formation responsible for the increase in milk fat%. Similar result were also reported by Bernadin (1972), Sharma *et al.* (2001), Prasad (2009), Verma *et al.* (2018), Salim Bahashwan (2014) and Sarkar *et al.* (2006) that milk composition traits were highest in hot humid season but lowest in compared to other seasons.

The daily yield of fat was observed to be higher during summer ( $p<0.05$ ) than winter season. Contradiction to result Kadzere *et al.* (2002) and Bouraoui *et al.* (2002) observed lower milk fat in the summer season. When THI value goes beyond 72, milk fat content declines. Zheng *et al.* (2009) observed that the higher environmental temperature significantly reduces the percentage of milk fat in milk.

#### Protein (%)


The milk protein values during the summer season and thermoneutral period were presented in Table 2 and Fig 4. It was observed that milk protein was significantly ( $p<0.05$ ) higher during a thermoneutral period than the summer season. The result was similar to Zheng *et al.* (2009) observed that heat stress significantly reduces the production of milk and the percentage of protein in milk. Bouraoui *et al.* (2002) observed lower milk protein in the summer season. When THI value goes beyond 72, milk protein content declines. Heat stress significantly reduced milk protein contents from 3.20% during the winter season to 3.07% during summer season (Gaafar *et al.* 2011). While Radhika *et al.* (2012) reported that, the protein content was significantly lower ( $p<0.05$ ) during winter than summer season.

**Table 2:** Milk composition during different seasons in lactating Kankrej cow.

Parameters	Seasons	Season-I Summer (May-June) Temp-Min-23.9, Max-42.1 RH-Min-50, Max-77	Season-II Thermoneutral (February-March) Temp-Min-11.8, Max-36.5 RH-Min-38, Max-68
Fat		3.56 $\pm$ 0.17 <sup>b</sup>	2.80 $\pm$ 0.23 <sup>a</sup>
Milk protein		2.47 $\pm$ 0.04 <sup>b</sup>	3.36 $\pm$ 0.03 <sup>a</sup>
Lactose		4.18 $\pm$ 0.06 <sup>a</sup>	4.50 $\pm$ 0.04 <sup>a</sup>
SNF		9.60 $\pm$ 0.10 <sup>b</sup>	8.47 $\pm$ 0.07 <sup>a</sup>
Density		38.29 $\pm$ 0.44 <sup>b</sup>	30.63 $\pm$ 0.38 <sup>a</sup>

Means with different superscripts (a, b) within a row differ significantly from each other.

(Source of Temperature (Temp) and Relative Humidity (RH) data- Department of Meteorology, Chimanbhai Patel College of Agriculture, Sardar Krushinagar Dantiwada Agriculture University, Sardar Krushinagar, Gujarat).



**Fig 3:** Milk fat (%) during different seasons in lactating Kankrej cow.




Fig 4: Milk protein level (%) during different seasons in lactating Kankrej cow.




Fig 5: Lactose concentration during different seasons in lactating Kankrej cow.

#### Lactose (%)

It was observed that the values of lactose during the summer season and thermoneutral period were shown in Table 2 and Fig 5. No significant change was found in lactose content during the summer season and thermoneutral season. The result was similar to Zheng *et al.* (2009) observed that the higher environmental temperature has no significant effect on the lactose content in the milk. Contraindication to result of Gaafar *et al.* (2011) reported that the heat stress significantly reduced milk lactose contents from 4.78% during the winter season to 4.59% during summer season.

#### Solid not fat (SNF%)

It was observed that the values of the solid not fat in milk during the summer season and thermoneutral period were presented in Table 2 and Fig 6. It was observed that solid not fat of milk was significantly ( $p < 0.05$ ) higher during the summer season than the thermoneutral

period. The result was contraindicated to Gaafar *et al.* (2011) who claims that higher temperature significantly reduced the milk SNF contents from 8.69% during the winter season to 8.34% during summer season. Similar results were also reported by Bernadin (1972) and Verma *et al.* (2018).

#### Density

It was observed that the values of milk density during the summer season and thermoneutral period were presented in Table 2 and Fig 7. It was observed that the density of milk was significantly ( $p < 0.05$ ) higher during the summer season as compared to the thermoneutral period.

In the present study, there were a significant ( $p < 0.05$ ) increase in fat%, concentration, solid not fat and density found during the summer season, while there was a significant ( $p < 0.05$ ) increase in milk protein level during the thermoneutral period.




Fig 6: Solid not fat (%) during different seasons in lactating Kankrej cow.




Fig 7: Density of milk during different seasons in lactating Kankrej cow.

## CONCLUSION

In present study, during the thermoneutral period the values of physiological responses and milk composition were found within the normal range so it may be considered as the physiologically most comfortable season for lactating Kankrej cow.

**Conflict of interest:** None.

## REFERENCES

- Altan, O., Pabuccuoglu, A., Altan, A., Konyalioglu, S. Bayraktar, H. (2003). Effect of heat stress on oxidative stress, lipid peroxidation and some stress parameters in broilers. *British Poultry Science*. 44(4): 545-550.
- Ammer, S., Lambert, C., Gauly, M. (2016). Comparison of different measuring methods for body temperature in lactating cows under different climatic conditions. *Journal of Dairy Research*. 83(2): 165-172.
- Anderson, B.H., Watson, D.L., Colditz, I.G. (1999). The effect of dexamethasone on some immunological parameters in cattle. *Veterinary Research Communication*. 23(7): 399-413.
- Bahashwan S. (2014). Effect of cold and hot season on fat, protein and lactose of Dhofari cows milk. *Net Journal of Agriculture Science*. 2(1): 47-49.
- Bernadin B. (1972). Seasonal variation in the composition and physicochemical and biological proportion of milk. *Dairy Sci. Abstr.* 34(7):207.
- Bewley, J.M., Einstein, M.E., Grott, M.W., Schutz, M.M. (2008). Comparison of reticular and rectal core body temperatures in lactating dairy cows. *Journal of Dairy Science*. 91(12): 4661-4672.
- Bouraoui, R., Lahmar, M., Majdoub, A., Djemali, M. and Belyea, R. (2002). The relationship of temperature-humidity index with milk production of dairy cows in a Mediterranean climate. *Anim. Res.* 51(6): 479-491.
- Frangiadaki, E., Golidi, E., Menegatos, I., Luzi, F. (2003). Comparison of does' performances under high and moderate temperature in a Greek commercial farm. *World Rabbit Science*. 11(3): 137-143.
- Gaafar, H.M.A., Gendy, M.E., Bassiouni, M.I., Shamiah, S.M., Halawa, A.A. and Hamd, M.A. (2011). Effect of heat stress on performance of dairy Friesian cow's milk production and composition. *Researcher*. 3(5): 85-93.



- Gaughan, J.B., Holt, S.M., Hahn, G.L., Mader, T.L., Eigenberg, R. (2000). Respiration rate-is it a good measure of heat stress in cattle? *Asian-Australasian Journal of Animal Sciences*. 13: 329-332.
- Gaughan, J.B., Mader, T.L., Holt, S.M., Josey, M.J., Rowan, K.J. (1999). Heat tolerance of Boran and Tuli crossbred steers. *Journal of Animal Science*. 77(9): 2398-2405.
- Hansen, J.R.S., Arechiga, C.F. (1999) Strategies for Managing Reproduction in the Heat Stressed Dairy Cow. *Journal of Dairy Science*. 77: 36-50.
- Haque, N., Ludri, A., Hossain, S.A., Ashutosh, M. (2012). Alteration of metabolic profiles in young and adult Murrah buffaloes exposed to acute heat stress *International Journal of Applied Animal Science*. 1(1): 23-29.
- Kadzere, C.T., Murphy, M.R., Silanikove, N. and Maltz, E. (2002). Heat stress in lactating dairy cows: A review. *Livest. Prod. Sci.* 77(1): 59-91.
- Korde, J.P., Singh, G., Varshney, V.P., Shukla, D.C. (2007). Effects of Long-term Heat Exposure on Adaptive Mechanism of Blood Acid-base in Buffalo Calves. *Asian-Australasian Journal of Animal Sciences*. 13: 329-332.
- Krishnan, G., Singh, G., Shukla, D.C. (2009). The Effect of Electrolyte Supplementation on Physiological Responses in Heat Stressed Male Buffalo Calves. *Indian Journal of Animal Science*. 79(1): 34-37.
- Marai, I.F.M., Habeeb, A.A.M., Daader, A.H., Yousef, H.M. (1995). Effect of Egyptian subtropical conditions and the heat stress alleviation techniques of water spray and diaphoretics on the growth and physiological functions of Friesian calves. *Journal of Arid Environment*. 30(2): 219-225.
- McDowell, R.E., Jones, R.G., Pant, H.C., Roy, A., Siegenthaler, E.J., Stouffer, J.R. (1972). Improvement of livestock production in warm climates. WH Freeman and Company, San Francisco.
- McLean, J.A. (1963). Measurement of cutaneous moisture vaporization from cattle by ventilated capsules. *The Journal of Physiology*. 167(3): 417-426.
- Nessim, M.G. (2004). Heat-induced biological changes as heat tolerance indices related to growth performance in buffaloes. Ph.D. Thesis (Unpublished). Ain Shams University, Cairo, Egypt.
- Nonaka, I., Takusari, N., Tajima, K., Suzuki, H., Higuchi, T., Kurihara, K.M. (2008). Effects of high environmental temperatures on physiological and nutritional status of prepubertal Holstein heifers. *Livestock Science*. 113: 14-23.
- Pereira, A.M.F., Baccari, F., Titto, E.A.L., Afonso, J.A. (2008). Effect of thermal stress on physiological parameters, feed intake and plasma thyroid hormones concentration in Alentejana, Mertolenga, Frisian and Limousine cattle breeds. *International Journal of Biometeorology*. 52: 199-208.
- Prasad, J. (2009). *Animal Husbandry and dairy Science* kalyani publisher Ludhiana.
- Radhika, G., Ajithkumar, S., Rani, A., Sathian, C.T., Anilkumar, K., Usha, A.P., Dinesh, C.N. (2012). Milk yield and composition of crossbred cows in the hilly Wayanad district of Kerala, India. *Indian J. Animal Sci.* 82: 1251-1254.
- Sarkar, U., Gupta, A.K., Sarkar, V., Mohanty, T.K., Raina, V.S., Prasad, S. (2006). Factors affecting test day milk yield and milk composition in dairy animals. *J. Dairying Foods HS*. 25: 129-132.
- Sharma, R.B., Kumar, M., Pathak, V. (2001). Effect of different seasons on cross-bred cow milk composition and paneer yield in sub-Himalayan region. Department of Livestock Products Technology, College of Veterinary and Animal Sciences CSK-HPKV, Palampur (H.P.), India.
- Taylor, N.A., Tipton, M.J., Kenny, G.P. (2014). Considerations for the measurement of core, skin and mean body temperatures. *Journal of Thermal Biology*. 46: 72-101.
- Verma, S., Singh, R.P. (2018). Effect of season on compositional quality of Gangatiri cow milk. *Journal of Pharmacognosy and Phytochemistry*. 7(2): 3813-3814.
- Werner, J., Mekjavic, I.B., Taylor, N.A.S. (2008). Concepts in Physiological Regulation: A Ther-moregulatory Perspective. In: *Physiological Bases of Human Performance during Work and Exercise*. [Taylor N.A.S., Groeller, H. (eds)], Churchill Livingstone, London, United Kingdom. 325-340.
- West, J.W., Hill, G.M., Fernandez, J.M., Mandebvu, P., Mullinix, B.G. (1999). Effects of dietary fiber on intake, milk yield and digestion by lactating dairy cows during cool or hot, humid weather. *Journal of Dairy Science*. 82: 2455-2465.
- Yadav, B., Pandey, V., Yadav, S., Singh, Y., Kumar, V., Sirohi, R. (2016). Effect of misting and wallowing cooling systems on milk yield, blood and physiological variable during heat stress in lactating Murrah buffalo. *Animal Science Technology*. 58: 2.
- Zhang, R., Wu, X., Bao, R., Liu, Q. (2010). Analysis on Physiological and Anti oxidative Indices of Fu'an Buffalo in Heat Stress and Non-heat Stress Seasons. *Acta Agriculturae Jiangxi*.
- Zheng, L., Chen, M. and Zhi-Cheng, G. (2009). Effects of heat stress on milk performance and fatty acids in milk fat of Holstein dairy cows. *J. Chin. Dairy Ind.* 37(9): 17-19.