



Genetic Diversity of Crossbred Cattle using Cytochrome Oxidase Subunit I (COI) Gene in South Sulawesi, Indonesia

Sri Utami^{1,2}, Ali Jamil³, Reviany V. Nidom^{4,5}, Muhlis Natsir⁶, Haris Prayitno⁶, Dhani Prakoso⁴, Arif N.M. Ansori⁴, Setyarina Indrasari^{4,7}, Chairul A. Nidom^{2,4,5,7}

10.18805/IJAR.BF-1600

ABSTRACT

Background: The genetic diversity information of local cattle and crossbred in South Sulawesi, Indonesia is essential for the establishment of conservation and breeding strategies. This research targeted to evaluate the phylogenetic relationship and genetic diversity of local and crossbred (CB) cattle.

Methods: The present research organized at the Animal Laboratory, Professor Nidom Foundation, Surabaya, Indonesia from February to October 2021. The genome was isolated from 20 whole blood samples from seven purebred belongs to Singosari National Artificial Insemination Center Malang, Indonesia, four crossbred in Gowa and nine crossbred in Sidrap, South Sulawesi, Indonesia. Partial sequences of mtDNA cytochrome oxidase subunit I (COI) gene were amplified employing the polymerase chain reaction technique.

Result: This study revealed that a lot of single nucleotide polymorphism (SNP) found in local cattle (Bali and Peranakan Ongole) and CB cattle in Gowa and Sidrap indicated various genetic diversity can be used as a marker candidate and potential to be developed into future research knowing the correlation of the SNP with the growth rate of CB. Thus, this information will be helpful for further strategy in breeding development program.

Key words: COI, Crossbreed, Genetic diversity, Multidimensional scaling, Phylogenetics.

INTRODUCTION

Since livestock producers in Indonesia rely heavily on cattle for their revenue, cattle are one of the most essential livestock products (Dakhlan *et al.*, 2022; Kuswati *et al.*, 2022; Pakpahan *et al.*, 2022; Prihandini *et al.*, 2022; Sudrajad *et al.*, 2022). The productivity of Indonesian livestock, especially local cattle, is still relatively low and is a major problem that must be resolved immediately (Mayulu *et al.*, 2010). The program to increase the local cattle population is not yet optimal due to the extensive traditional rearing system, high production of livestock slaughter, limited feed, shrinking grazing area and decreased genetic quality (Kumar *et al.*, 2017a; Pundir *et al.*, 2019). The productivity of beef cattle can be increased through genetic and environmental quality improvements, in its application through a breeding program (Agung *et al.*, 2019; Jobirov *et al.*, 2022; Kibona *et al.*, 2022). Moreover, some breeders' cross local cattle with Limousine semen or Simmental, Brahman to get a breed of cattle that have a better production performance than the local cattle and to get the proceeds of sale the cattle is high. In fact, the maintenance of crossbred also has constraints in reproduction for low heifers compared to local cows (Malau-Aduli *et al.*, 2021).

The genetic diversity of native cattle breeds has recently been challenged by a lack of information on other cattle breeds, the development of native and local cow breeds, particularly in South Sulawesi, Indonesia and the import of exotic breeds (Sari *et al.*, 2016; Sutarno and Setyawan, 2016). Making successful conservation plans is challenging due to their existing genetic variety, prevalence of inbreeding and genetic blood combination. Major risks include genetic erosion

¹Agriculture Quarantine Major Service of Surabaya, Ministry of Agriculture, Surabaya, Indonesia.

²Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia.

³Directorate General of Facilities and Infrastructure, Ministry of Agriculture, Jakarta, Indonesia.

⁴Professor Nidom Foundation, Surabaya, Indonesia.

⁵Riset AIRC Indonesia, Surabaya, Indonesia.

⁶Agriculture Quarantine of Jayapura, Jayapura, Indonesia.

⁷PNF-Animal Diagnostic-Laboratory, Surabaya, Indonesia.

Corresponding Author: Chairul A. Nidom, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia.
Email: nidomca@fkh.unair.ac.id

How to cite this article: Utami, S., Jamil, A., Nidom, R.V., Nasir, M., Prayitno, H., Prakoso, D., Ansori, A.N.M., Indrasari, S. and Nidom, C.A. (2022). Genetic Diversity of Crossbred Cattle Using Cytochrome Oxidase Subunit I (COI) Gene in South Sulawesi, Indonesia. Indian Journal of Animal Research. doi: 10.18805/IJAR.BF-1600.

Submitted: 21-10-2022 **Accepted:** 29-12-2022 **Online:** 01-02-2023

and the loss of breed-specific genetic variation. It is commonly recognized that local cattle, such as those from Bali, have a significant number of favorable features, including the capacity to withstand hot weather conditions, poor fodder quality and resistance to internal parasites and infectious illnesses (Mohamad *et al.*, 2012; Bakae *et al.*, 2022; Liu *et al.*, 2022; Trujano-Chavez *et al.*, 2022; Xu *et al.*, 2020).

The mtDNA COI is a gene that is often used as a DNA barcode to analyze the diversity of genetic in cattle (Hidayati

et al., 2016). This gene was also used to reconstruct phylogenetics at the species level of evolution (Cai and Ma, 2016). Regarding the above points, thus, explored the COI gene to assess the genetic diversity and phylogenetic relationships of the local cattle and CB populations in Gowa and Sidrap District, Indonesia. Supplying information of genetic in crossbreds is vital for future breeding programs in order to support policies and programs to elevate the quality of beef cattle in South Sulawesi Province, Indonesia.

MATERIALS AND METHODS

Blood samples collection and DNA extraction

This study organized at the Animal Laboratory, Professor Nidom Foundation, Surabaya, Indonesia from February to October 2021. This study approved by the Institutional Animal Care and Use Committee - Professor Nidom Foundation (IACUC-PNF), Surabaya, Indonesia (Number: 010221/IACUC/VII/2021). Twenty blood samples were isolated from three regions representing seven purebred cattle belongs to Singosari National Artificial Insemination Center, Malang, Indonesia and four CB cattle from Gowa

District and nine CB cattle from Sidrap, South Sulawesi, Indonesia (Fig 1 and Table 1). Samples isolated from jugular vein using 3 mL syringe. Those blood samples were kept in room temperature and preserved in EDTA for laboratory analysis. The DNA was extracted using gSYNC™ DNA extraction kit and stored at -20°C before further analysis.

Quantitative and qualitative test of DNA isolation

1.5% agarose gel electrophoresis in 1XTAE solution was used for a quantitative and qualitative evaluation of the extracted DNA. For 40 minutes, electrophoresis was carried out at 100 V. By using spectrophotometry on all samples, purity level measurement and concentration of the extracted DNA are achieved (Ansori *et al.*, 2021).

PCR amplification

Gene COI HCO 2198 and LCO 1490 was amplified employing two primers: Forward 5' TAAACTTCAGGGT GACCAAAAAATCA 3' and Reverse 5' GGTCAAATCATA AAGA TATTGG 3' referred to Poniente *et al.* (2022). The amplification conditions were as follows initial denaturation (95°C for 5 mins), denaturation (95°C for 45 sec with 35 cycles), annealing (60°C for 45 sec), extension (72°C for 60 sec) and final extension (72°C for 5 mins). The PCR products were sequenced employing ABI 310 xL Genetic Analyser (Applied Biosystems, Inc) referred to Ansori *et al.* (2021) and Prihandini *et al.* (2022).

Genetic, phylogenetic tree and statistical analysis

All samples (mtDNA COI gene sequences) were aligned using the ClustalW. MEGA 11 software was used to estimate the COI gene diversity metrics, such as the number of polymorphic sites genetic distances (Kimura two-parameter model algorithm) and the resulting distance matrices were used to create a neighbor-joining (NJ) tree (1000 bootstrapping repeats) (Ansori *et al.*, 2021; Naseer *et al.*, 2018; Tamura *et al.*, 2021). In the present research, two samples of *Bos taurus* (GenBank Accession Number: HM.102289.1 and HQ.860420.1) were used in the

Table 1: Samples of purebreed and CB cattle.

Purebreed	CB cattle gowa district	CB cattle sidrap district
S1 : Ongole	Sp A : Lim - PO	Sp1 : Lim - Bali
S2 : Bali	Sp B : Angus - PO	Sp2 : Sim - Bali
S3 : Limousine	Sp E : Brah - PO	Sp5 : Bali - Lim - Lim
S4 : Madura	Sp F : Sim - PO	Sp6 : Bali - Lim - Brah
S5 : Angus		Sp7 : Bali - Brah
S6 : Brahman		Sp8: Bali-Lim-Madura
S7 : Sim		Sp9 : Bali - Brah - Sim
		Sp10 : Bali - Madura
		Sp11: Bali - Sim - Lim

Note: Lim: Limousine; PO: Peranakang ongole; Sim: Simmental; Brah: Brahman.




Fig 1: Sampling location of this study.

to 0.297. The result was in accordance with Kumar *et al.* (2017b) who stated that the genetic diversity of COI gene in the Bovidae was between 0.0 and 1.92%. In this study showed that seven purebred cattle might be grouped into *B. taurus* because it had a genetic distance of 0.000 (Angus, Brahman and Simmental), while Limousine has a genetic distance of 0.002 and Madura with a value of 0.013. Bali and PO cattle have a longer genetic distance were 0.055 and 0.095. Among the CB (Angus-PO) is very far to the GenBank reference (0.248) and other CB and purebred. The closest genetic distance of CB (Bali-Brahman) and (Bali-Madura) is 0.077 each. We revealed that most of CB cattle should be grouped into *B. indicus* (Bali and PO) because it had a genetic distance closer than to *B. taurus*.

A Neighbor-joining (NJ) tree was generated in order to verify the phylogenetic relationship of 20 of the COI gene using MEGA 11 based on nucleotide diversity (Fig 5). The result showed that all these samples were grouped into three main clades (A, B and C) and outgroup (D). Clade A is consist of CB cattle (Angus-PO) with bootstrap 94%. CB cattle (Angus-PO) which separated from other groups of CB cattle.

B clade was supported by 100% bootstrap value and consist purebred cattle Angus, Brahman and Simmental with the closest genetic distance was 0.00 and 100% homolog to *B. taurus*, while Limousine has a genetic distance of 0.002 and Madura with a value of 0.013. Cluster C has a bootstrap value of 48%, the members of this cluster are mostly CB cattle except PO and Bali. The genetic distance of PO and Balinese cows is 0.055 and 0.095, respectively. Considering the result, based on the DNA barcode, most of CB cattle were in the same cluster with *B. indicus* as well as Bali and PO cattle.

Statistical analysis

The result of Pearson's correlation analysis could be shown on Table 2 and Table 3. Results of Pearson's correlation

Table 2: Pearson's corellation analysis of nucleotide (snp) to hematology and blood chemistry description.

Variable	Correlation to SNP	P value
SNP (Nucleotide)	1	
WBC	0.542	0.017*
RBC	-0.650	0.003**
HGB	-0.336	0.160
HCT	-0.619	0.005**
PLT	0.336	0.123
MCV	0.145	0.554
MCHC	0.541	0.017*
MCH	0.617	0.005**
Total protein	-0.211	0.386
Albumin	-0.407	0.084
Globulin	-0.076	0.757
SGOT	0.284	0.239
SGPT	-0.129	0.598
ALP	-0.088	0.720
Total cholesterol	-0.069	0.784

Description: *Significant (p-value <0.05). **Very Significant (p-value <0.01). RBC- Red blood cell, WBC- White blood cell, HGB- Hemoglobin, HCT- hematokrit, PLT- Platelets, MCV- Mean corpuscular volume, MCH- Mean corpuscular hemoglobin, MCHC- Mean corpuscular hemoglobin concentration. SGOT- Serum glutamic oxaloacetic transaminase, SGPT- Serum glutamic pyruvate transaminase, ALP- Alkaline phosphatase.

Table 3: Pearson's correlation analysis of amino acid (snp) to hematology and blood chemistry description.

Variable	Correlation to SNP	P value
SNP (Nucleotide)	1	
WBC	0.479	0.033*
RBC	-0.702	0.001**
HGB	-0.590	0.006**
HCT	-0.619	0.005**
PLT	0.310	0.183
MCV	-0.005	0.982
MCHC	0.479	0.032*
MCH	0.263	0.282
Total protein	-0.202	0.394
Albumin	-0.545	0.013*
Globulin	-0.007	0.977
SGOT	0.209	0.377
SGPT	-0.006	0.981
ALP	-0.081	0.735
Total cholesterol	-0.232	0.339

Description: *Significant (p-value <0.05). **Very Significant (p-value <0.01). RBC- Red blood cell, WBC- White blood cell, HGB- Hemoglobin, HCT- hematokrit, PLT- Platelets, MCV- Mean corpuscular volume, MCH- Mean corpuscular hemoglobin, MCHC- Mean corpuscular hemoglobin concentration. SGOT- Serum glutamic oxaloacetic transaminase, SGPT- Serum glutamic pyruvate transaminase, ALP- Alkaline phosphatase.




Fig 5: Phylogenetic tree purebreed and CB cattle.

analysis that was illustrated in Table 2 showed that there was a very significant (p -value <0.01) and significant (p -value <0.05) correlation between SNP (nucleotide) and several hematological parameters. These hematological parameters were WBC, RBC, HCT, MCH and MCHC. Three parameters had a very significant correlation to SNP, namely RBC (0.003), HCT (0.005) and MCHC (0.005). Other parameters had a significant correlation to SNP were WBC (0.017) and MCH (0.017). Whereas Pearson's correlation analysis results in the following Table 3 described that several hematological parameters had a very significant correlation to SNP (amino acid). These parameters were RBC (0.001), HGB (0.006) and HCT (0.001)

Other parameters had a significant correlation to SNP were WBC (0.033) and MCHC (0.032). Only albumin (0.013) as a blood chemistry parameter had a significant correlation to SNP (amino acid). These hematological and blood chemistry

parameters that had a very significant and significant correlation to SNP (nucleotide and amino acid) would be used as a dependent variable to construct multidimensional scaling (MDS) map. The results of multidimensional scaling map would be illustrated in the following figures.

Fig 6 above shows that all samples were divided into three groups. Angus-PO CB (SpB) was localized separately (A). The second group members were purebred (B). The last group members were crossbred (C), except S1 (Ongole) was purebred. Results of the MDS map (Fig 6) shows that each group has mostly the same members configuration as the phylogenetic tree groups.

Fig 7 above described that the MDS map configuration based on the Amino Acid correlation to hematological and blood chemistry parameters was mostly the same as the MDS map based on the Nucleotide correlation to both variables (Fig 7), so both MDS maps had similar




Fig 6: MDS map based on the hematological parameters correlation to SNP nucleotides.




Fig 7: MDS map based on the hematological and blood chemistry parameter correlation to SNP amino acids.

configuration members as the phylogenetic tree for describing the relationship among samples. MDS map method could be an appropriate technique to support and complete Neighbor-joining for constructing the phylogenetic tree.

CONCLUSION

This study revealed that a lot of SNP found in local cattle (Bali and PO) and CB cattle in Gowa and Sidrap indicated various genetic diversity could be a marker candidate and potential to be developed in future research knowing the correlation of the SNP with the growth rate of CB. This study also shows that the correlation of SNP to hematological and blood chemistry description can be an appropriate potential indicator and parameter to describe healthy animal status. This information will be helpful for further strategy in breeding development program.

Conflict of interest: None.

REFERENCES

- Agung, P.P., Saputra, F., Zein, M., Wulandari, A.S., Putra, W., Said, S. and Jakaria, J. (2019). Genetic diversity of Indonesian cattle breeds based on microsatellite markers. *Asian-Australasian Journal of Animal Sciences*. 32(4): 467-476.
- Ansori, A.N.M., Kusala, M.K.J., Nidom, R.V., Indrasari, S., Zarkasie, K., Santoso, K.P. and Nidom, C.A. (2021). Pathological and molecular characterization of Newcastle disease virus isolated from *Gallus gallus* in Java, Indonesia. *Indian Journal of Animal Research*. 55(8): 930-935.
- Bakae, T., Monau, P.I., Nsoso, S.J. and Kgwatalala, P.M. (2022). Assessment of genetic diversity and relationship of the two Sanga type cattle of Botswana based on microsatellite markers. *Tropical Animal Health and Production*. 54(4): 210. DOI: 10.1007/s11250-022-03212-9.
- Cai, L. and Ma, H. (2016). Using nuclear genes to reconstruct angiosperm phylogeny at the species level: A case study with Brassicaceae species. *Journal of Systematics and Evolution*. 54(4): 438-452.
- Dakhlan, A., Adhianto, K., Sulastri, Kurniawati, D., Ermawati, R. and Doni, S.T. (2022). Mapping growth hormone gene of body weight Krui cattle in Pesisir Barat Regency Lampung, Indonesia. *Pakistan Journal of Biological Sciences*. 25(8): 741-747.
- Hidayati, Misrianti, R. and Ali, A. (2016). Phylogenetic tree of Kuantan cattle by DNA barcoding. *Indonesian Journal of Animal and Veterinary Sciences*. 21(1): 41-48.
- Jobirov, F., Yuejie, Z. and Kibona, C.A. (2022). Evaluating profitability of beef cattle farming and its determinants among smallholder beef cattle farmers in the Baljovan District of Khatlon region, Tajikistan. *PloS One*. 17(9): e0274391.
- Kibona, C.A., Yuejie, Z. and Tian, L. (2022). Factors that influence beef meat production in Tanzania. A Cobb-Douglas production function estimation approach. *PloS One*. 17(8): e0272812.
- Kumar, S., Ganguly, I., Singh, S., Sodhi, M. and Bhakat, M. (2017a). Y-chromosome variation in Indian native cattle breeds and crossbred population. *Indian Journal of Animal Research*. 51(1): 1-7.
- Kumar, V., Sharma, N. and Sharma, A. (2017b). DNA barcoding of the Indian blackbuck (*Antelope cervicapra*) and their correlation with other closely related species. *Egyptian Journal of Forensic Sciences*. 7: 31. <https://doi.org/10.1186/s41935-017-0034-6>.
- Kuswati, K., Furqon, A., Septian, W.A. and Susilawati, T. (2022). Polymorphism of leptin gene (single nucleotide polymorphisms c.73T>C) and its association with body weight and body measurements in Madura cattle. *Veterinary World*. 15(3): 775-781.
- Liu, Y., Cheng, H., Wang, S., Luo, X., Ma, X., Sun, L., Chen, N., Zhang, J., Qu, K., Wang, M., Liu, J., Huang, B. and Lei, C. (2022). Genomic diversity and selection signatures for weining cattle on the border of Yunnan-Guizhou. *Frontiers in Genetics*. 13: 848951.
- Malau-Aduli, A., Curran, J., Gall, H., Henriksen, E., O'Connor, A., Paine, L., Richardson, B., van Slidregt, H. and Smith, L. (2021) Genetics and nutrition impacts on herd productivity in the Northern Australian beef cattle production cycle. *Veterinary and Animal Science*. 15: 100228.
- Mayulu, H., Sunarso, Sutrisno, C.I. and Sumarsono. (2010). Beef cattle farm development policy in Indonesia. *Journal of Agricultural Research and Development*. 29(1): 34-41.
- Mohamad, K., Olsson, M. andersson, G., Purwantara, B., van Tol, H.T., Rodriguez-Martinez, H., Colenbrander, B. and Lenstra, J.A. (2012) The origin of Indonesian cattle and conservation genetics of the Bali cattle breed. *Reproduction in Domestic Animals*. 47(Suppl 1): 18-20.
- Naseer, J., Anjum, K.M., Khan, W.A., Imran, M., Ishaque, M., Hafeez, S., Munir, M.A., Nazir, M.A. (2018). Phylogenetic analysis-based studies on genetic variation of Cytochrome b gene of Indian peafowl (*Pavo cristatus*) in Pakistan. *Indian Journal of Animal Research*. 52(3): 343-346.
- Pakpahan, S., Widayanti, R. and Artama, W.T. (2022). Selection signatures and formation of the Samosir goat breed through the cultures of the Batak Toba Tribe in Samosir Island, Indonesia. *Veterinary World*. 15(4): 1044-1050.
- Poniente, J.A., Pereda, J.M.R., Peña, J.T.D., Ventolero, M.F.H., Santos, M.D. (2022). Mitochondrial DNA-based species testing of confiscated aquatic wildlife in the Philippines. *Forensic Science International: Animals and Environments*. 2: 100051.
- Prihandini, P.W., Primasari, A., Aryogi, A., Luthfi, M. and Hariyono, D. (2022). Genetic polymorphisms of the 5' untranslated regions of the HSP70 gene in Indonesian cattle populations. *Veterinary World*. 15(1): 168-172.
- Pundir, R.K., Singh, P.K., Dangi, P.S., Kumar, A., Borah, S., Mahanta, N. and Mette, S.L. (2019). Phenotypic characterisation of indigenous cattle of Meghalaya State. *Indian Journal of Animal Research*. 53(1): 59-62.
- Sari, E.M., Jianlin, H., Noor, R.R., Sumantri, C. and Margawati, E.T. (2016). Phylogenetic analysis of Aceh cattle breed of Indonesia through mitochondrial D-Loop region. *Journal of Genetic Engineering and Biotechnology*. 14(1): 227-231.
- Sudrajat, P., Kusminanto, R.Y., Volkandari, S.D. and Cahyadi, M. (2022). Genomic structure of Bali cattle based on linkage disequilibrium and effective population size analyses using 50K single nucleotide polymorphisms data. *Veterinary World*. 15(2): 449-454.

- Sutarno and Setyawan, A.D. (2016). The diversity of local cattle in Indonesia and the efforts to develop superior indigenous cattle breeds. *Biodiversitas*. 16: 275-295.
- Tamura, K., Stecher, G. and Kumar, S. (2021) MEGA11: Molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution*. 38(7): 3022-3027.
- Trujano-Chavez, M.Z., Ruíz-Flores, A., López-Ordaz, R. and Pérez-Rodríguez, P. (2022). Genetic diversity in reproductive traits of Braunvieh cattle determined with SNP markers. *Veterinary Medicine and Science*. 8(4): 1709-1720.
- Xu, L.X., Liu, J., He, G.Z., Zhang, L. and Zhou, W.Z. (2020). Detection of genetic polymorphism of ANGPTL4 and POMC gene and correlation with growth traits in Guanling cattle. *Indian Journal of Animal Research*. 54(11): 1343-1346.