



The Potential Influence of *Coleus forskolii*, Green Coffee and Green Tea on Body Weight, Reproductive Performance, Hematological Profiles in Mice

A.A. Mohammed¹, M.S. El. Sayed², I. AlGherair¹, S. Al-Suwaiegh¹

10.18805/IJAR.BF-1916

ABSTRACT

Background: *Coleus forskolii*, green coffee and green tea have been shown to effect on body functions and health.

Methods: Twenty males and thirty female albino mice of 33.86 ±0.27g BW were classified into two groups; control group fed basal control diet versus treated group fed basal control diet containing multi-ingredient supplement (*Coleus forskolii*, green coffee bean and green tea; 1.5% per supplement)for ten weeks. Initial and final body weights were recorded of control and supplemented groups. Germinal vesicle oocytes were harvested from ovaries after 48:0h of PMSG injection for investigating oocyte quality. Furthermore, female mice were mated with fertile males and pregnancy rate and offspring numbers were recorded. In addition, the male mice of both groups were sedated using 26.6 mg/kg BW xylazine for recording rectal temperature, pulse rate and SPO2. Besides, blood samples were collected for complete blood picture and plasma biochemistry analyses.

Result: The results showed that multi-ingredient supplement caused significant decrease in body weight gain versus increase (P<0.05) of rectal temperature. Reproductive performances represented by oocyte quality and pregnancy rate were significantly decreased in multi-ingredient supplement group. Furthermore, blood cells and plasma profiles (urea, creatinine, glucose, aspartate aminotransferase, alanine aminotransferase and lactate dehydrogenase) were decreased in multi-ingredient supplement groups except total protein if compared to control group. It could be concluded that multi-ingredient supplement resulted in significant decrease of body weight gain, reproductive performances and blood cell profiles as well.

Key words: *Coleus forskolin*, Glucose, Green coffee bean, Green tea, Oocyte, Reproductive.

INTRODUCTION

Dietary supplements are given to stimulate health and alleviate body dysfunctions in mammalian species (Mohammed 2012, 2018, 2019; Mohammed and Al-Khamis 2024; Mohammed *et al.*, 2018, 2020, 2021, 2024a,b,c,d; Al Masruri *et al.*, 2022a,b; Al-Saiady *et al.*, 2024). *Coleus forskolii*, green coffee bean and green tea were found to increase body weight loss, diabetes, thermogenesis and cardiovascular health (Litoschet *et al.*, 1982; Godard *et al.*, 2005; Thom, 2007; Revuelta-Iniesta *et al.*, 2014; Huang *et al.*, 2014; Roshni and Rekha, 2024). Therefore, the potential effects of those ingredients were explored on changes of body weight gain, reproductive performances concerning oocyte quality and pregnancy rate, blood and plasma profiles (Mohammed *et al.*, 2024d,e).

Coleus forskolii is a medicinal plant cultivated in India and other countries of Asia and eastern Africa (Tunget *et al.*, 2021). The demand for *C. forskolii* plant is increasing because it is importance in dietary and beverage supplements, pharmaceuticals and cosmetics usages (Roshni and Rekha 2024). *C. forskolii* plant was used for forskolin compound production (Mitra *et al.*, 2020; Kundur and Shyam 2024). Dietary *C. forskolin* plant and forskolin compound were found to increase lipolysis, improve diabetes and cardiovascular health and increase blood flow to the brain (Amezcuca *et al.*, 2022; Abbasi *et al.*, 2023; Roshni and Rekha 2024).

¹Department of Animal and Fish Production, College of Agriculture and Food Sciences, King Faisal University, P.O. Box 402, Al-Ahsa 31982, KSA.

²Avian Research Center, King Faisal University, P.O. Box 402, Al-Ahsa 31982, KSA.

Corresponding Author: A.A. Mohammed, Department of Animal and Fish Production, College of Agriculture and Food Sciences, King Faisal University, P.O. Box 402, Al-Ahsa 31982, KSA.
Email: aamohammed@kfu.edu.sa

ORCID: 0000-0002-1825-7248, 0000-0002-0903-1036, 0009-0000-9880-3255, 0000-0003-4392-5508

How to cite this article: Mohammed, A.A., Sayed, M.S.E., AlGherair, I. and Al-Suwaiegh, S. (2024). The Potential Influence of *Coleus forskolii*, Green Coffee and Green Tea on Body Weight, Reproductive Performance, Hematological Profiles in Mice. Indian Journal of Animal Research. 1-8. doi: 10.18805/IJAR.BF-1916.

Submitted: 01-11-2024 **Accepted:** 19-12-2024 **Online:** 13-01-2025

The effects of coffee extract on body functions were investigated in several studies including blood glucose and insulin values post exercise (Beam *et al.*, 2015). No significant effects of green coffee bean extract for blood glucose and insulin values were obtained. In addition, coffee extract effects on lipid metabolism were explored in several studies (Choi *et al.*, 2016; Takahashi *et al.*, 2017; Farias-Pereira *et al.*, 2018). The doses were given in diet (1-2%) or water (0.1-0.5%) or according to body weight

(200-1000 mg/kg BW). The results showed decreases in body weight, glucose, triglycerides, cholesterol, HDL, LDL, insulin and leptin values versus increases of ATP levels.

In addition, green tea was shown to reduce appetite and increase thermogenesis. Besides, blood pressure and lipid profiles were changed due to green tea supplementation (Onakpoya *et al.*, 2014). Dietary green tea was significantly reduced systolic blood pressure, total cholesterol and LDL-cholesterol values. Collectively, the ingredients of *Coleus forskolii* (Mohammed *et al.*, 2024), green coffee bean (Thom 2007; Revuelta-Iniesta *et al.*, 2014) and green tea (Huang *et al.*, 2014) were known to increase lipolysis and thermogenesis via a variety of biological pathways (Litosch *et al.*, 1982; Godard *et al.*, 2005; Thom 2007; Revuelta-Iniesta *et al.*, 2014; Huang *et al.*, 2014). They have been associated with changing body composition (Barrea *et al.*, 2019) and may simultaneously influence the main pathways underlying obesity. Therefore, the aims of the current study were to explore the changes in body weight gain, rectal temperature, pulse rate and SPO₂ of control and multi-ingredient supplement groups. Besides, reproductive performance concerning oocyte quality and pregnancy rate were recorded as well. In addition, the changes in hematological profiles and plasma biochemistry values were investigated as well.

MATERIALS AND METHODS

The experimental procedures were approved by the ethical committee of King Faisal University [KFU-REC-2024-OCT-

EA241670]. The *C. forskolin* dried leaves were obtained from farm located in Gizan area of KSA. The leaves were arranged in a single layer on a baking sheet of oven and they were exposed to 50-60°C for sixteen hours. In addition, green coffee bean (*Coffe arabica*) and green tea were purchased from the market. The experimental procedures were carried out in the experimental animal lab of Agriculture and Food Sciences College, KFU University.

Site of study and animal management

The current study was executed from August to October 2024 in animal lab of animal and fish production department. Twenty males and thirty females albino mice were used for the study (body weight; 33.86±0.27 g) (Fig 1). Mice were kept in groups of five mice per cage (40×24× 18 cm) of control and the treated groups. The animals were given pellet diet (Arasco, KSA), which composed of 2.90% fat, 22.0% protein, 3.20% fiber, 1.0% mixture of vitamins and minerals and energy of 3300.00 kcal/kg. The supplemented doses of *C. forskolin*, green coffee bean and green tea were 1.5% of each supplement according to the recommended doses of previous studies (Shimoda *et al.*, 2006; Murase *et al.*, 2010; Farias-Pereira *et al.*, 2018; Mohammed *et al.*, 2024 d,e). The ground basal diet was mixed with 1.5% powders of each supplemented ingredient and pelleted. Animals had free access to water and diets. Mice were kept controlled under 12 h light cycle and 12 h dark starting at 8 a.m. The relative humidity (%) and temperature (°C) values during the study were controlled




Fig 1: Experimental design of multi-ingredient supplement impact on thermo-tolerance parameters, blood and plasma profiles in DX anesthetized mice.

to $25 \pm 2.50^\circ\text{C}$ and $45 \pm 9.0\%$, respectively. The duration of study was twelve weeks.

Monitoring oocyte quality and reproductive performance

Five females of each group were injected with 7.5 IU of PMSG to synchronize ovarian follicle development. The injected females were sacrificed through cervical dislocation 48h after PMSG injection. Ovaries were removed and germinal vesicle oocytes were released by puncturing of ovarian follicles with 30-G sterile needles under a stereomicroscope. The oocytes were released into TCM199 medium supplemented with BSA. The diameters of denuded GV oocytes were determined to the nearest micrometer using 0.01 mm eyepiece on a compound biological microscope (Mohammed, 2009; Mohammed and Al-Hozab, 2019). For reproductive performance, 10 females of each group were bred with male of proven fertility for eight weeks. The pregnant females and delivered pups were recorded.

Monitoring body weight, rectal temperature, heart rate, partial pressure of oxygen and values

Body weights (g) were recorded at the beginning and end of study (Sartorius balance, Azulmart-KSA). The male mice of both groups were sedated using 26.6 mg/kg BW xylazine for immediate recording of rectal temperature, pulse rate and SPO₂. Rectal temperatures were monitored using clinical thermometer (Citizen) (Mohammed, 2018; Mohammed *et al.*, 2024e,f). Pulse rate and partial pressure of oxygen were recorded using pulse oximeter device (CMS60D-VET).

Blood sample collection and analyses

Blood samples were collected from the orbital sinus of control and multi-ingredient supplement groups at the end of experiment (Hoff, 2000). The obtained blood samples were analyzed using automatic hematology analyzer (Mythic 5Vet PRO Hematology Analyzer) and biochemistry analyzer (Skyla VB1). The resulting blood parameters include red and white blood cell count, hematocrit and hemoglobin. The resulting biochemistry parameters include total proteins, liver enzymes, urea, creatinine and mineral values.

Statistical analysis

Body weight and rectal temperature, SPO₂ and pulse rate, oocyte quality and reproductive performance and blood and plasma profile values of control and multi-ingredient supplement groups were statistically analyzed using General Linear Model procedure of SAS (SAS 2008) according to the following model:

$$Y_{ij} = \mu + T_i + e_{ij}$$

Where:

μ = Mean.

T_i = Effects of multi-ingredient supplement.

e_{ij} = Standard error.

Duncan's multiple range test (1955) was used to compare between means of control and multi-ingredient supplement groups.

RESULTS AND DISCUSSION

Body weight and temperature, SPO₂ and pulse rate, oocyte quality and reproductive performance and blood and plasma biochemistry profiles of multi-ingredient supplement and control groups are presented in Table (1-4).

Body weight, rectal temperature, SPO₂ and pulse rate

Changes of body weight, rectal temperature, SPO₂ and pulse rate are presented in Table (1). Value of final body weight (g) was significantly decreased in multi-ingredient supplement group compared to that of control group. On the other hand, hyperthermia of rectal temperature ($^\circ\text{C}$) was recorded in multi-ingredient supplement group. Additionally, pulse rate and SPO₂ were not differed between both groups.

Monitoring oocyte quality and reproductive performance

Values of oocyte quality and reproductive performance are presented in Table (2). Percentages of GV oocytes with diameter 70.0 and 80.0 μm were lower in multi-ingredient supplement group compared to those of control group versus the percentage of GV oocytes with diameter 60.0 μm if compared to control group. Simultaneously, pregnancy rate and number of pups per female were decreased ($P < 0.001$) in multi-ingredient supplement group compared to control one.

Blood and plasma profiles

Values and blood and plasma profiles are presented in Table (3-4). Values of blood parameters were significantly improved in control group compared to multi-ingredient supplement group (Table 3). Regarding to plasma profiles, total protein values were increased whereas urea, creatinine, glucose, aspartate aminotransferase, alanine aminotransferase and lactate dehydrogenase values were significantly decreased in multi-ingredient supplement group versus control group.

Results of the current study are presented in Table (1-4) indicating the effects of dietary (1.5%) *C. forskohlii*, green coffee bean and green tea for each supplement on body weight, reproductive performance, hematological profiles in mice. Generally, the obtained results indicated significant decrease of body weight gain, blood and plasma profiles in addition to increase of rectal temperature and plasma total protein in the multi-ingredient supplement group compared to control group. Furthermore, the reproductive performances represented by oocyte quality and pregnancy rate and delivered pups were significantly decreased in the multi-ingredient supplement group compared to control group. These effects could be owing to the common ingredients in *C. forskohlii*, green coffee bean and green tea including botanicals (herbs and other plant components), caffeine, dietary fibers and minerals. Both *C. forskohlii*, green coffee bean and green tea contain various compounds, some of which can be potentially harmful if consumed in excess amount.

Dietary *C. forskolii* and its extracted compound, forskolin, have been explored for a variety of potential health benefits (Amezcuca *et al.*, 2022; Roshni and Rekha 2024). These benefits may include body weight loss, improved diabetes and cardiovascular health, increased muscle mass and enhanced blood flow to the brain (Abbasi *et al.*, 2023; Mohammed *et al.*, 2024d,e). The primary active ingredient in green coffee is chlorogenic acid. This compound is believed to be responsible for many of the

potential health benefits as anti-obesity, regulating glucose and lipid metabolism, antioxidant and anti-inflammatory impacts (Tajik *et al.*, 2017; Yu *et al.*, 2022; Nguyen *et al.*, 2024). The primary active ingredient in green tea is a group of compounds called polyphenols, specifically catechins (Farhan 2022). Among these catechins, epigallocatechin gallate is considered the most potent and abundant compound. These compounds are responsible for many of the health benefits attributed to green tea as antioxidant,

Table 1: Changes of body weight (g), rectal temperature (°C), pulse rate and SPO₂ (%) and glucose values (mg/dl) of multi-ingredient supplement and control groups in mice.

Parameters	Treat	Control	SE	P value
Initial BW, g	33.88	33.84	1.18	0.97
Final BW, G	38.75 ^b	43.10 ^a	1.55	<0.001
Rectal temperature, °C	36.68 ^a	36.46 ^b	0.11	<0.001
Pulse rate/min.	165.7	165.3	6.24	0.97
SPO ₂ , %	82.3	81.6	3.17	0.96

a, b: Values between groups with different superscripts significantly differ at P<0.05.

SE: Standard error.

Table 2: Changes of Oocyte quality and reproductive performance of multi-ingredient supplement and control groups in mice.

Parameters	Treat	Control	SE	P value
Total collected oocytes/5 females	70.0 ^b	80.0 ^a	0.72	<0.01
No. collected oocytes/female	14.0 ^b	16.0 ^a	0.72	<0.02
Diameter, 60 µm	61.4 ^a	46.25 ^b	3.10	<0.001
Diameter, 70 µm	25.7 ^b	35.00 ^a	2.66	<0.001
Diameter, 80 µm	12.90 ^b	18.75 ^a	2.55	<0.001
Pregnancy rate, %	30.0 ^a (3/10)	80.0 ^a (8/10)	1.40	<0.001
Pups number/female	8.0 ^a	9.125 ^a	0.74	<0.01

a, b: Values between groups with different superscripts significantly differ at P<0.05.

SE: Standard error.

Table 3: Changes of blood profiles of multi-ingredient supplement and control groups in mice.

Parameters	Treat	Control	SE	P value
RBCs, 10 ⁶ /µl	4.50 ^b	6.40 ^a	0.18	<0.0001
PCV, %	22.7 ^b	29.7 ^a	1.21	<0.0001
Hb, g/dl	8.00 ^b	10.0 ^a	0.16	<0.0001
MCV, fl	50.6 ^a	46.4 ^b	3.41	<0.07
MCH, pg	18.0 ^a	15.8 ^b	1.12	<0.05
MCHC, g/dl	36.0	34.0	2.36	<0.07
RDW, %	14.9 ^b	17.0 ^a	0.98	<0.04
WBCs, 10 ³ /ml	5.25 ^b	8.73 ^a	1.32	<0.0001
Neutrophils, 10 ³ /µl	2.451 ^b	2.604 ^a	0.09	<0.01
Lymphocyte, 10 ³ /µl	2.002 ^b	4.505 ^a	0.14	<0.0001
Monocyte, 10 ³ /µl	0.437 ^b	1.219 ^a	0.001	<0.0001
Eosinophils, 10 ³ /µl	0.309 ^b	0.299 ^a	0.001	<0.06
Basophils, 10 ³ /µl	0.051 ^b	0.103 ^a	0.0002	<0.0001
PLT, 10 ³ /µl	1016 ^b	1293 ^a	15.19	<0.001

a, b: Values between groups with different superscripts significantly differ at P<0.05.

SE: Standard error; RBCs: Red blood cells; PCV: Packed cell volume; Hb: Hemoglobin; MCV: Mean corpuscular volume; MCH: Mean corpuscular hemoglobin; MCHC: Mean corpuscular hemoglobin concentration; RDW: Red cell distribution width; WBCs: White blood cells; PLT: Platelet.

Table 4: Changes of profiles of multi-ingredient supplement and control groups in mice.

Parameters	Treat	Control	SE	P value
Total protein, g/dl	5.1 ^a	4.9 ^b	0.15	0.01
Urea, mg/dl	24.0 ^a	28.0 ^a	1.04	<0.0001
Creatinine, mg/dl	0.50 ^b	0.70 ^a	0.03	<0.0001
Glucose, mg/dl	81.0 ^b	104.0 ^a	2.50	<0.0001
Aspartate Aminotransferase, U/l	180.0	205.0	13.3	<0.06
Alanine Aminotransferase, U/l	216.0	261.0	13.5	<0.04
Lactate dehydrogenase, U/l	513	551	21.9	0.07
Iron, µmol/L	296	312	10.83	0.09

a, b: Values between groups with different superscripts significantly differ at P<0.05.

SE: Standard error.

weight management, improved diabetes and heart health properties (Du *et al.*, 2012; Legeay *et al.*, 2015; Mohammed and Al-Hozab 2019).

Collectively, the percentage 1.5% of each supplement resulted in lower body weight gain, hyperthermia and hypoglycemia if compared to basal control diet. Furthermore, values of blood cells and plasma profiles (urea, creatinine, glucose, aspartate aminotransferase, alanine aminotransferase and lactate dehydrogenase) were decreased in multi-ingredient supplement groups except total protein if compared to control group. It seems that such supplements gave adverse effects in some parameters due to excess supplemented amount or harmful ingredients as caffeine, tannins and oxalates (Noonan and Savage 1999; Duda-Chodak and Tarko 2023). Tannins are polyphenols found in both green tea and green coffee offer some health benefits, while excessive tannin intake can interfere with iron absorption and may worsen digestive issues (Delimont *et al.*, 2017; Duda-Chodak and Tarko 2023). Excess caffeine can affect various aspects of female reproductive performances including slight decrease estrogen levels, possible delay in conception, increased risk of miscarriage, stillbirth and lower birth weight (Wen *et al.*, 2001; Karypidis *et al.*, 2006; Weng *et al.*, 2008; Greenwood *et al.*, 2010; Wikoff *et al.*, 2017; Bu *et al.*, 2020), where we found lower quality of GV oocytes and pregnancy rate in this study. In addition, the lower reproductive performance in supplement group could be also attributed to decrease of steroid hormone values due to the changes in lipid metabolism, low glucose level and body weight gain (Senosy *et al.*, 2018; Mohammed *et al.*, 2022).

CONCLUSION

It could be concluded that *C. forskolii*, green coffee bean and green tea multi-ingredient supplement modulates body weight gain, thermo-tolerance response, blood and plasma profiles. On the other hand, multi-ingredient supplement resulted in negative effects reproductive efficiency represented by oocyte quality, pregnancy and delivered pups. Further studies are still required for exploring multi-ingredient supplement effects through the level of supplementation and the resulting physiological body functions and reproductive performance.

ACKNOWLEDGEMENT

The authors want to thank and acknowledge Deanship of Scientific Research, King Faisal University, Saudi Arabia for funding and support (KFU242726).

Disclaimers

The views and conclusions expressed in this article are solely those of the authors and do not necessarily represent the views of their affiliated institutions. The authors are responsible for the accuracy and completeness of the information provided, but do not accept any liability for any direct or indirect losses resulting from the use of this content.

Informed consent

The Ethical Committee of Deanship of Scientific Research, King Faisal University, Saudi Arabia, approved all animal procedures, experimental animal care and handling techniques for experiments (KFU242726).

Conflict of interest

The authors declare that there are no conflicts of interest regarding the publication of this article. No funding or sponsorship influenced the design of the study, data collection, analysis, decision to publish, or preparation of the manuscript.

REFERENCES

- Abbasi, M., Zhou, F., Hu, Q. and Wang, S. (2023). P31-001-23 Anti-obesity and metabolic effects of forskolin in obese C57BL/6J mice. *Current Developments in Nutrition*. 7(1): 101-567.
- Al Masruri, H., Al Zeidi, R., Al Mufarji, A. and Mohammed A.A. (2022a). *Moringa oleifera* leaves supplementation to anaesthetized mice associated with changes of thermo-tolerance parameters, blood and plasma indices. *Fresenius Environmental Bulletin*. 8: 8179-8187.
- Al Masruri, H., Al Zeidi, R., Al Mufarji, A., Mohammed, A.A., Al Madani, A. and Mohammed, H. (2022b). Leverage of *Moringa oleifera* supplementation on performances, biochemical and milk profiles in mammals. *Advances Animal Veterinary Sciences*. 10(9): 2043-2050.

- Al-Saiady, M., Al-Shaheen, T., El-Waziry, A., Mohammed, A.E.A. (2024). Effects of extruded flaxseed and Salmate® inclusion in the diet on milk yield and composition, ruminal fermentation and degradation and kinetic flow of digesta and fluid in lactating dairy cows in the subtropics. *Veterinary World*. 17(3): 540-549.
- Amezcuca, I.J., Blas, S.R., Municio, M.D., Soria, A.C., Matute, A.I.R. and Sanz, M.L. (2022). Development of a multianalytical strategy for detection of frauds in *Coleus forskolii* supplements. *Journal of Chromatography A*. 1676: 463-198.
- Barrea, L., Altieri B., Polese B., De Conno B., Muscogiuri G., Colao A., Savastano S. (2019). Nutritionist and obesity: Brief overview on efficacy, safety and drug interactions of the main weight-loss dietary supplements. *International Journal of Obesity Supplements*. 9: 32-49.
- Beam, J.R., Gibson, A.L., Kerksick, C.M., Conn, C.A., White, A.C., Mermier, C.M. (2015). Effect of post-exercise caffeine and green coffee bean extract consumption on blood glucose and insulin concentrations. *Nutrition*. 31(2): 292-297.
- Bu, F.L., Feng, X., Yang, X.Y., Ren, J., Cao, H.J. (2020). Relationship between caffeine intake and infertility: A systematic review of controlled clinical studies. *BMC Womens Health*. 20(1): 125.
- Choi, B.K., Park, S.B., Lee, D.R., Lee, H.J., Jin, Y.Y., Yang, S.H., Suh, J.W. (2016). Green coffee bean extract improves obesity by decreasing body fat in high-fat diet-induced obese mice. *Asian Pacific Journal of Tropical Medicine*. 9: 635-643.
- Delimont, N.M., Haub, M.D., Lindshield, B.L. (2017). The Impact of tannin consumption on iron bioavailability and status: A Narrative Review. *Current Developments in Nutrition*. 1(2): 1-12.
- Du, G.J., Zhang, Z., Wen, X.D., Yu, C., Calway, T., Yuan, C.S., Wang, C.Z. (2012). Epigallocatechin Gallate (EGCG) is the most effective cancer chemopreventive polyphenol in green tea. *Nutrients*. 4(11): 1679-1691.
- Duda-Chodak, A., Tarko, T. (2023). Possible side effects of polyphenols and their interactions with medicines. *Molecules*. 28(6): 2536.
- Duncan, D.B. (1955). Multiple range and multiple Ftest. *Biometrics*. 11: 1.
- Farhan, M. (2022). Green Tea Catechins: Nature's Way of Preventing and Treating Cancer. *International Journal of Molecular Sciences*. 23(18): 10713.
- Farias-Pereira, R., Oshiro, J., Kim, K.H., Park, Y. (2018). Green coffee bean extract and 5-O-caffeoylquinic acid regulate fat metabolism in *Caenorhabditis elegans*. *Journal Functional Foods*. 48: 586-593.
- Godard, M.P., Johnson, B.A., Richmond, S.R. (2005). Body Composition and hormonal adaptations associated with forskolin consumption in overweight and obese men. *Obesity Research*. 13: 1335-1343.
- Greenwood, D.C., N. Alwan, S. Boylan, J.E. Cade, J. Charvill, K.C. Chipps, M.S. Cooke, V.A. Dolby, A.W.M. Hay, S. Kassam, et al. (2010). Caffeine intake during pregnancy, late miscarriage and stillbirth. *European Journal of Epidemiology*. 10: 1007.
- Henderson, S., Magu, B., Rasmussen, C., Lancaster, S., Kerksick, C., Smith, P., Melton, C., Cowan, P., Greenwood, M., Earnest, C. et al. (2005). Effects of *coleus forskolii* supplementation on body composition and hematological profiles in mildly overweight women. *Journal of the International Society of Sports Nutrition*. 2(2): 54-62.
- Hoff, J. (2000). Methods of blood collection in the mouse. *Laboratory Animals*. 29: 47-53.
- Huang, J., Wang, Y., Xie, Z., Zhou, Y., Zhang, Y., Wan, X. (2014). The anti-obesity effects of green tea in human intervention and basic molecular studies. *European Journal of Clinical Nutrition*. 68: 1075-1087.
- Karypidis, A.H., Soderstrom, T., Nordmark, A., Granath, F., Cnattingius, S., Rane, A. (2006). Association of cytochrome P450 1B1 polymorphism with first-trimester miscarriage. *Fertility Sterility*. 86: 1498-1503.
- Kundur, S., Shyam, P. (2024). Quantification of total phenols in *Coleus forskolii* and inhibition of aldose reductase by rosmarinic acid. *Journal of Molecular Structure*. 1312: 138-435.
- Legeay, S., Rodier, M., Fillon, L., Faure, S., Clere, N. (2015). Epigallocatechin Gallate: A Review of Its Beneficial Properties to Prevent Metabolic Syndrome. *Nutrients*. 7: 5443-5468.
- Litosch, I., Hudson, T.H., Mills, I., Li, S.Y., Fain, J.N. (1982). Forskolin as an activator of cyclic AMP accumulation and lipolysis in rat adipocytes. *Molecular Pharmacology*. 22: 109-115.
- Mitra, M., Gantait, S., Mandal, N. (2020). *Coleus forskolii*: advancements and prospects of *in vitro* biotechnology. *Applied Microbiology Biotechnology*. 104(6): 2359-2371.
- Mohammed, A.A. (2009). Ovarian morphology and oocyte quality in relation to *in vitro* embryo production in ruminant animals. *Assiut Journal of Agricultural Sciences*. 40(2): 1-12.
- Mohammed, A.A. (2019). Nigella Sativa oil improves physiological parameters, oocyte quality after ovarian transplantation and reproductive performance of female mice. *Pakistan Journal Zoology*. 51(6): 2225-2231.
- Mohammed, A.A., Al-Hozab A.A. (2019). +(-)catechin raises body temperature, changes blood parameters, improves oocyte quality and reproductive performance of female mice. *Indian Journal of Animal Research*. 54(5): 543-548. doi: 10.18805/IJAR.B-981.
- Mohammed, A.A., Al-Hozab, A., Alshaheen, T. (2018). Effects of diazepam and xylazine on changes of blood oxygen and glucose levels in mice. *Advances in Animal and Veterinary Sciences* 6(3): 121-127.
- Mohammed, A.A. (2012). Ovarian tissue transplantation in mice and rats: Comparison of Ovaries Age. *Pakistan Journal of Zoology*. 50(2): 481-486.
- Mohammed, A.A. (2018). Development of oocytes and preimplantation embryos of mice fed diet supplemented with *dunaliella salina*. *Advances Animal Veterinary Sciences*. 6(1): 33-39.
- Mohammed, A.A., Al-Khamis, S. (2024). Effects of dietary microalgae inclusion on feed utilization, morphological and chemical characteristics and body health in fish: Advantages and Disadvantages. *Indian Journal of Animal Research*. 58(9): 1452-1459. doi: 10.18805/IJAR.BF-1748.

- Mohammed, A.A., Abdelnabi1, M.A., Modlinski, J.A. (2012). Evaluation of anesthesia and reproductive performance upon diazepam and xylazine injection in rats. *Animal Science Papers and Reports*. 30(3): 285-292.
- Mohammed, A.A., Alshaheen, T. and Suwaiegh, S. (2020). Effects of *Myo-inositol* on physiological and reproductive traits through blood parameters, oocyte quality and embryo transfer in mice. *Indian Journal of Animal Research*. 10.18805/ijar.B-1300.
- Mohammed, A.A., AlHizab F., AlSuwaiegh S., Alshaheen T., Kassab A., Hamdon H., Senosy W. (2021). Effects of propylene glycol on ovarian restoration, reproductive performance, metabolic status and milk production during transition period of ewes in subtropics. *Fresenius Environmental Bulletin*. 2021: 8192.
- Mohammed, A.A., Al-Suwaiegh, S., AlGherair, I., Alessa, F. and Alhujaili, W.F. (2024a). Promising roles of *zingiber officinale* and its derivatives on promoting health and protecting from disorders. *Indian Journal of Animal Research*. doi:10.18805/IJAR.BF-1788.
- Mohammed, A.A., AlGherair, I., AlSuwaiegh, S., Al-Khamis, S., Alessa, F., Al-Madni, A. and Al-Ghamdi, A. (2024b). Effects of nutritive and non-nutritive feed supplements on feed utilization, growth and reproductive performances in mammals. *Indian Journal of Animal Research*. 58(4): 615-621. doi: 10.18805/IJAR.BF-1891.
- Mohammed, A.A., Al-Suwaiegh, S., AlGherair, I., Al-Khamis, S., Alessa, F., Al-Awaid, S., Alhujaili, W.F., Mohammed, A. and Mohammed, A. (2024c). The Potential impacts of antioxidant micronutrients on productive and reproductive performances of mammalian species during stressful conditions. *Indian Journal of Animal Research*. 58(6): 1031-1038. doi: 10.18805/IJAR.BF-1773.
- Mohammed, A.A., Al-Suwaiegh, S., Al-Gherair, I., Al-Khamis, S., Al-Awaid, S., Mohammed, A., Mohammed, A. (2024d). The adjuvant impacts of antioxidant micronutrients on ovarian follicle development, oocyte maturation and embryo development of mammalian species: A review. *Indian Journal of Animal Research*. 58(10): 1651-1657. doi: 10.18805/IJAR.BF-1816.
- Mohammed, A.A., AlSuwaiegh, S., Al-Gherair, I. (2024e). Potential impacts of dried green coleus forskolin leaves on changes of body and liver weight, glucose and rectal temperature, serum profiles in mice. *Indian Journal of Animal Research*. doi: 10.18805/IJAR.BF-1891.
- Mohammed, A.A., Al-Suwaiegh, S. and AlGherair, I. (2024f). Potential impacts of dried green coleus forskolin leaves on changes of body and liver weight, glucose and body temperature, serum profiles in mice. *Indian Journal of Animal Research*. doi: 10.18805/IJAR.BF-1839.
- Murase, T., Misawa, K., Minegishi, Y., Aoki, M., Ominami, H., Suzuki, Y., Shibuya, Y., Hase, T. (2010). Coffee polyphenols suppress diet-induced body fat accumulation by downregulating SREBP-1c and related molecules in C57BL/6 J mice. *Am. American Journal of Physiology-Endocrinology and Metabolism*. 300: E122-E133.
- Nguyen, V., Taine, E.G., Meng, D., Cui, T., Tan, W. (2024). Chlorogenic acid: A systematic review on the biological functions, mechanistic actions and therapeutic potentials. *Nutrients*. 16: 924.
- Noonan, S.C., Savage, G.P. (1999). Oxalate content of foods and its effect on humans. *Asia Pacific Journal of Clinical Nutrition*. 8(1): 64-74.
- Onakpoya, I., Spencer, E., Heneghan, C., Thompson, M. (2014). The effect of green tea on blood pressure and lipid profile: A systematic review and meta-analysis of randomized clinical trials. *Nutrition, Metabolism and Cardiovascular Diseases*. 24(8): 823-836.
- Revuelta-Iniesta, R., Al-Dujaili E.A.S. (2014). Consumption of Green Coffee Reduces Blood Pressure and Body Composition by Influencing 11 β -HSD1 Enzyme Activity in Healthy Individuals: A Pilot Crossover Study Using Green and Black Coffee. *BioMed Research International*. 482-704.
- Roshni, P.T., Rekha, P.D. (2024). Biotechnological interventions for the production of forskolin, an active compound from the medicinal plant, *Coleus forskohlii*. *Physiology Molecular Biology Plants*. 30(2): 213-226.
- SAS (2008). SAS user's guide: Basics. Statistical Analysis System Institute, Inc., Cary, NC, USA.
- Shimoda, H., Seki, E., Aitani, M. (2006). Inhibitory effect of green coffee bean extract on fat accumulation and body weight gain in mice. *BMC Complement. Alternative medicine*. 6: 9.
- Tajik, N., Tajik, M., Mack, I., Enck, P. (2017). The potential effects of chlorogenic acid, the main phenolic components in coffee, on health: A comprehensive review of the literature. *European Journal of Nutrition*. 56(7): 2215-2244.
- Takahashi, K., Yanai, S., Shimokado, K., Ishigami, A. (2017). Coffee consumption in aged mice increases energy production and decreases hepatic mTOR levels. *Nutrition*. 38: 1-8.
- Thom, E. (2007). The effect of chlorogenic acid enriched coffee on glucose absorption in healthy volunteers and its effect on body mass when used Long-term in overweight and obese People *Journal of International Medical Research*. 35: 900-908.
- Tung, Y.C., Shih, Y.A., Nagabhushanam, K., Ho, C.T., Cheng, A.C., Pan, M.H. (2021). *Coleus forskohlii* and *Garcinia indica* extracts attenuated lipid accumulation by regulating energy metabolism and modulating gut microbiota in obese mice, *Food Research International*. 142: 110-143.
- Wen, W., Shu, X.O., Jacobs, D.R., Brown, J.E. (2001). The associations of maternal caffeine consumption and nausea with spontaneous abortion. *Epidemiology*. 12: 38-42.
- Weng, X., Odouli, R., Li, D.K. (2008). Maternal caffeine consumption during pregnancy and the risk of miscarriage: A prospective cohort study. *American Journal of Obstetrics and Gynecology*. 198: 279.e1-279.e8.
- Wikoff, D., Welsh, B.T., Henderson, R., Brorby, G.P., Britt, J., Myers, E., Goldberger, J., Lieberman, H.R., O'Brien, C., Peck, J. et al. (2017). Systematic review of the potential adverse effects of caffeine consumption in healthy adults, pregnant women, adolescents and children. *Food and Chemical Toxicology*. 109: 585-648.

- Yu, Y., Zhang, Z., Chang, C. (2022). Chlorogenic acid intake guidance: Sources, health benefits and safety. *Asia Pacific Journal of Clinical Nutrition*. 31(4): 602-610.
- Senosy, W., Kassab, A.Y., Hamdon, H.A., Mohammed, A.A. (2018). Influence of organic phosphorus on reproductive performance and metabolic profiles of anoestrous Farafr ewes in subtropics at the end of breeding season. *Reproduction in Domestic Animals*. 53(4): 904-913.
- Senosy, W., Kassab, A.Y., Hamdon, H.A., Mohammed, A.A. (2018). Effects of feeding green microalgae on ovarian activity, reproductive hormones and metabolic parameters of Boer goats in arid subtropics. *Theriogenology*. 96: 16-22.
- Mohammed, A.A., Al Mufarji, A. Alawaid, S. (2022). Developmental potential of ovarian follicles in mammals: involvement in assisted reproductive techniques. *Pakistan Journal Zoology*. 2022: 1-11.