



Relationship between Forage Particle Size and Rate of *in vitro* Digestion Parameters of Alfalfa-based Lamb in Complete Pelleted Diets

Abdulrahman S. Alharthi¹, Faisal A. Alshamiry¹, Ahmed A. Alghonaim¹,
Hani H. Al-Baadani¹, Ibrahim A. Alhidary¹

10.18805/IJAR.BF-1923

ABSTRACT

Background: Intensive feeding systems for lamb production are based on high concentrations of feed, depending on the type of forage, which can cause digestive disturbances; however, forage particle length (FPL) and size (FPS) can affect the fermentation process and the productive performance of lambs. Intensive feeding systems for lamb production are based on high concentrations of feed, depending on the type of forage, which can cause digestive disturbances; however, forage particle length (FPL) and size (FPS) can affect the fermentation process and the productive performance of lambs. Understanding how the lambs meet these challenges requires a conceptual integration of forage structure, mechanical particle size of forage and microbial fermentation. This synthesis will hopefully contribute to the development of a biomimetic co-treatment process for translating the unique characteristics of digestion into an effective component of biorefineries for the industrial conversion of cellulosic in forage biomass to useful products. Therefore, this study aimed to evaluate the relationship among FPL, degradability of feed *in vitro* and digestibility of organic materials.

Methods: As a laboratory simulation of rumen functions, four alfalfa FPS lengths were studied within the components of integrated feed pellets 0.1 mm, common in the Saudi animal feed market, 5, 10 and 15 mm using the same feed composition, but with different lengths.

Result: Cumulative gas production in PSL₈ and PSL₁₂ lambs was higher compared with the other two groups, indicating that lengths >15 mm and <5 mm affected gas production rates. The effect of FPL on rumen parameters had no effect on the pH of the rumen fluid. There were no significant differences in all values of the main volatile fatty acids acetate, propionate and butyrate and digestibility increased when the FPL ranged from 5 to 10 mm as in the PSL₅ and PSL₁₀ groups. Hydrogen levels were significantly higher in the PSL₁ and PSL₁₀ groups than in the PSL₅ and PSL₁₅ groups. Based on forage size, groups PSL₅ to PSL₁₀ showed the possibility of using this type of feed for growing lambs and small ruminants.

Key words: Fatty acids, Forage particle size, *In vitro* digestion, Lamb, Ruminal fermentation.

INTRODUCTION

Geopolitical changes and their impact on food supply chains in addition to global warming, high temperatures and drought (Jenkins, 1987; Izadbakhsh *et al.*, 2024) may pose a threat to animal husbandry in general and growing animals in particular (Letourneau *et al.*, 2024). The mechanisms of coping with heat stress are low compared to those in larger animals (Galanakis, 2023). The problem of greenhouse gas emissions is increasing, especially those produced by ruminant animals (Giamouri *et al.*, 2023), which may account for up to one fifth of total production (FAO, 2022), posing many environmental and health challenges, which may be due to Overfeeding and waste (Alexander *et al.*, 2017), so feeding strategies and controlling feed particle size may contribute to reducing waste and the rate of raid emission. greenhouse effect (Yan *et al.*, 2024), yet both the agricultural and livestock sectors must continue to provide food to cover growing human needs (Jararweh *et al.*, 2023; Fanzo and Miachon 2023); however, certain nutritional strategies and techniques can help overcome the obstacles and harmful effects of heat stress (Li *et al.*, 2021). Increasing benefit levels from provided rations and reducing waste to achieve

¹Department of Animal Production, College of Food and Agriculture Science, King Saud University, P.O. Box 2460, Riyadh 11451, Saudi Arabia.

Corresponding Author: Ibrahim A. Alhidary, Department of Animal Production, College of Food and Agriculture Science, King Saud University, P.O. Box 2460, Riyadh 11451, Saudi Arabia.
Email: ialhidary@ksu.edu.sa

How to cite this article: Alharthi, A.S., Alshamiry, F.A., Alghonaim, A.A., Al-Baadani, H.H. and Alhidary, I.A. (2025). Relationship between Forage Particle Size and Rate of *in vitro* Digestion Parameters of Alfalfa-based Lamb in Complete Pelleted Diets. Indian Journal of Animal Research. 1-9. doi: 10.18805/IJAR.BF-1923.

Submitted: 10-11-2024 **Accepted:** 16-12-2024 **Online:** 22-01-2025

the highest return at the lowest costs (Caihong *et al.*, 2023) will achieve sustainability in the production of red meat such as lamb (Zhong *et al.*, 2018; Foster *et al.*, 2023). Commercial livestock activities have expanded and intensive feeding has been used to produce meat from lambs such as total mixed rations (Pérez-Ruchel *et al.*, 2017), which are highly concentrated to achieve nutritional goals and the highest weight gain at the lowest possible

cost (Mahgoub *et al.*, 2000). In addition, the use of concentrated pellets instead of whole grains may improve the productive characteristics of growing lambs, such as growth rate and carcass quality (Alshamiry *et al.*, 2023).

Previous studies on the effect of forage particle size (FPS) considered one of the more modern methods confirm that it may affect the activity of the digestive system and microbial flora in the rumen (Ichihara and Fukubayashi, 2010; Qin *et al.*, 2023) and consequently absorption (Cui *et al.*, 2019; Foster *et al.*, 2023). Any change in FPS may lead to digestive disturbances and fermentation in the rumen. Alterations in FPS may result in digestive disturbances and fermentation inside the rumen (Chen *et al.*, 2021), the activity of the microbial flora in the rumen correlates with the surface area and the rate of forage material decomposition (Zebeli *et al.*, 2007; Weimer and Hall, 2020). Consequently, affecting the fattening rate and general performance of lambs (Francisco *et al.*, 2020). A comparison of ration substrates on rumen microbial fermentation does not depend solely on fiber content (Mirheidari *et al.*, 2020), although it is necessary to maintain the health, dynamism and activity of the rumen and to maintain the internal environment for the functioning of different types of bacterial communities (Letourneau *et al.*, 2024).

Understanding the biomass accumulation in ruminants is critical for improved earnings and long-term livestock production (Ekeocha *et al.*, 2023). Similarly, the protein and fat content of the rations, FPS and forage particle length (FPL) and the rate of cutting and grinding, are important factors (Li *et al.*, 2021).

According to Aissa *et al.* (2021), the maturation stage of forage material can impact digestibility and gas generation by influencing the percentage of dry matter and nutritional content, including protein and fiber.

The nutritional value of animal forage depends on its chemical composition, digestibility coefficient, fiber content and particle size (López *et al.*, 2005). To achieve this, there are many ways to estimate nutritional value but *in vitro* methods may be the best (Foster *et al.*, 2023) and have been used to estimate nutrient concentrations, digestibility (Gomide *et al.*, 2023) and the extent of particle size effects. Furthermore, the ratio of concentrate to roughage may influence fermentation efficiency, gas production and microbial protein synthesis (Zhang *et al.*, 2022). To predict digestion efficiency and make assumptions about the feed used, nutritional value and quality should be compared with *in vivo* experiments (Foster *et al.*, 2023).

In vitro techniques are also considered simple and relatively inexpensive compared to other forage evaluation methods (Vinyard and Faciola, 2022). Subsequently, there is a need to know more about the effects of the FPS and FPL on the intake, growth rate and refusal and wasted forage. Therefore, this study aimed to evaluate the relationship among FPL, degradability of feed *in vitro* and digestibility of organic materials.

MATERIALS AND METHODS

Animal welfare and ethics clearance

This study was undertaken at King Saud University (KSU), Riyadh, Saudi Arabia and the animals and procedures adopted in this study were in accordance with the Animal Welfare Act of Practice for the Care and Use of Animals for Scientific Purposes and approved by the Research Ethics Committee, KSU (Ethics Reference No: KSU-SE-22-26).

Animals and experimental design

A total of 40 Naimi male lambs, with an average body weight of 23 ± 2 kg and ~ 3 months old, were used in the 84-day trial. Initially, there was a 14-day adaptation period. The lambs were weighed, ear-tagged, vaccinated against clostridial diseases and treated for internal and external parasites as well as other vaccines against the most common infectious diseases in Saudi Arabia. Thereafter, the animals were randomly divided into four groups of 24 lambs each and housed in shaded pens, provided with a feed trough and a 10-L plastic water bucket. All lambs were offered the same complete pelleted diet (the basal diet) mostly corn and alfalfa hay containing 2.7 Mcal of MEM and 13.5% CP/kg. Dry matter (DM) content at maintenance level (2.5% of the initial body weight) was measured twice daily at 08:00 and 15:00. On day 1 of the experimental period, lambs were randomly assigned to one of four dietary treatments (four groups in each treatment): complete pelleted diet contained chopped alfalfa hay with lengths either 1 mm (the basal diet; PSL₁; control), 5 mm ((PSL₅), 10 mm (PSL₁₀) or 15 mm (PSL₁₅). Forage particle size is controlled industrially by controlling the distance between the drills or knives by the Mesh size and it is difficult to achieve completely equal sizes, but it represents 70% of the target according to the Pennsylvania scale.

All experimental diets were formulated in the facility at the Experimental Station of Animal Production Department, College of Food and Agriculture Sciences, King Saud University, to meet or exceed the nutritional requirements (National Research Council, 2007) of growing lambs and then provided as complete pelleted diets (Table 1).

Measurements and sample collection

Feeding analysis

Feed from each treatment was sampled before the study and monthly during the study and samples were frozen at -4°C for analysis. At the end of the study, feed samples were pooled (5%) and analyzed for nutrient composition at KSU laboratories. DM and ash content were determined by drying samples in an oven at 100°C for 24 h and by incinerating samples at 550°C for 3 h in a muffle furnace, respectively. The crude protein content of each sample was measured using an elemental analyzer. Neutral detergent fiber (NDF) and acid detergent fiber (ADF) were determined according to methods described by (Van Soest *et al.*, 1991) and AOAC method no. 973.18 C (2000), respectively.

Rumen fermentation profile

Rumen fluid samples (~50 mL) from six lambs/treatment were collected using an oral stomach tube before the morning feeding (0 h) and 3 h after feeding on day 45. The samples were analyzed immediately for pH using a microprocessor pH-meter (Model pH 211, Hanna® Instruments, Woonsocket, RI, USA) and then strained through four layers of cheesecloth. They were then transferred into plastic tubes and acidified with 2 mL of concentrated sulfuric acid (H₂SO₄) and frozen at -20°C for further analysis. The samples were then thawed and analyzed for ammonia (NH₃-N) using a TECO® Diagnostics kit and total proportion volatile fatty acids (VFAs) as described by (Jenkins, 1987), using 2-ethylbutyric acid (EBA) as an internal standard and gas chromatography mass spectrometry.

In vitro batch culture fermentation measurements

Feed samples were collected, dried to process DM and stored in sterile bags at 4°C until apparent digestibility could be estimated using the method of (Prachansuwan *et al.*, 2019). In four ANKOM gas jars per treatment as replicates (20 gas jars/treatment), 70 mL of strained rumen fluid was combined with 130 mL of pre-warmed buffer medium as batch rumen cultures. Finely powdered feed (200 g; based on DM) was weighed in a Dacron bag (Ankom Inc., Fairport, NY, USA), added to each ANKOM jar and incubated for 24 h to determine the apparent digestibility of DM, OM, ADF and NDF. The samples and feed treatments were analyzed for DM, ash analysis for calculating OM (100_Ash) and NDF according to the methods of AOAC, 2000. All dietary treatment jars, which were previously sealed, were filled with CO₂ and then connected to a Tedlar gas collection bag (Santa®, CA, USA.). The jars were placed in a water bath (Thermo® Fisher Scientific, Model 2873.0, USA) at 39°C for 24 h until the release process of the gases in the collection bag was completed. The jars were shaken every 2 h for 30 s. All CO₂ GP values are expressed as milliliters per time (2 h) and total GP (24 h) (Johnson *et al.*, 2009).

VFAs and Ammonia-N (NH₃-N) in the culture jars were determined at the end of each experiment. Samples (5 mL) from each jar were taken and placed directly placed in an ice bath and then stored at -20°C until analysis. The samples were mixed with 1 mL of 25% metaphosphoric acid and centrifuged at 20,000×g for 10 min at 4°C to produce a clear supernatant. The supernatant (1 mL) was filtered with a PTFE syringe filter (0.2 µm) and transferred to a 1.5 mL glass chromatography vial (Agilent). Acetic acid (C2), propionic acid (C3), butyric acid (C4), isobutyric acid (iso-C4), valeric acid (Val) and isovaleric acid (iso-Val) were analyzed by gas chromatography (Shimadzu Scientific Instruments Inc., Columbia, MD, USA), using EBA as an internal standard (Ichihara and Fukubayashi, 2010). VFA separation was performed as described previously (Cui *et al.*, 2019). VFAs are expressed as mM/1.0 ml sample. A sample was collected from each of the jars and centrifuged

at 12,000×g for 15 min at 4°C to measure the ammonia N concentration (NH₃-N) using a spectrophotometer (Perkin® Elmer, Waltham, USA.). The pH values of the culture jar sample were determined directly during sampling using a digital pH meter (Hanna® Instruments pH 211; Woonsocket, RI, USA-2014).

Statistical analysis

Statistical analyses were performed using SAS software v. 9.4 (SAS Institute Inc., Cary, NY, USA-2022). The gas production dataset was scaled down to a 1-h resolution and nonlinear regression analysis was performed using the SAS MIXED procedure. Significant differences in estimated parameters of gas production kinetics among the substrates or treatments were assessed using likelihood ratio 95 % confidence intervals. The variables were compared among the four forage particle sizes using the Kruskal-Wallis test with the NPAR1_WAY procedure in the gas component dataset. *P*<0.05 was considered statistically significant and Duncan's multiple range test was used to analyze these statistical differences.

RESULTS AND DISCUSSION

In vitro digestion and fermentation

The estimated nutrient digestibility, pH, ruminal gas production and fermentation for dietary treatments with different lengths of alfalfa hay during the *in vitro* digestion

Table 1: Ingredients and chemical composition of the basal diet¹.

Item, unit	Contain
Ingredients, % of dietary DM	
Corn	34.89
Alfalfa hay	23.29
Soybean meal	9.13
Wheat bran	17.34
Wheat straw	6.76
Limestone	2.67
Molasses	2.12
Dicalcium phosphate	1.62
Salt	1.12
Acid buffer	0.91
Vitamins and minerals premix ²	0.15
Nutrient composition, DM basis	
Dry Matter, (%)	90.34
Ash, %	8.17
Crude protein, %	13.93
Ether extract, %	1.54
Neutral detergent fiber, %	30.21
Acid detergent fiber, %	14.11
Metabolizable energy, Mcal/kg	2.79

¹Alfalfa hay was chopped to either 1, 5, 10 or 15-mm lengths and incorporated into the complete pelleted diets.

²Contained per kg, 10000 IU vitamin A, 1000 IU, vitamin D, 20 IU, vitamin E, 300 mg Mg, 24 mg Cu, 0.6 mg Co, 1.2 mg I, 60 mg Mn, 0.3 mg Se, 60 mg Zn.

process are presented in Table 2 and Fig 1. The apparent OM digestibility was affected ($P<0.05$) by dietary treatment. The mixture of loose alfalfa hay with concentrate pellets (PSL₁₅) resulted in a decrease in apparent digestibility for OM of 70.84% ($P<0.01$) compared to those of other experimental diets. The total gas production over the 24 h incubation period and ruminal fermentation (VFAs or Ammonia-N) did not differ ($P<0.05$) between treatments and they were similar across treatments, averaging 201.25 mL, 33.39 mM and 8.22 mM, respectively. The ruminal pH value was greater ($P<0.01$) in PSL₁₀ and PSL₁₅ than in other treatments.

The addition of forage at lengths of 5-10 mm to the complete pelleted diet increased ($P<0.05$) Iso-valerate and valerate molar percentages, respectively, during the *in vitro* digestion compared to the PSL₁₀ diet. However, the PSL₁ and PSL₁₅ diets had intermediate iso-valerate and valerate molar percentages (Table 2).

Fiber digestibility

The mean ruminal fluid pH and fermentation profiles of growing lambs fed pelleted diets containing different staple lengths of alfalfa hay are shown in Table 3. The pH values of ruminal fluid, total VFA concentrations and proportions of the majority of individual VFAs in the ruminal fluid were similar across dietary treatments and did not differ ($P>0.05$) between treatments. Feeding in the group control diet (PSL₁) resulted in an increase ($P>0.05$) in butyric acid molar percentage of VFAs (9.17%) compared with other groups.

In terms of polynomial contrast effects, there was a quadratic response ($P>0.05$) found in propionic and acetic acid molar percentages, whereas a linear response was observed ($P>0.05$) in acetate to propionate ratio as alfalfa hay length increased (Table 3 and Fig 1).

In vitro fermentation may play an important role in the study of feed efficiency and highlight the differences between feed types that affect the quantity and quality of feed available to ruminants (Bueno *et al.*, 2020; Vastolo *et al.*, 2022). Estimates and measurements related to the kinetics of nutrient decomposition in the rumen of ruminants are important indicators through which nutrients are classified as well as their carbohydrate and protein contents. This may then aid our understanding of digestion rates (Basto-Silva *et al.*, 2022; Paixão *et al.*, 2022). Compatibility between the decomposition of carbohydrates and nitrogen in the rumen increases the level of microbial protein synthesis and reduces energy loss during fermentation in the rumen. Accordingly, through laboratory digestion, they may be able to predict animal performance based on the nutritional composition of the feed (Santos *et al.*, 2016; Wei *et al.*, 2021).

Alfalfa feed is one of the most common types of feed used when manufacturing pelleted feed because of its high protein content (Al-Saiady *et al.*, 2010; Blume *et al.*, 2021; Srisaikhram and Rupitak, 2021) and palatability in animals (Vahdani *et al.*, 2014). To determine the nutritional value of alfalfa with the rest of the feeds included in the formulation

Table 2: Nutrient digestibility and fermentation characteristics produced from the *in vitro* incubation of dietary treatments with rumen fluid of growing lambs fed pelleted diets containing different staple lengths of alfalfa hay.

Variables, unit ²	Dietary treatments ¹				SEM	P value
	SLF ₁	SLF ₅	SLF ₁₀	SLF ₁₅		
DM digestibility, %	78.73	81.84	79.64	77.12	1.37	0.20
OM digestibility, %	77.13 ^a	80.45 ^a	78.51 ^a	70.84 ^b	1.50	0.01
NDF digestibility, %	57.31	55.52	58.14	55.49	2.83	0.88
Ruminal pH	6.50 ^b	6.51 ^b	6.58 ^a	6.61 ^a	0.04	0.01
Total gas production, mL	199	209	205	192	5.41	0.22
CO ₂ , ml	155	165	160	150	4.23	0.16
CH ₄ , ml	40.44	41.40	41.95	39.58	1.10	0.48
H ₂ , ml	2.92 ^a	1.99 ^b	2.97 ^a	2.10 ^b	0.07	0.01
Total VFA, mM	33.55	37.29	32.07	32.79	2.21	0.40
Acetate	15.83	17.47	15.31	15.33	0.91	0.35
Propionate	8.91	9.87	8.50	8.90	0.71	0.59
Butyrate	4.72	5.33	4.41	4.39	0.02	0.29
Iso-Butyrate	0.42	0.47	0.39	0.43	0.36	0.29
Iso-Valerate	0.81 ^{ab}	0.93 ^a	0.74 ^b	0.88 ^{ab}	0.04	0.02
Valerate	0.64 ^{ab}	0.76 ^a	0.60 ^b	0.71 ^{ab}	0.04	0.04
Acetate: Propionate ratio	1.78	1.77	1.80	1.72	0.05	0.81
Ammonia-N, mM	7.30	7.89	8.15	9.50	0.55	0.11

^{a,b} Within a row, means without a common superscript differ ($P<0.05$). ¹SLF₁ = A complete pelleted diet contained 1 mm length of alfalfa hay (Control); SLF₅ = A complete pelleted diet contained 5 mm length of alfalfa hay; SLF₁₀ = A complete pelleted diet contained 10 mm length of alfalfa hay; and SLF₁₅ = A complete pelleted diet contained 15 mm length of alfalfa hay. VFA = Volatile fatty acid.

when their lengths and sizes differ and to determine the amount of gas resulting from fermentation that may lead to fixed indicators and standards for evaluating the quality of manufactured feeds, this study showed that modification of FPL and FPS without changing the chemical composition may affect cumulative gas production (Zebeli *et al.*, 2008). The obtained results showed that the values of cumulative gas production in PSL₅ and PSL₁₀ lambs were higher compared with the other two groups, indicating that lengths greater than 5 and less than 15 mm affected the rate of gas production, which is consistent with the findings of Li *et al.* (2018). There is a relationship between the FPS and the time spent in the rumen; the larger the FPS, the more it

delays in the digestive system and the smaller the FPS, the faster it exits and in both cases it affects the efficiency of digestion, fermentation and decomposition of the feed, so logically it seems that the average FPS combines between them in achieving the appropriate time while increasing the rate of efficiency of forage utilization (Beauchemin *et al.*, 2003; Jiang *et al.*, 2019; Chelotti *et al.*, 2024). Rapid fermentation may increase gas build-up but simultaneously reduce nitrogen supply, which can negatively affect microbial activity (Magalhães *et al.*, 2019). FPL effects on rumen parameters had no effect on the pH of rumen fluid at different time points, which is consistent with the results reported by (Costa *et al.*, 2017) and (Santos-




Fig 1: Cumulative CO₂ gas production during the *in vitro* digestion for FPS from 2 to 24 hours of incubation (the x-axis).

Table 3: Effects of feeding different staple lengths of alfalfa hay on pH values of rumen fluid and ruminal fermentation characteristics of growing lambs fed pelleted diets.

Measurement, unit ²	Dietary treatments ¹				SEM	<i>P</i> value
	SLF ₁	SLF ₅	SLF ₁₀	SLF ₁₅		
Ruminal pH						
Before feeding, 0 h	6.19	6.17	6.29	6.69	0.17	0.14
After feeding, 3 h	5.99	5.95	5.93	5.90	0.09	0.91
VFA composition, %						
Total VFA, mM	32.86	34.87	31.03	33.14	1.43	0.14
Lactic acid	42.01	43.42	43.31	44.49	0.98	0.34
Formic acid	1.57	1.65	1.62	1.76	0.17	0.07
Acetic acid	28.40	29.99	28.50	27.41	1.09	0.23
Propionic acid	18.84	17.60	18.63	19.08	0.87	0.18
Butyric acid	9.17 ^a	7.34 ^b	7.94 ^b	7.26 ^b	1.21	0.05
Acetate: Propionate ratio	1.51 ^b	1.70 ^a	1.53 ^b	1.44 ^b	0.12	0.02

^{a,b} Within a row, means without a common superscript differ ($P < 0.05$). ¹SLF₁ = A complete pelleted diet contained 1 mm length of alfalfa hay (Control); SLF₅ = A complete pelleted diet contained 5 mm length of alfalfa hay; SLF₁₀ = A complete pelleted diet contained 10 mm length of alfalfa hay and SLF₁₅ = A complete pelleted diet contained 15 mm length of alfalfa hay.. VFA = Volatile fatty acid.

Silva *et al.*, 2019). There were no significant differences in the main VFAs acetate, propionate and butyrate nor in gas productivity. OM digestibility was significantly higher in the PSL₁, PSL₅ and PSL₁₀ groups than in PSL₁₅ group. This may be due to the liberation of microbial flora activity as the length of the forage and the percentage of fibers in the ration affect the digestibility of OM, by shifting the flow of the hydrogenation pathways towards VFAs, which results in the liberation of part of the laboratory according to the reported (Wang *et al.*, 2018). As shown by (Shen *et al.*, 2017), digestibility increased when the length of the forage particles ranged from 8 to 12 mm, as in the PSL₈ and PSL₁₂ groups, this result is consistent with what the researchers reported (Vidya *et al.*, 2024). The hydrogen levels were significantly higher in the PSL₁ and PSL₁₀ groups than in the PSL₅ and PSL₁₅ groups. The higher hydrogen values may be due to the liberation of H₂ as the particle length and the percentage of fibers in the ratio affected the production of methane by shifting the H₂ flow towards propionate, resulting in the liberation of part of the hydrogen in the laboratory (Wang *et al.*, 2018; Bhatt *et al.*, 2023). (Ramirez *et al.*, 2016) stated that dietary modifications may modify rumen kinetics, which may affect the synthesis and deposition of fats in ruminants that feed on particles of different forage lengths. Efforts to control forage particle size, when formulating animal diets, may help reduce methane production, as well as improve animal productivity, by reducing the amount of waste, residue and feces and succeed in mitigating animal gas emissions (Lakhani *et al.*, 2023).

CONCLUSION

Based on feed size, the values of PSL₅ to PSL₁₀ showed that these feed types could be used for feeding growing lambs and small ruminants, which can be explained by the amount of time required for the microbial flora, its ability to deteriorate and decompose the feed and by increasing the surface area. However, further studies are necessary to determine the particle size and length of forage before the manufacture of complete pelleted feed in growing animal diets, the study furthermore recommends conducting it on *in vivo* animals and other breeds of sheep and goats. Further experimental studies are required, to investigate the effects of various particle sizes of different forage on succeed production parameters in ruminants.

ACKNOWLEDGEMENT

The authors extend their appreciation to the National Plan for Science, Technology and Innovation (MAARIFAH), King Abdulaziz City for Science and Technology, Kingdom of Saudi Arabia (award number 2-17-04-001-0038).

Author contributions

Faisal A. Alshamiry, Hani H. Al-Baadani and Ahmed A. Alghonaim: Methodology, formal analysis and writing original draft. Abdulrahman S. Alharthi, Faisal A. Alshamiry and conceptualization, methodology and data curation; Ibrahim A. Alhidary: Investigation, review and editing and

project administration. All the authors have read and agreed to the published version of the manuscript.

Funding

The authors extend their appreciation to the National Plan for Science, Technology and Innovation (MAARIFAH), King Abdulaziz City for Science and Technology, Kingdom of Saudi Arabia (award number 2-17-04-001-0038).

Institutional Review Board Statement

The study was conducted at King Saud University, Riyadh, Saudi Arabia and the use of animals and procedures adopted here were in accordance with the Animal Welfare Act of Practice for the Care and Use of Animals for Scientific Purposes and approved by the Research Ethics Committee of King Saud University (Ethics Reference No: KSU-SE-22-26).

Disclaimers

The views and conclusions expressed in this article are solely those of the authors and do not necessarily represent the views of their affiliated institutions. The authors are responsible for the accuracy and completeness of the information provided, but do not accept any liability for any direct or indirect losses resulting from the use of this content.

Informed consent

All animal procedures for experiments were approved by the Committee of Experimental Animal care and handling techniques were approved by the University of Animal Care Committee.

Conflict of interest

The authors declare that there are no conflicts of interest regarding the publication of this article. No funding or sponsorship influenced the design of the study, data collection, analysis, decision to publish, or preparation of the manuscript.

REFERENCES

- Aïssa, A., Mustapha, C.B., Alvir, M.R., Ayed, M. H., Znaïdi, I.E.A., Moujahed, N. and Kraiem, K. (2021). Relationship between *in vivo*, *in vitro* parameters and chemical composition to predict the nutritive value of some legume forages. *Agricultural Science Digest*. 41(2). doi: 10.18805/ag.D-252.
- Alexander, P., Brown, C., Arneth, A., Finnigan, J., Moran, D. and Rounsevell, M.D. (2017). Losses, inefficiencies and waste in the global food system. *Agricultural Systems*. 153: 190-200.
- Al-Saiady, M.Y., Abouheif, M.A., Aziz Makkawi, A., Ibrahim, H.A. and Al-Owaimer, A.N. (2010). Impact of particle length of alfalfa hay in the diet of growing lambs on performance, digestion and carcass characteristics. *Asian-Australasian Journal of Animal Sciences*. 23(4): 475-482.
- Alshamiry, F.A., Alharthi, A.S., Al-Baadani, H.H., Aljumaah, R.S. and Alhidary, I.A. (2023). Growth rates, carcass traits, meat yield and fatty acid composition in growing lambs under different feeding regimes. *Life*. 13(2): 409.

- AOAC International. (2000). Official methods of analysis of AOAC International. 17: 1-2).
- Basto-Silva, C., Enes, P., Oliva-Teles, A., Capilla, E. and Guerreiro, I. (2022). Dietary protein/carbohydrate ratio and feeding frequency affect feed utilization, intermediary metabolism and economic efficiency of gilthead seabream (*Sparus aurata*) juveniles. *Aquaculture*. 554: 738-182.
- Beauchemin, K.A., Colombatto, D., Morgavi, D.P. and Yang, W.Z. (2003). Use of exogenous fibrolytic enzymes to improve feed utilization by ruminants. *Journal of Animal Science*. 81(14_suppl_2): E37-E47.
- Bhatt, R.S., Sarkar, S., Soni, L. and Sahoo, A. (2023). Effect of supplementation of roasted oilseeds and phytochemical-rich herbages on nutrient utilization, methane emission and growth performance in finisher lambs. *Environmental Science and Pollution Research*. 30(33): 80094-80105.
- Blume, L., Hoischen-Taubner, S. and Sundrum, A. (2021). Alfalfa-a regional protein source for all farm animals. *Landbauforschung*. 71(1). doi: 10.3220/LBF161589 4157 000.
- Bueno, I.C., Brandi, R.A., Fagundes, G.M., Benetel, G. and Muir, J.P. (2020). The role of condensed tannins in the *in vitro* rumen fermentation kinetics in ruminant species: Feeding type involved? *Animals*. 10(4): 635.
- Caihong, H. (2023). Methods to improve the economic benefits of raising Mutton Sheep. *Agricultural and Forestry Economics and Management*. 6(2): 68-73.
- Chelotti, J.O., Martinez-Rau, L.S., Ferrero, M., Vignolo, L.D., Galli, J.R., Planisich, A.M. and Giovanini, L.L. (2024). Livestock feeding behaviour: A review on automated systems for ruminant monitoring. *Biosystems Engineering*. 246: 150-177.
- Chen, H., Wang, C., Huasai, S. and Chen (2021). Effects of dietary forage to concentrate ratio on nutrient digestibility, ruminal fermentation and rumen bacterial composition in Angus cows. *Scientific Reports*. 11(1): 170-23.
- Costa, M., Alves, S.P., Francisco, A., Almeida, J., Alfaia, C.M., Martins, S.V. and Bessa, R.J.B. (2017). The reduction of starch in finishing diets supplemented with oil does not prevent the accumulation of trans-10 18: 1 in lamb meat. *Journal of Animal Science*. 95(8): 3745-3761.
- Cui, K., Qi, M., Wang, S., Diao, Q. and Zhang, N. (2019). Dietary energy and protein levels influenced the growth performance, ruminal morphology and fermentation and microbial diversity of lambs. *Scientific Reports*. 9(1): 166-12.
- Ekeocha, A.H., Aganga, A.A., Odumboni, A.A. and Makinwa, O.M. (2023). Growth Rate and Biomass Accumulation in Forage Maize (*Zea mays*), Forage Millet (*Echinochloa utilis*), Elephant Grass (*Pennisetum purpureum*) and Gamba Grass (*Andropogon gayanus*). *Indian Journal of Agricultural Research*. 57(4). doi:10.18805/IJArE.AF-672.
- Fanzo, J. and Miachon, L. (2023). Harnessing the connectivity of climate change, food systems and diets: Taking action to improve human and planetary health. *Anthropocene*. 42: 100-381.
- Foster, J.L., Smith, W.B., Rouquette, F.M. and Tedeschi, L.O. (2023). Forages and pastures symposium: an update on *in vitro* and *in situ* experimental techniques for approximation of ruminal fiber degradation. *Journal of Animal Science*. 101: skad097.
- Francisco, A.E., Janiëek, M., Dentinho, T., Portugal, A.P., Almeida, J.M., Alves, S.P. and Santos-Silva, J. (2020). Effects of alfalfa particle size and starch content in diets on feeding behaviour, intake, rumen parameters, animal performance and meat quality of growing lambs. *Meat Science*. 161: 107-964.
- Galanakis, C.M. (2023). The “vertigo” of the food sector within the triangle of climate change, the post-pandemic world and the Russian-Ukrainian war. *Foods*. 12(4): 721.
- Giamouri, E., Zisis, F., Mitsiopolou, C., Christodoulou, C., Pappas, A.C., Simitzis and Tsiplakou, (2023). Sustainable strategies for greenhouse gas emission reduction in small ruminants farming. *Sustainability*. 15(5): 41-18.
- Gomide, DR., Pereira, R.A., Silva, R.B., Carvalho, J.T., Lara, M.A. and Pereira, M.N. (2023). Effect of particle size of silage of flint corn grain on dairy cows fed tropical pasture: Performance, intake, ruminal fermentation and digestibility. *Animals*. 13(12): 19-32.
- Ichihara, K.I. and Fukubayashi, Y. (2010). Preparation of fatty acid methyl esters for gas-liquid chromatography [S]. *Journal of Lipid Research*. 51(3): 635-640.
- Izadbakhsh, M. H., Hashemzadeh, F., Alikhani, M., Ghorbani, G.R., Khorvash, M., Heidari, M. and Ahmadi, F. (2024). Effects of dietary fiber level and forage particle size on growth, nutrient digestion, Ruminal fermentation and behavior of Weaned Holstein Calves under heat stress. *Animals*. 14(2): 275.
- Jararweh, Y., Fatima, S., Jarrah, M. and AlZu'bi, S. (2023). Smart and sustainable agriculture: Fundamentals, enabling technologies and future directions. *Computers and Electrical Engineering*. 110: 108-799.
- Jenkins, T.C. (1987). Effect of fats and fatty acid combinations on ruminal fermentation in semi-continuous *in vitro* cultures. *Journal of Animal Science*. 64(5): 1526-1532.
- Jiang, F.G., Lin, X.Y., Yan, Z.G., Hu, Z.Y., Wang, Y. and Wang, Z.H. (2019). Effects of forage source and particle size on chewing activity, ruminal pH and saliva secretion in lactating Holstein cows. *Animal Science Journal*. 90(3): 382-392.
- Johnson, M.C., Devine, A.A., Ellis, J.C. and Grunden, A. (2009). Effects of antibiotics and oil on microbial profiles and fermentation in mixed cultures of ruminal microorganisms. *Journal of Dairy Science*. 92(9): 4467-4480.
- Lakhani, P., Kumar, R., Madan, J. and Sindhu, S. (2023). Ruminant methane mitigation-strategy from bench top to field condition: A review. *Agricultural Reviews*. 44(2): 182-190. doi: 10.18805/ag.R-2152.
- Letourneau, J., Carrion, V.M., Zeng, J., Jiang, S., Osborne, O.W., Holmes, Z.C. and David, L.A. (2024). Interplay between particle size and microbial ecology in the gut microbiome. *The ISME Journal*. 18(1): wræ168.

- Li, Y., Guo, Y.L., Zhang, C.X., Cai, X.F., Liu, P. and Li, C.L. (2021). Effects of physical forms of starter feed on growth, nutrient digestibility, gastrointestinal enzyme activity and morphology of pre-and post-weaning lambs. *Animal*. 15(1): 100-044.
- Li, Z., Liu, N., Cao, Y., Jin, C., Li, F., Cai, C. and Yao, J. (2018). Effects of fumaric acid supplementation on methane production and rumen fermentation in goats fed diets varying in forage and concentrate particle size. *Journal of Animal Science and Biotechnology*. 9: 1-9.
- López, S., Davies, D.R., Giráldez, F.J., Dhanoa, M.S., Dijkstra, J. and France, J. (2005). Assessment of nutritive value of cereal and legume straws based on chemical composition and *in vitro* digestibility. *Journal of the Science of Food and Agriculture*. 85(9): 1550-1557.
- Magalhães, A.L.R., Teodoro, A.L., Gois, G.C., Campos, F.S., Souza, J.D.S.R. andrade, A.P. and Nascimento, D.B.D. (2019). Chemical and mineral composition, kinetics of degradation and *in vitro* gas production of native cactus. *Journal of Agricultural Studies*. 7(4): 119-137.
- Mahgoub, O., Lu, C.D. and Early, R.J. (2000). Effects of dietary energy density on feed intake, body weight gain and carcass chemical composition of Omani growing lambs. *Small Ruminant Research*. 37(1-2): 35-42.
- Mirheidari, A., Torbatinejad, N.M., Shakeri, P. and Mokhtarpour, A. (2020). Effects of biochar produced from different biomass sources on digestibility, ruminal fermentation, microbial protein synthesis and growth performance of male lambs. *Small Ruminant Research*. 183: 106-042.
- National Research Council. (2007). *Nutrient Requirements of Small Ruminants: Sheep, Goats, Cervids and New World Camelids*. National Academy Press.
- Oecd, F.A.O. (2022). *OECD-FAO Agricultural Outlook. 2022-2031*.
- Paixão, T.R., de Almeida, V.V.S., Oliveira, A.C., da Silva, A.P.G., Silva, J.W.D., Santos, L.V. and Silva, R.R. (2022). Intake, digestibility, ruminal parameters and performance in lamb fed with increasing levels of red propolis extract. *Tropical Animal Health and Production*. 54(6): 364.
- Pérez-Ruchel, A., Repetto, J.L. and Cajarville, C. (2017). Supplementing high-quality fresh forage to growing lambs fed a total mixed ration diet led to higher intake without altering nutrient utilization. *Animal*. 11(12): 2175-2183.
- Prachansuwan, A., Kriengsinyos, W., Judprasong, K., Kovitvadi, A. and Chundang, P. (2019). Effect of different pre-boiling treatment on *in vitro* protein and amino acid digestibility of mung beans [*Vigna radiata* (L.) Wilczek]. *Malaysian Journal of Nutrition*. 25(3): 361-375.
- Qin, M., Su, C.W., Umar, M., Lobonp, O.R. and Manta (2023). Are climate and geopolitics the challenges to sustainable development? Novel evidence from the global supply chain. *Economic Analysis and Policy*. 77: 748-763.
- Ramirez, H.R., Harvatine, K.J. and Kononoff, P.J. (2016). Forage particle size and fat intake affect rumen passage, the fatty acid profile of milk and milk fat production in dairy cows consuming dried distillers grains with solubles. *Journal of Dairy Science*. 99(1): 392-398.
- Santos, S., Rotta, P.P., Costa, E., Silva, L., Menezes, A.C.B., Pina, D. and Valadares Filho, S. (2016). Protein ruminal degradation of feeds and microbial protein synthesis. *Nutrient Requirements of Zebu and Crossbred Cattle*. 43.
- Santos-Silva, J., Francisco, A., Alves, S.P., Portugal, P., Dentinho, T., Almeida, J. and Bessa, R.J. (2019). Effect of dietary neutral detergent fibre source on lambs growth, meat quality and biohydrogenation intermediates. *Meat Science*. 147: 28-36.
- Shen, J., Liu, Z., Yu, Z. and Zhu, W. (2017). Monensin and nisin affect rumen fermentation and microbiota differently *in vitro*. *Frontiers in Microbiology*. 8: 11-11.
- Srisaikhram, S. and Rupitak, Q. (2021). A preliminary study on growth, yield and nutritive value of four varieties of alfalfa and the utilization of alfalfa dehydrated pellets in a total mixed ratio in meat goat diet. *CMUJ. Nat. Sci.* 20(1): e202-1003.
- Vahdani, N., Moravej, H., Rezayazdi, K., Dehghan-Banadki, M. (2014). Evaluation of Nutritive Value of Grass Pea Hay in Sheep Nutrition and Its Palatability as Compared with Alfalfa. *JAST*. 16 (3): 537-550.
- Van Soest, P.V., Robertson, J.B. and Lewis, B.A. (1991). Methods for dietary fiber, neutral detergent fiber and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science*. 74(10): 3583-3597.
- Vastolo, A., Matera, R., Serrapica, F., Cutrignelli, M.I., Neglia, G., Kiatti, D.D. and Calabrò, S. (2022). Improvement of rumen fermentation efficiency using different energy sources: *In vitro* comparison between Buffalo and cow. *Fermentation*. 8(8): 351.
- Vidya, B., Venkateshwarlu, M., Nagalakshmi, D., Chandra, A.S., Preetham, V.C. and Kumari, N.N. (2024). Rumen Fermentation and Microbial Nitrogen Supply in Native Sheep Fed Crop Residue based Densified Feed Blocks Varying in Particle Size of Roughage. *Asian Journal of Dairy and Food Research*. 43(1). doi:10.18805/ajdfr.DR-1896.
- Vinyard, J.R. and Faciola, A.P. (2022). Unraveling the pros and cons of various *in vitro* methodologies for ruminant nutrition: A review. *Translational Animal Science*. 6(4): txac130.
- Wang, K., Nan, X., Chu, K., Tong, J., Yang, L., Zheng, S. and Xiong, B. (2018). Shifts of hydrogen metabolism from methanogenesis to propionate production in response to replacement of forage fiber with non-forage fiber sources in diets *in vitro*. *Frontiers in Microbiology*. 9: 27-64.
- Wei, Z., Xie, X., Xue, M., Valencak, T.G., Liu, J. and Sun, H. (2021). The effects of non-fiber carbohydrate content and forage type on rumen microbiome of dairy cows. *Animals*. 11(12): 35-19.
- Weimer, P.J. and Hall, M.B. (2020). The potential for biomimetic application of rumination to bioreactor design. *Biomass and Bioenergy*. 143: 105-822.
- World Health Organization. (2022). *The state of food security and nutrition in the world 2022: Repurposing food and agricultural policies to make healthy diets more affordable* (Vol. 2022). Food and Agriculture Org.
- Yan, X., Ying, Y., Li, K., Zhang, Q. and Wang, K. (2024). A review of mitigation technologies and management strategies for greenhouse gas and air pollutant emissions in livestock production. *Journal of Environmental Management*. 352: 120-028.

- Zebeli, Q., Tafaj, M., Weber, I., Dijkstra, J., Steingass, H. and Drochner, W. (2007). Effects of varying dietary forage particle size in two concentrate levels on chewing activity, ruminal mat characteristics and passage in dairy cows. *Journal of Dairy Science*. 90(4): 1929-1942.
- Zebeli, Q., Tafaj, M., Weber, I., Steingass, H. and Drochner, W. (2008). Effects of dietary forage particle size and concentrate level on fermentation profile, *in vitro* degradation characteristics and concentration of liquid-or solid-associated bacterial mass in the rumen of dairy cows. *Animal Feed Science and Technology*. 140(3-4): 307-325.
- Zhang, Z., Wang, S., Qi, R., Shahzad, K., Shi, L., Zhang, X. and Wang, M. (2022). *In vitro* study of *Urtica cannabina* and *Leymus chinensis* on rumen microbial fermentation and gas production. *Indian Journal of Animal Research*. 56(2): 187-191. doi:10.18805/IJAR.B-1291.
- Zhong, R.Z., Fang, Y., Zhou, D.W., Sun, X.Z., Zhou, C.S. and He, Y.Q. (2018). Pelleted total mixed ration improves growth performance of fattening lambs. *Animal Feed Science and Technology*. 242: 127-134.