



Ameliorative Effects of Dandelion Root Extract (*Taraxacum officinale*) against Silver Nanoparticles-stimulated Hepatotoxicity in Male Albino Rats

Rabab Mohamed Aljarari¹, Safa H. Qahl¹, Sarah Ayidh Al-Sulami¹, Reem Yahya Alzahri¹

10.18805/IJAR.BF-1956

ABSTRACT

Background: Dandelion roots have considerable health benefits including promoting liver health and combating inflammation. Therefore, our study investigated the impacts of dandelion root extract (*Taraxacum officinale*) on AgNPs-triggered hepatotoxicity in rats.

Methods: Herein, 48 adult male albino Wistar rats were randomly distributed into six groups (8 rats/group). As controls, rats from the first group were employed. *T. officinale* (500, 250 mg/kg b.w./day) was administered orally to the rats of the second and third groups. The AgNPs (1 mg/kg b.w./day) were intraperitoneally injected into the fourth group. Rats of the fifth and sixth groups were treated with (*T. officinale* 500+ AgNPs) and (*T. officinale* 250+ AgNPs). After four weeks, biochemical, histopathological and immunohistochemical evaluations were conducted on the blood samples and liver tissues.

Result: Here, we have manifested that exposure to AgNPs significantly elevated liver enzyme levels alkaline phosphatase (ALP), aspartate aminotransferase (AST), alanine aminotransferase (ALT) and lactate dehydrogenase (LDH), total protein, albumin, Malondialdehyde (MDA) and caspase-3 whereas total bilirubin levels, superoxide dismutase (SOD) and glutathione (GSH), revealed significant decline linked to hepatic damage. Treatment of rats with *T. officinale* showed a pronounced attenuation of the damage caused by AgNPs associated with improvement of biochemical, hepatic histopathological and immunohistochemical alterations. This study shows that sub-chronic oral administration of *T. officinale* in rats protects against AgNPs-stimulated hepatotoxicity. The positive implications may be ascribed to the antioxidant capabilities of *T. officinale*.

Key words: Antioxidants, Dandelion roots (*Taraxacum officinale*), Hepatotoxicity, Oxidative stress, Silver nanoparticles.

INTRODUCTION

Nanoparticles (NPs) are at the forefront of the quickly evolving field of nanotechnology and have attracted significant attention for their use in both the treatment of cancer and as an efficient drug delivery method (Albrahim and Alonazi, 2020). Compared to the original bulk substance, these NPs' average size falls between 1 and 100 nm (Ema *et al.*, 2017). These sizes have distinct chemical and physical characteristics that lead to advanced magnetic, electrical, optical, mechanical and structural qualities (Lekha *et al.*, 2021). Silver NPs (AgNPs) have attracted attention among the many current nanomaterials because of their distinctive physical and biochemical characteristics (Zhang *et al.*, 2016). Also, their exceptional antibacterial activity makes them beneficial for applications in many areas, including biomedicines (Lee and Jun, 2019). However, their concentration in the environment is unknown and they pose a potential threat to the human body organs, including the liver (Xiong *et al.*, 2022). The implications of AgNP accumulation in the liver are inflammation, necrosis and apoptosis (Al-Doaiss *et al.*, 2020).

Oxidative stress is an essential mechanism of AgNP hepatotoxicity, reactive oxygen species (ROS) generation and proinflammatory cytokines secretion (Mishra *et al.*, 2016; Blanco *et al.*, 2018; Mao *et al.*, 2018). Salama and her colleagues manifested that oxidative stress, inflammation and ultimately cellular death can be the result of elevated

¹Department of Biological Science, College of Science, University of Jeddah, Jeddah, Saudi Arabia.

Corresponding Author: Rabab Mohamed Aljarari, Department of Biological Science, College of Science, University of Jeddah, Jeddah, Saudi Arabia. Email: rmaljarary@uj.edu.sa

How to cite this article: Aljarari, R.M., Qahl, S.H., Al-Sulami, S.A. and Alzahri, R.Y. (2025). Ameliorative Effects of Dandelion Root Extract (*Taraxacum officinale*) against Silver Nanoparticles-Stimulated Hepatotoxicity in Male Albino Rats. Indian Journal of Animal Research. 1-11. doi: 10.18805/IJAR.BF-1956.

Submitted: 12-01-2025 **Accepted:** 28-02-2025 **Online:** 15-05-2025

ROS levels, which disrupt intracellular redox balance (Salama *et al.*, 2023). Excessive synthesis of the ROS may lead to heightened lipid peroxidation, damage of mitochondria and apoptosis, beyond the capacity of natural cellular antioxidant defense mechanisms, But the body has a strategy by which it tries to overcome ROS and defend the body through antioxidants the antioxidant defense system and the most important antioxidant enzymes GSH and SOD, which in turn reduce the toxic effects (Flores-López *et al.*, 2019).

Recent years have seen a significant increase in interest in natural product-based medications for preventing and treating liver conditions (Amin *et al.*, 2016; Ayaz *et al.*, 2017; Hamza *et al.*, 2018; Khajuria *et al.*, 2018; Ashktorab *et al.*, 2019). The dandelion *Taraxacum officinale*

(*T. officinale*) is a perennial herbaceous plant in the Asteraceae family. In many nations, it has been employed as a system for both traditional and contemporary herbal treatment. It has been utilized as a folk remedy for treating chronic liver disorders in many different nations. Moreover, it exhibited numerous health advantages because of its diuretic (R̄cz-Kotilla *et al.*, 1974) and anti-oxidant *in vivo* (Choi *et al.*, 2010) and *In vitro* (Park *et al.*, 2011; Ivanov *et al.*, 2018; Milek *et al.*, 2019), anticancer (Saratale *et al.*, 2018), such as human leukemia, pancreatic, colorectal and prostate cancer cells (Nguyen *et al.*, 2019; Ovadje *et al.*, 2011; Ovadje *et al.*, 2012; Ovadje *et al.*, 2016) and inflammatory (Di Napoli and Zucchetti, 2021). In the Arabian regions, dandelion *T. officinale*, an edible plant, has been utilized as a traditional herbal remedy to cure liver disorders (Schutz *et al.*, 2006). *T. officinale* contains flavonoids and polyphenolic compounds, including sesquiterpene lactones, taraxerol and chlorogenic and chicoric acids, as well as vitamins A, B, C, D, E, lecithin, inositol and minerals like sodium, iron, magnesium, copper, zinc, calcium, phosphorus, silicon and manganese (Pereira *et al.*, 2016; Qadir *et al.*, 2022). It has been shown that the root extract of *T. officinale* may offer protection against certain toxic liver injuries (Cai *et al.*, 2017).

Therefore, we aimed to ascertain the ameliorative impact of *T. Officinale* against AgNPs-triggered hepatotoxicity and explore the underlying mechanisms.

MATERIALS AND METHODS

Dandelion (*T. officinale*) root extract was supplied as herbal supplement capsules (800 mg/capsule) and purchased from (Herbal Factors Company, Coquitlam, Canada). Each capsule was suspended in 10 ml of distilled water shortly before administration (Park *et al.*, 2010). Phosphate buffered saline (PBS) was obtained from the pharmaceutical Solutions Industry, Jeddah, Saudi Arabia). Diethyl ether was obtained from Sigma-Aldrich (GMBH, Munich, Germany). Formaldehyde was purchased from (Riedel-Detain, Sleaze, Germany). The ALT, AST, ALP, LDH, albumin, GSH, SOD, MDA, total bilirubin and protein, were evaluated by Enzyme-Linked Immunosorbent Assay (ELISA) kits (My BioSource, San Diego, U.S.A).

In brief, a cold bath maintained at 6°C to 10°C was used to combine 185 mL of type 1 water (Milli Q) with 5 mL of both 0.05 M sodium citrate (TSC, Sigma Aldrich CAS 6132-04-3) and silver nitrate (AgNO₃, PANREAC CAS 7761-88-8). The mixture was centrifugated at 3000 RPM for a duration of 3 minutes. Afterward, 5 mL of 0.05 M sodium borohydride (NaBH₄, Sigma-Aldrich CAS 16940-66-2) was progressively introduced. Typically, 1.25 M sodium hydroxide (NaOH, PANREAC CAS 1310-73-2) was employed to adjust the pH to 10. The resulting nanoparticles (NPs) were stored in amber bottles at 4°C and were utilized as mentioned in a previous study (Quintero-Quiroz *et al.*, 2019).

The samples' size and morphology were verified with transmission electron microscopy (TEM) utilizing a Tecnai F20 Super Twin Turbo-molecular Pump (TMP) and a Tecnai

G2 200KV TEM (FEI). The samples were generated by depositing a drop of each suspension, about 60 nm thick, onto a carbon membrane (Monteiro *et al.*, 2011) (Fig 1). The sample was examined employing an X-ray diffractometer (XRD) using CuK α radiation, scanning a 2 θ range from 10° to 80° (Zhang *et al.*, 2016) (Fig 2).

Here, male albino Wistar strain rats (*Rattus norvegicus*), with a weight range of 140-180 g were employed. The rats were acquired from the experimental animal facility of the Faculty of Pharmacy at King Abdulaziz University, Jeddah, Saudi Arabia. All methods received approval from the Animal Care and Use Committee at King Abdul-Aziz University, Faculty of Pharmacy, Jeddah, Saudi Arabia (Approval number: PH-1444-16). Rats were acclimatized for 7 days before the beginning of the experiment. Standard plastic (polypropylene) cages were used to house the rats, which were maintained in controlled laboratory conditions with a humidity of 65%, a temperature of 20 $\pm$ 1°C and a 12:12 h light / dark cycle. Throughout the acclimation phase, the rats were fed daily with normal commercial chow ad libitum and given unrestricted access to water.

Typically, 48 rats were allocated into six experimental groups (n = 8) and were treated as follows:

- Group 1: (Negative control) received an oral administration of saline solution (0.9% NaCl) for four weeks daily.
- Group 2: received *T. officinale* root aquatic extract (500 mg/kg b.w.) orally daily for four weeks.
- Group 3: received *T. officinale* root aquatic extract (250 mg/kg b.w.) orally daily for four weeks.
- Group 4: (positive control), was given AgNPs (1mg/kg b.w.) by intraperitoneal injection (I/P), daily for four weeks.
- Group 5: Received *T. officinale* root aquatic extract (500 mg/kg b.w.) and AgNPs (1mg/kg b.w.); with one-hour intervals between the two administrations; daily for four weeks.
- Group 6: Received *T. officinale* root aquatic extract (250 mg/kg b.w.) and AgNPs (1mg/kg b.w.) fo.; with one-hour intervals between the two administrations; daily for four weeks.

The selected dose of AgNPs was selected based on previous research articles (Aboelwafa *et al.*, 2022; Assar *et al.*, 2022; Yousef *et al.*, 2022). Also, the two doses of *T. officinale* root aquatic extract were selected according to Domitrović *et al.* (2010); Fallah *et al.* (2010); Şenocak and Yıldırım (2017); Shaaban *et al.* (2023) for the low dose (250 mg/kg) and according to Fallah *et al.* (2010); Aremu *et al.* (2019); Hamza *et al.* (2020) for the high dose (500 mg/kg).

After the end of the experiment (at 38th day), rats were weighed and anaesthetized utilizing diethyl ether. Blood samples were obtained from the retro-orbital venous plexuses. The serum underwent centrifugation at 2500 rpm for 15 min and was thereafter stored for biochemical analysis at -80°C. The abdomen was dissected and then the liver was isolated, weighed and dissected into two parts. The first section was produced and fixed in 10%

formalin for histopathological and immunohistochemical analysis. A separate portion was homogenized in 4 volumes of phosphate buffer (pH 7.4) for biochemical examination. The liver-to-body weight index was determined by calculating the ratio of liver weight to the rat's final body weight after weighing both the body and liver. The ALT, AST, ALP, LDH, albumin, total bilirubin and protein levels in serum were ascertained by commercial kits, per protocols.

For biochemical evaluation of GSH, SOD and MDA, a part of liver homogenate was centrifuged at 12000 g at 4°C for 20 min. Afterwards, the supernatant was separated and stored at -80°C and the activities of oxidative stress markers were investigated by commercial kits, following the protocols.

Liver tissues fixed in formalin were embedded in paraffin blocks, sliced into 5-micrometer thickness, placed on glass slides and stained with hematoxylin and eosin (HandE) (Bancroft and Gamble, 2008). The prepared liver slices were examined using an Intellisite Ultra-Fast Scanner (Digital Pathology Slide Scanner, Philips FMT0225).

Furthermore, 5-µm formalin-fixed, paraffin-embedded slices were examined with immunohistochemical staining via the streptavidin-biotin technique utilizing caspase 3 antibodies as an indicator for programmed cell death (apoptosis), following the aforementioned protocol (Ghonimi *et al.*, 2022).

The statistical analysis was performed with the Statistical Package for Social Science (SPSS program for Windows, version 25). Data were expressed as mean $\pm$ standard error. The difference between different experimental groups was assessed with One Way ANOVA (Tukey test). P-value <0.05 is deemed statistically significant.

RESULTS AND DISCUSSION

Body weight

The initial body weight of different groups ranged from (140 gm to 181 gm). The smallest weight was in the *T. officinale* 250 group (140 gm) in comparison to the control rats (178 gm). However, AgNPs intoxicated rats, (*T. officinale* 500 + AgNPs) and (*T. officinale* 250 + AgNPs) treated rats showed




Fig 1: Transmission electron microscope (TEM) micrograph showed silver nanoparticles (AgNPs) with spherical shapes and monodispersed behaviour (Scale 100 nm).




Fig 2: Show the x-ray diffraction pattern of AgNPs with an average crystal particle size of 12 nm.

a substantial increment in their initial body weight compared to control. Moreover, no significant difference in ultimate body weight among the various groups was observed. However, the AgNPs intoxicated rats displayed a marked reduction in their body weight compared to other treated groups (Fig 3).

Liver weight index

As depicted in Fig 4 revealed a non-significant difference between different treated groups. However, the liver weight index of AgNP-intoxicated rats declined compared to the control group. Also, (*T. officinale* 500 + AgNPs) treated rats exhibited a reduction in the liver weight index. However, it was not significant.

Liver function analyses

The outcomes manifested that the (ALT, AST, ALP, LDH, albumin and total protein) levels were significantly raised in the serum of the AgNPs group, unlike the control rats ($P < 0.05$). Treatment with (*T. officinale* 500+ AgNPs) and (*T. officinale* 250+ AgNPs) significantly mitigated (ALT, AST,

ALP, LDH, albumin and total protein) levels contrasted with AgNPs group ($P < 0.05$). The total bilirubin level in serum revealed a significant decline in the AgNPs group, unlike the control rats ($P < 0.05$). Nevertheless, treatment with (*T. officinale* 500+ AgNPs) and (*T. officinale* 250+ AgNPs) caused significant increases in total bilirubin compared with the AgNPs group ($P < 0.05$; Table 1).

Liver oxidative stress markers

Here, we found that the AgNPs group experienced a significant decline in the GSH and SOD levels compared to the control rats ($P < 0.05$). Treatment with (*T. officinale* 500+ AgNPs) and (*T. officinale* 250+ AgNPs) caused significant increases in GSH and SOD levels compared with the AgNPs group ($P < 0.05$). The AgNPs group exhibited significantly elevated MDA levels in comparison with control rats ($P < 0.05$). Nevertheless, treatment of rats with (*T. officinale* 500+ AgNPs) and (*T. officinale* 250+ AgNPs) caused significant decreases in MDA levels compared with the AgNPs group ($P < 0.05$, Fig 5a-c).




Fig 3: Changes in body weight (g) after four weeks of different experimental groups. Data are mean $\pm$ SE of eight animals in each group.

Table 1: Levels of serum ALT, AST, ALP, LDH, Albumin, Total bilirubin and Total protein of different experimental groups.

Group	ALT (U/L)	AST (U/L)	ALP (U/L)	LDH (U/L)	Albumin (g/dl)	Total bilirubin (mg/dl)	Total protein (g/dl)
Control	18.30 $\pm$ 1.56	25.57 $\pm$ 2.06	47 $\pm$ 2.31	150.33 $\pm$ 8.51	6.53 $\pm$ 0.23	4.51 $\pm$ 0.45	0.37 $\pm$ 0.04
<i>T. officinale</i> 500	18.57 $\pm$ 2.23	22.17 $\pm$ 1.71	52.67 $\pm$ 6.77	163.33 $\pm$ 8.82	6.00 $\pm$ 0.26	4.11 $\pm$ 0.31	0.40 $\pm$ 0.09
<i>T. officinale</i> 250	12.37 $\pm$ 0.47	14.73 $\pm$ 0.94	47.33 $\pm$ 1.45	145.67 $\pm$ 2.96	5.87 $\pm$ 0.26	5.7 $\pm$ 0.76	0.81 $\pm$ 0.19
AgNPs	99.67 $\pm$ 12.78*	114.90 $\pm$ 13.62*	147.33 $\pm$ 8.09*	492 $\pm$ 28.18*	9.83 $\pm$ 0.26*	1.73 $\pm$ 0.23*	1.40 $\pm$ 0.09*
<i>T. officinale</i> 500 + AgNPs	14.40 $\pm$ 1.91#	17.57 $\pm$ 2.06#	52.67 $\pm$ 6.17#	150 $\pm$ 12.06#	5.9 $\pm$ 0.26#	4.57 $\pm$ 0.38#	0.61 $\pm$ 0.13#
<i>T. officinale</i> 250 + AgNPs	16.40 $\pm$ 2.51#	20.47 $\pm$ 2.42#	59.33 $\pm$ 9.39#	168.67 $\pm$ 10.84#	6.17 $\pm$ 0.41#	5.13 $\pm$ 0.47#	0.52 $\pm$ 0.16#

*Indicates a significant difference versus control. # Indicates a significant difference versus AgNPs.

Liver histopathological examination

Histopathological examination of liver sections is illustrated (Fig 6a-f). Histological studies revealed that the control group showed normal hepatic architecture (Fig 6a). The liver structure of the group treated with (*T. officinale* 500 mg) showed that the liver histology exhibited a structure comparable to that of the control group, except for minor histological alterations (Fig 6b). The structure of the liver treated with (*T. officinale* 250 mg) exhibited a resemblance to the control group, characterized by relatively normal hepatocytes with rounded vesicular nuclei and eosinophilic cytoplasm containing basophilic granules (Fig 6c). The

alterations in the liver of the AgNPs group demonstrated severe damage in the liver which was revealed as inflammatory, degenerative, necrotic and hyperplastic alterations. The AgNPs-intoxicated group showed obvious histopathological changes; these include degenerations of hepatocytes leaving empty spaces, dilated hyperemic and congested blood sinusoids, degenerated hepatocytes and some of them appeared with eosinophilic cytoplasm and deeply stained pyknotic nuclei. The portal tract demonstrated congested and dilated portal vein, mononuclear cell infiltrations and necrotic empty spaces left by degenerated hepatocytes (Fig 6d). In rats treated




Fig 4: Liver weight index (%) of different experimental groups. Data are mean±SE of eight animals in each group.




Fig 5(a-c): Levels of liver homogenate (a) GSH, (b) SOD, (c) MDA of different experimental groups.

with (*T. officinale* 500+AgNPs) and (*T. officinale* 250+AgNPs) the hepatic cellularity had significantly improved, according to light microscopic studies.

Immunohistochemical examination of Caspase-3

Immunostaining of caspase-3 in the liver manifested adverse caspase-3 expression in the control, (*T. officinale* 500) and (*T. officinale* 250) groups (Fig 7a-c). Conversely, the AgNPs group revealed positive caspase-3 expression in hepatic cells (Fig 7d). Moreover, mitigated caspase-3 expression was manifested in the liver of (*T. officinale* 500+AgNPs) group (Fig 7e). Also, moderate caspase-3 expression was revealed in the liver of rats treated with (*T. officinale* 250+AgNPs; Fig 7f).

The unique physicochemical features of AgNPs make them potentially useful in cosmetics, biosensing, imaging, home products, medicine and research laboratories (Assar *et al.*, 2022). The increasing usage of AgNPs in the last decade has resulted in several issues related to the

environment and human health (Syafiuddin *et al.*, 2017). Therefore, a greater understanding of AgNP toxicity and its underlying mechanisms is a must. Natural product-derived drugs have garnered significant interest in preventing and treating liver conditions (Li *et al.*, 2018). *T. officinale* was considered one of the beneficial plants in maintaining liver homeostasis (Ignat *et al.*, 2021).

Here, we aimed to ascertain the hepatoprotective implications of *T. officinale* versus AgNPs-triggered toxicity in adult male albino rats. Rats exposed to AgNPs showed significant body loss, liver function reduction, antioxidant defence system disruption and histopathological alterations of the liver architecture associated with extensive immunoreactivity against the anti-caspase-3 antibody.

Body weight change is a marker of a drug or any agent's substantial toxicity (Zainal *et al.*, 2020). Herein, our findings demonstrated that the body weight was decreased in rats intoxicated with AgNPs contrasted with the control group. Similar studies manifested that exposure to AgNPs can




Fig 6 (A-F): Photomicrograph sections in the hepatic tissue of rats (Hand E; X400- scale bar 50).

result in weight loss (El-Naggar *et al.*, 2021; Assar *et al.*, 2022; Olugbodi *et al.*, 2023). Conversely, concomitant administration of different doses of *T. officinale* (500 and 250 mg/kg) with AgNPs results in effective management of weight loss compared to rats in the AgNPs group. This finding was aligned with an earlier study (Tan *et al.*, 2017) that suggested the increased growth performance of juvenile golden pompano *Trachinotus ovatus* after dietary *T. officinale* extracts were administered. Also, Adelakun *et al.* reported that treatment of diabetic rats with *T. officinale* extract raised the rats' body weight (Adelakun *et al.*, 2024). Concerning liver weight index, the current study showed no significant change in the liver weight index in AgNPs-intoxicated rats compared to the control group. These results agreed with the previous study by Rezaei *et al.* (2018). Also, this outcome was consistent with a previous study by Fahmy *et al.* (2020), which reported that rats administrated different oral doses of AgNPs (5, 25 and 50 mg/kg b.w.) five days a week for three months did not exhibit any appreciable changes in their absolute or relative liver weight. Furthermore, Assar *et al.* (2022) reported that adult rats treated with AgNPs for a period of 15 and 30 days showed a decrease in liver weight. This decrease in liver weight might be attributed to a rise in lipid peroxidation, leading to structural changes in lipid vacuoles. On the other hand, (*T. officinale* 500 mg + AgNPs) treated rats displayed a decrease in the liver weight index compared to the AgNPs group. These results agreed with the previous study by Mahesh *et al.* (2010).

The intracellular enzyme release is a significant marker of hepatocyte injury (Abdelkader *et al.*, 2020). Rats intoxicated with AgNPs exhibited significantly elevated ALT, AST, ALP, LDH, albumin and total protein levels, with a significant decline in total bilirubin levels in comparison

with control rats. Similarly, previous research suggested that the toxic effects of AgNPs on rat bodies can affect liver functions. This may be attributed to the AgNPs' free radicals that cause hepatocytes to release ALT and allow it to enter the blood (Abd El-Maksoud *et al.*, 2019; Salama *et al.*, 2023). Rats responded to external stimuli by phagocytosing AgNPs with a rise in the number of WBCs (Behzadi *et al.*, 2017). However, treatment with various dosages of *T. officinale* (500 and 250 mg/kg) showed a marked reduction in the liver function enzymes associated with marked elevation in the serum total bilirubin. A previous study demonstrated that *T. officinale* improved the liver function of rats subjected to CCL4 (Hamza *et al.*, 2020). Also, Devaraj discussed the hepatoprotective properties of dandelion (*T. officinale*), especially against various chronic liver diseases (Devaraj, 2016).

Furthermore, in this study, exposure of rats to AgNPs disrupts the antioxidant redox system, including GSH and SOD which aligned with prior research (Ansar *et al.*, 2017). The AgNPs intoxicated rats exhibited significantly raised MDA levels. The AgNP toxicity leads to oxidative stress that results in enhancing MDA production in hepatocytes (Fatemi *et al.*, 2017). These findings corroborate prior research (Moradi-Sardareh *et al.*, 2018; El-Naggar *et al.*, 2021). However, treatment with various doses of *T. officinale* (500 and 250 mg/kg) results in significantly reduced MDA levels linked to a marked increment in the GSH and SOD levels. These outcomes were consistent with earlier research that showed that dandelion (*T. officinale*) treatment before monosodium glutamate exposure led to a significant rise in the GSH levels and a significant decline in the MDA levels (Hussein and Jawad, 2023). Also, Hamza and his colleagues demonstrated the ameliorative impact of dandelion (*T. officinale*) on CCL4-triggered liver oxidative stress as it showed the best efficacy in normalizing the hepatic content of MDA and SOD activities (Hamza *et al.*, 2020).




Fig 7 (A-F): Photomicrographs of the immunohistochemical stain of the liver against anti-caspase-3 antibody.

All of these outcomes were verified by the histopathological investigation of the liver. The AgNPs-intoxicated group showed obvious histopathological changes; these include degenerations of hepatocytes leaving empty spaces, dilated hyperemic and congested blood sinusoids, degenerated hepatocytes and some of them appeared with eosinophilic cytoplasm and deeply stained pyknotic nuclei. The portal tract revealed congested and dilated portal veins, mononuclear cell infiltrations and necrotic empty spaces left by degenerated hepatocytes. Our outcomes are in accordance with prior research that demonstrated that exposure to AgNPs for 28 days led to congestion of central venules, abnormal morphology of hepatocytes and the blood sinusoids that appear infiltrated by inflammatory cells (Olugbodi *et al.*, 2023). Also, Hamza and his colleagues reported that AgNPs induced degenerative alterations in the liver, kidney and cardiac tissue of male rats (Hamza *et al.*, 2020). Treatment with (*T. officinale* 500 mg and 250 mg) demonstrated a liver structure comparable to that of the control group, with very minor histological alterations. Furthermore, the livers of rats administered (*T. officinale* 500 mg + AgNPs) exhibited significant protection of hepatic cells against necrotic cell death, in contrast to the extensive liver damage found in the AgNPs-treated group. Most of the hepatocytes appeared normal in histological structures. Administration of (*T. officinale* 250 mg+ AgNPs) showed alleviated inflammation, edema, leucocytic cell infiltration, necrotizing hepatocytes and connective tissue fibre propagation which is produced by AgNP intoxication. The histological architecture of the liver of rats treated with (*T. officinale* 500 mg+ AgNPs) group appeared more likely similar to the control group compared to that of (*T. officinale* 250 mg+ AgNPs) group. Our findings agree with the findings of earlier research observed that the hepatic injuries were significantly reduced by *T. officinale* root extract pretreatments (Pfingstgraf *et al.*, 2021). Microscopic examination of the hepatic tissue of rats co-treated with *T. officinale* demonstrated an enhancement in liver histoarchitecture, like that of the control rat. The impacts of *T. officinale* roots can be attributed to their high inulin content (Jalili *et al.*, 2020). Here, we demonstrated that the hepatoprotective abilities of *T. officinale* are more effective in inhibiting liver fibrosis and inflammation. This could be due to the variations in classes and contents of polyphenolic compounds. The roots of *T. officinale* include several phytochemicals, encompassing sesquiterpene lactones, polysaccharides, terpenoids, and phenolic compounds having antioxidant properties (Qadir *et al.*, 2022; Ürüsan, 2023).

Regarding the immunohistochemistry reactivity of liver tissue to the anti-caspase-3 antibody, the AgNPs intoxicated group displayed severe immunoreactivity confirming widespread apoptosis. These outcomes aligned with previous studies (Shehata *et al.*, 2022; Yousef *et al.*, 2022). Conversely, administration of *T. officinale* at various doses (500 and 250 mg/kg) has protective effects against apoptosis as confirmed in a previous study (Abdel-magied *et al.*, 2019).

CONCLUSION

This study demonstrated that *T. officinal* roots can effectively protect against AgNP-triggered hepatotoxicity in male rats through their antioxidant potential activity.

Informed consent

The experimental design was approved by the Animal Care and Use Committee at King Abdul-Aziz University, Faculty of Pharmacy, Jeddah, Saudi Arabia (Approval number: PH-1444-16).

Conflict of interest

The authors declare no conflict of interest.

REFERENCES

- Abd El-Maksoud, E.M., Lebda, M.A., Hashem, A.E., Taha, N.M. and Kamel, M.A. (2019). Ginkgo biloba mitigates silver nanoparticles- induced hepatotoxicity in Wistar rats via improvement of mitochondrial biogenesis and antioxidant status. *Environmental Science and Pollution Research*. 26: 25844-25854. <https://doi.org/10.1007/s11356-019-05835-2>.
- Abdel-Magied, N., Abdel Fattah, S.M. and Elkady, A.A. (2019). Differential effect of *Taraxacum officinale* L. (dandelion) root extract on hepatic and testicular tissues of rats exposed to ionizing radiation. *Molecular Biology Reports*. 46: 4893-4907. <https://doi.org/10.1007/s11033-019-04939-9>.
- Abdelkader, N.F., Elyamany, M., Gad, A.M., Assaf, N., Fawzy, H.M. and Elesawy, W.H. (2020). Ellagic acid attenuates liver toxicity induced by valproic acid in rats. *Journal of Pharmacological Sciences*. 143: 23-29. <https://doi.org/10.1016/j.jphs.2020.01.007>.
- Aboelwafa, H.R., Ramadan, R.A., Ibraheim, S.S. and Yousef, H.N. (2022). Modulation effects of eugenol on nephrotoxicity triggered by Silver nanoparticles in adult rats. *Biology*. 11: 1719.
- Adelakun, S.A., Akomaye, A.J., Omotoso, O.D. and Arowosegbe, O.A. (2024). Anti-hepatopathy and anti-nephropathy activities of *Taraxacum officinale* in a rat model of Streptozotocin diabetes-induced hepatorenal toxicity and dyslipidemia via attenuation of oxidative stress, inflammation, apoptosis, electrolyte imbalances and mitochondrial dysfunction. *Aspects of Molecular Medicine*. 3: 100034.
- Al-Doaiss, A., Jarrar, Q. and Moshawih, S. (2020). Hepatic histopathological and ultrastructural alterations induced by 10 nm silver nanoparticles. *IET nanobiotechnology*. 14: 405-411. <https://doi.org/10.1049/iet-nbt.2020.0039>.
- Albrahim, T. and Alonazi, M.A. (2020). Role of Beetroot (*Beta vulgaris*) Juice on Chronic Nanotoxicity of Silver Nanoparticle-Induced Hepatotoxicity in Male Rats. *International Journal of Nanomedicine*. 15: 3471-3482. <https://doi.org/10.2147/ijn.S248078>.
- Amin, A., Hamza, A.A., Daoud, S., Khazanehdari, K., Hrou, A.A., Baig, B., Chaiboonchoe, A., Adrian, T.E., Zaki, N. and Salehi-Ashtiani, K. (2016). Saffron-based crocin prevents early lesions of liver cancer: *In vivo*, *in vitro* and network analyses. *Recent Patents on Anti-Cancer Drug Discovery*. 11: 121-133. <https://doi.org/10.2174/1574892810666151102110248>.
- Ansar, S., Alshehri, S.M., Abudawood, M., Hamed, S.S. and Ahamad, T. (2017). Antioxidant and hepatoprotective role of selenium against silver nanoparticles. *International Journal of Nanomedicine*. 12: 7789-7797. <https://doi.org/10.2147/ijn.S136748>.

- Aremu, O.O., Oyediji, A.O., Oyediji, O.O., Nkeh-Chungag, B.N. and Rusike, C.R.S. (2019). *in vitro* and *in vivo* antioxidant properties of taraxacum officinale in N- ω -Nitro-L-Arginine Methyl ester (L-NAME)-induced hypertensive rats. *Antioxidants*. 8: 309.
- Ashktorab, H., Soleimani, A., Singh, G., Amin, A., Tabatabaei, S., Latella, G., Stein, U., Akhondzadeh, S., Solanki, N., Gondré-Lewis, M.C., Habtezion, A. and Brim, H. (2019). Saffron: The golden spice with therapeutic properties on digestive diseases. *Nutrients*. 11: 943. <https://doi.org/10.3390/nu11050943>.
- Assar, D.H., Mokhbatly, A.A., Ghazy, E.W., Elbially, Z.I., Gaber, A.A., Hassan, A.A., Nabil, A. and Asa, S.A. (2022). Silver nanoparticles induced hepatotoxicity via the apoptotic/antiapoptotic pathway with activation of TGF β -1 and α -SMA triggered liver fibrosis in Sprague Dawley rats. *Environmental Science and Pollution Research*. 29: 80448-80465. <https://doi.org/10.1007/s11356-022-21388-3>.
- Ayaz, N.O., Ramadan, K.S., Farid, H.E. and Alnahdi, H.S. (2017). Protective role and antioxidant activity of arabic gum against trichloroacetate- induced toxicity in liver of male rats. *Indian Journal of Animal Research*. 51(2): 303-309. doi: 10.18805/ijar.10976.
- Bancroft, J.D. and Gamble, M. (Eds.). (2008). *Theory and Practice of Histological Techniques*. Elsevier Health Sciences.
- Behzadi, S., Serpooshan, V., Tao, W., Hamaly, M.A., Alkawarek, M.Y., Dreaden, E.C., Brown, D., Alkilany, A. M., Farokhzad, O.C. and Mahmoudi, M. (2017). Cellular uptake of nanoparticles: Journey inside the cell. *Chemical Society Reviews*. 46: 4218-4244. <https://doi.org/10.1039/c6cs00636a>.
- Blanco, J., Tomas-Hernandez, S., García, T., Mulero, M., Gomez, M., Domingo, J.L. and Sánchez, D.J. (2018). Oral exposure to silver nanoparticles increases oxidative stress markers in the liver of male rats and deregulates the insulin signalling pathway and p53 and cleaved caspase 3 protein expression. *Food and Chemical Toxicology*. 115: 398-404. <https://doi.org/10.1016/j.fct.2018.03.039>.
- Cai, L., Wan, D., Yi, F. and Luan, L. (2017). Purification, preliminary characterization and hepatoprotective effects of polysaccharides from dandelion root. *Molecules*. 22: 1409. <https://doi.org/10.3390/molecules22091409>.
- Choi, U.K., Lee, O.H., Yim, J.H., Cho, C.W., Rhee, Y.K., Lim, S.I. and Kim, Y.C. (2010). Hypolipidemic and antioxidant effects of dandelion (*Taraxacum officinale*) root and leaf on cholesterol-fed rabbits. *International journal of molecular sciences*. 11: 67-78.
- Devaraj, E. (2016). Hepatoprotective properties of dandelion: Recent update. *Journal of Applied Pharmaceutical Science*. 6: 202-205.
- Di Napoli, A. and Zucchetti, P. (2021). A comprehensive review of the benefits of *Taraxacum officinale* on human health. *Bulletin of the National Research Centre*. 45: 110.
- Domitrović, R., Jakovac, H., Romić, Ž., Rahelić, D. and Tadić, Ž. (2010). Antifibrotic activity of *Taraxacum officinale* root in carbon tetrachloride-induced liver damage in mice. *Journal of ethnopharmacology*. 130: 569-577.
- El-Naggar, S., Khandil, R. and Salem, M. (2021). Phoenix dactylifera seed extract ameliorates the biochemical toxicity induced by silver nanoparticles in mice. *International Journal of Cancer and Biomedical Research*. 5: 73-84.
- Ema, M., Okuda, H., Gamo, M. and Honda, K. (2017). A review of reproductive and developmental toxicity of silver nanoparticles in laboratory animals. *Reproductive Toxicology*. 67: 149-164. <https://doi.org/10.1016/j.reprotox.2017.01.005>.
- Fahmy, E., Sharkawy, A., Ibraheim, T.A. and Yahia, D. (2020). Effects of silver nitrate nanoparticles on the oxidative status in sprague dawley rats. *Assiut Veterinary Medical Journal*. 66: 85-91.
- Fallah, H.H., Zarrei, M.A., Ziai, S.A., Mehrazma, M., Alavian, S.M., Kianbakht, S. and Mehdizadeh, M. (2010). The effects of *Taraxacum officinale* L. and *Berberis vulgaris* L. root extracts on carbon tetrachloride induced liver toxicity in rats. *Journal of Medicinal Plants*. 9: 45-52.
- Fatemi, M., Moshtaghian, J., Ghaedi, K., Jafari Dinani, N. and Naderi, G. (2017). Effects of silver nanoparticle on the developing liver of rat pups after maternal exposure. *Iranian Journal of Pharmaceutical Research*. 16: 685-693.
- Flores-López, L.Z., Espinoza-Gómez, H. and Somanathan, R. (2019). Silver nanoparticles: Electron transfer, reactive oxygen species, oxidative stress, beneficial and toxicological effects. Mini review. *Journal of Applied Toxicology*. 39: 16-26. <https://doi.org/10.1002/jat.3654>.
- Ghonimi, W.A.M., Alferah, M.A.Z., Dahran, N. and El-Shetry, E.S. (2022). Hepatic and renal toxicity following the injection of copper oxide nanoparticles (CuO NPs) in mature male Westar rats: Histochemical and caspase 3 immunohistochemical reactivities. *Environmental Science and Pollution Research*. 29: 81923-81937. <https://doi.org/10.1007/s11356-022-21521-2>.
- Hamza, A.A., Heeba, G.H., Elwy, H.M., Murali, C., El-Awady, R. and Amin, A. (2018). Molecular characterization of the grape seeds extract's effect against chemically induced liver cancer: *in vivo* and *in vitro* analyses. *Scientific Reports*. 8: 1270. <https://doi.org/10.1038/s41598-018-19492-x>.
- Hamza, A.A., Mohamed, M.G., Lashin, F.M. and Amin, A. (2020). Dandelion prevents liver fibrosis, inflammatory response and oxidative stress in rats. *The Journal of Basic and Applied Zoology*. 81: 1-13.
- Hussein, I. and Jawad, R.A. (2023). Study of the dandelion on some physiological parameters against induced by monosodium glutamate in male white rabbits (*Lepus arcticus*). In *AIP Conference Proceedings* (Vol. 2414, No. 1). AIP Publishing.
- Ignat, M.V., Coldea, T.E., Salany, L.C. and Mudura, E. (2021). Plants of the spontaneous flora with beneficial action in the management of diabetes, hepatic disorders and cardiovascular disease. *Plants (Basel)*. 10: 216. <https://doi.org/10.3390/plants10020216>.
- Ivanov, I., Petkova, N., Tumbarski, J., Dincheva, I., Badjakov, I., Denev, P. and Pavlov, A. (2018). GC-MS characterization of n-hexane soluble fraction from dandelion (*Taraxacum officinale* Weber ex FH Wigg.) aerial parts and its antioxidant and antimicrobial properties. *Zeitschrift für Naturforschung C*. 73: 41-47.
- Jalili, C., Taghadosi, M., Pazhouhi, M., Bahrehmand, F., Miraghaee, S., Pourmand, D. and Rashidi, I. (2020). An overview of therapeutic potentials of *Taraxacum officinale* (dandelion): A traditionally valuable herb with a reach historical background. *WCRJ World Cancer Res. J.* 7: e1679.

- Khajuria, P., Raghuwanshi, P., Rastogi, A., Koul, A.L., Zargar, R. and Kour, S. (2018). Hepatoprotective effect of seabuckthorn leaf extract in streptozotocin induced diabetes mellitus in Wistar rats. *Indian Journal of Animal Research*. 52(12): 1745-1750. doi: 10.18805/ijar.B-3439.
- Lee, S.H. and Jun, B.H. (2019). Silver nanoparticles: Synthesis and application for nanomedicine. *International Journal of Molecular Sciences*. 20: 865. <https://doi.org/10.3390/ijms20040865>.
- Lekha, D.C., Shanmugam, R., Madhuri, K., Dwarampudi, L.P., Bhaskaran, M., Kongara, D., Tesfaye, J.L., Nagaprasad, N., Bhargavi, V.N. and Krishnaraj, R. (2021). Review on silver nanoparticle synthesis method, antibacterial activity, drug delivery vehicles and toxicity pathways: Recent advances and future aspects. *Journal of Nanomaterials*. 2021: 4401829.
- Li, Q., Li, H. J., Xu, T., Du, H., Huan Gang, C.L., Fan, G. and Zhang, Y. (2018). Natural medicines used in the traditional tibetan medical system for the treatment of liver diseases. *Front Pharmacol*. 9: 29. <https://doi.org/10.3389/fphar.2018.00029>.
- Mahesh, A., Jeyachandran, R., Cindrella, L., Thangadurai, D., Veerapur, V. and Muralidhara Rao, D. (2010). Hepatocurative potential of sesquiterpene lactones of *Taraxacum officinale* on carbon tetrachloride induced liver toxicity in mice. *Acta Biologica Hungarica*. 61: 175-190.
- Mao, B.H., Chen, Z.Y., Wang, Y.J. and Yan, S.J. (2018). Silver nanoparticles have lethal and sublethal adverse effects on development and longevity by inducing ROS-mediated stress responses. *Scientific Reports*. 8: 2445. <https://doi.org/10.1038/s41598-018-20728-z>.
- Milek, M., Marciněáková, D. and Legáth, J. (2019). Polyphenols content, antioxidant activity and cytotoxicity assessment of *Taraxacum officinale* extracts prepared through the micelle-mediated extraction method. *Molecules*. 24: 1025.
- Mishra, A.R., Zheng, J., Tang, X. and Goering, P.L. (2016). Silver nanoparticle-induced autophagic-lysosomal disruption and NLRP3-Inflammasome Activation in HepG2 Cells Is Size-dependent. *Toxicological Sciences*. 150: 473-487. <https://doi.org/10.1093/toxsci/kfw011>.
- Monteiro, D.R., Gorup, L.F., Silva, S., Negri, M., de Camargo, E.R., Oliveira, R., Barbosa, D.B. and Henriques, M. (2011). Silver colloidal nanoparticles: Antifungal effect against adhered cells and biofilms of *Candida albicans* and *Candida glabrata*. *Biofouling*. 27: 711-719. <https://doi.org/10.1080/08927014.2011.599101>.
- Moradi-Sardareh, H., Basir, H.R.G., Hassan, Z.M., Davoudi, M., Amidi, F. and Paknejad, M. (2018). Toxicity of silver nanoparticles on different tissues of Balb/C mice. *Life sciences*. 211: 81-90. <https://doi.org/10.1016/j.lfs.2018.09.001>.
- Nguyen, C., Mehaidli, A., Baskaran, K., Grewal, S., Pupulin, A., Ruvinov, I., Scaria, B., Parashar, K., Vegh, C and Pandey, S. (2019). Dandelion root and lemongrass extracts induce apoptosis, enhance chemotherapeutic efficacy and reduce tumour xenograft growth *in vivo* in prostate cancer. Evidence Based Complementary and Alternative Medicine. 2019: 2951428.
- Olugbodi, J.O., Lawal, B., Bako, G., Onikanni, A.S., Abolenin, S.M., Mohammud, S.S., Ataya, F.S. and Batiha, G. E. (2023). Effect of sub-dermal exposure of silver nanoparticles on hepatic, renal and cardiac functions accompanying oxidative damage in male Wistar rats. *Scientific Reports*. 13: 10539.
- Ovadge, P., Chatterjee, S., Griffin, C., Tran, C., Hamm, C. and Pandey, S. (2011). Selective induction of apoptosis through activation of caspase-8 in human leukemia cells (Jurkat) by dandelion root extract. *Journal of Ethnopharmacology*. 133: 86-91.
- Ovadge, P., Chochkeh, M., Akbari-Asl, P., Hamm, C. and Pandey, S. (2012). Selective induction of apoptosis and autophagy through treatment with dandelion root extract in human pancreatic cancer cells. *Pancreas*. 41: 1039-1047.
- Ovadge, P., Ammar, S., Guerrero, J.A., Arnason, J.T. and Pandey, S. (2016). Dandelion root extract affects colorectal cancer proliferation and survival through the activation of multiple death signalling pathways. *Oncotarget*. 7: 73080.
- Park, C.M., Cha, Y.S., Youn, H.J., Cho, C.W. and Song, Y.S. (2010). Amelioration of oxidative stress by dandelion extract through CYP2E1 suppression against acute liver injury induced by carbon tetrachloride in sprague dawley rats. *Phytotherapy Research*. 24: 1347-1353.
- Park, C.M., Park, J.Y., Noh, K.H., Shin, J.H. and Song, Y.S. (2011). *Taraxacum officinale* Weber extracts inhibit LPS-induced oxidative stress and nitric oxide production via the NF- κ B modulation in RAW 264.7 cells. *Journal of ethnopharmacology*. 133: 834-842.
- Pereira, C., Barros, L. and Ferreira, I.C. (2016). Extraction, identification, fractionation and isolation of phenolic compounds in plants with hepatoprotective effects. *Journal of the Science of Food and Agriculture*. 96: 1068-1084.
- Pfingstgraf, I.O., Taulescu, M., Pop, R.M., Orăsan, R., Vlase, L., Uifalean, A., Todea, D., Alexescu, T., Toma, C. and Pîrvu, A.E. (2021). Protective effects of *Taraxacum officinale* L. (Dandelion) root extract in experimental acute on chronic liver failure. *Antioxidants (Basel)*. 10: 504. <https://doi.org/10.3390/antiox10040504>.
- Qadir, I., Nazir, S., Sheikh, M.A., Naaz, F., Bashir, S., Ovais, S., Khan, N.A. and Masoodi, M. H. (2022). *Taraxacum officinale*: The esculent dandelion as herbal medicine. In *Edible plants in health and diseases: Volume II: Phytochemical and Pharmacological Properties* Springer. (pp. 299-326).
- Quintero-Quiroz, C., Acevedo, N., Zapata-Giraldo, J., Botero, L.E., Quintero, J., Zárate-Triviño, D., Saldarriaga, J. and Pérez, V.Z. (2019). Optimization of silver nanoparticle synthesis by chemical reduction and evaluation of its antimicrobial and toxic activity. *Biomater Res*. 23: 27. <https://doi.org/10.1186/s40824-019-0173-y>.
- Rβcz-Kotilla, E., Rac, G. and Solomon, A. (1974). The action of *Taraxacum officinale* extracts on the body weight and diuresis of laboratory animals. *Planta Medica*. 26: 212-217.
- Rezaei, A., Farzinpour, A., Vaziry, A. and Jalili, A. (2018). Effects of silver nanoparticles on hematological parameters and hepatorenal functions in laying japanese Quails. *Biological Trace Element Research*. 185: 475-485.
- Salama, B., Alzahrani, K.J., Alghamdi, K.S., Al-Amer, O., Hassan, K.E., Elhefny, M.A. et al. (2023). Silver nanoparticles enhance oxidative stress, Inflammation and apoptosis in liver and kidney Tissues: Potential protective role of thymoquinone. *Biological Trace Element Research*. 201: 2942-2954. <https://doi.org/10.1007/s12011-022-03399-w>.

- Saratale, R.G., Benelli, G., Kumar, G., Kim, D.S. and Saratale, G.D. (2018). Bio-fabrication of silver nanoparticles using the leaf extract of an ancient herbal medicine, dandelion (*Taraxacum officinale*), evaluation of their antioxidant, anticancer potential and antimicrobial activity against phytopathogens. *Environmental Science and Pollution Research*. 25: 10392-10406.
- Schutz, K., Carle, R. and Schieber, A. (2006). *Taraxacum*-a review on its phytochemical and pharmacological profile. *Journal of Ethnopharmacology*. 107: 313-323.
- Şenocak, E.A. and Yıldırım, B.A. (2017). Ratlarda parasetamol ile oluşturulan hepatotoksiste üzerine *Taraxacum officinale* etanol ekstraktının etkisi. *Harran Üniversitesi Veteriner Fakültesi Dergisi*. 6: 11-18.
- Shaaban, H.M., Mehana, E.S.E., Oda, S.S., Tohamy, H.G. and El-Karim, D.R.G. (2023). Clinico-pathological evaluation of hepatoprotective effect of dandelion and Ginkgo biloba extract against thioacetamide induced liver damage. *Alexandria Journal of Veterinary Sciences*. 76: 90-94.
- Shehata, A.M., Salem, F.M., El-Saied, E.M., Abd El-Rahman, S.S., Mahmoud, M.Y. and Noshay, P.A. (2022). Evaluation of the ameliorative effect of zinc nanoparticles against silver nanoparticle-induced toxicity in liver and kidney of rats. *Biological Trace Element Research*. 1-11.
- Syafiuddin, A., Salmiati, Salim, M.R., Beng Hong Kueh, A., Hadibarata, T. and Nur, H. (2017). A review of silver nanoparticles: Research trends, global consumption, synthesis, properties and future challenges. *Journal of the Chinese Chemical Society*. 64: 732-756.
- Tan, X., Sun, Z., Chen, S., Chen, S., Huang, Z., Zhou, C., Zou, C., Liu, Q., Ye, H., Lin, H., Ye, C. and Wang, A. (2017). Effects of dietary dandelion extracts on growth performance, body composition, plasma biochemical parameters, immune responses and disease resistance of juvenile golden pompano *Trachinotus ovatus*. *Fish and Shellfish Immunology*. 66: 198-206. <https://doi.org/10.1016/j.fsi.2017.05.028>.
- Ürüşan, H. (2023). Effects of dandelion (*Taraxacum officinale*) supplementation on productive performance, egg quality traits, serum biochemical parameters and liver fat rate in laying hens. *Indian Journal of Animal Research*. 57(8): 1018-1024. doi: 10.18805/IJAR.BF-1662.
- Xiong, P., Huang, X., Ye, N., Lu, Q., Zhang, G., Peng, S., Wang, H. and Liu, Y. (2022). Cytotoxicity of metal-based nanoparticles: From mechanisms and methods of evaluation to pathological manifestations. *Advanced Science*. 9: 2106049. <https://doi.org/10.1002/advs.202106049>.
- Yousef, H.N., Ibraheim, S.S., Ramadan, R.A. and Aboelwafa, H.R. (2022). The ameliorative role of eugenol against silver nanoparticles-induced hepatotoxicity in male wistar rats. *Oxidative Medicine and Cellular Longevity*. 2022: 3820848. <https://doi.org/10.1155/2022/3820848>.
- Zainal, Z., Ong, A., Yuen May, C., Chang, S.K., Abdul Rahim, A. and Khaza'ai, H. (2020). Acute and subchronic oral toxicity of oil palm puree in sprague-dawley rats. *Int J. Environ Res Public Health*. 17. <https://doi.org/10.3390/ijerph17103404>.
- Zhang, X.F., Liu, Z.G., Shen, W. and Gurunathan, S. (2016). Silver nanoparticles: Synthesis, characterization, properties, applications and therapeutic approaches. *International Journal of Molecular Sciences*. 17: 1534. <https://doi.org/10.3390/ijms17091534>.