



Effects of Different Doses of Dia-VTM MB_{PRO} on Growth Performance, Serum Metabolites and Antioxidant Profiles of Broiler Chickens

Y. Alyousef¹, M. Shawky², G. Rayan¹, H. Darrag³, H. Najib¹, A. Mohammed¹

10.18805/IJAR.BF-1984

ABSTRACT

Background: *Saccharomyces cerevisiae* and/or essential oils have become increasingly important in poultry production due to improved gut health, enhanced growth performance and boosted immune system.

Methods: A total of 160 one-day old Ross-308 broiler chicks were randomly distributed into four groups, each consisting of four replicates with ten birds per replicate. The chicks were fed a basal diet containing *Saccharomyces cerevisiae* and essential oils at concentrations of 0 MB_{PRO}, 200 MB_{PRO}, 400 MB_{PRO} and 800 MB_{PRO} mg/kg feed respectively. Body weight gain (g), feed intake and feed conversion ratio were recorded. In addition, carcass traits were determined through dressing percentage and carcass parts. Furthermore, blood samples were collected for determination of serum metabolites and antioxidant profiles.

Result: The results showed that dietary *Saccharomyces cerevisiae* and essential oils incorporation improved ($P < 0.001$) body weight gain and feed conversion ratio. Carcass traits remained unaffected by the combination treatments if compared to control. Serum metabolites and antioxidant profiles were improved with the combination treatments. It could be concluded that *Saccharomyces cerevisiae* and essential oils incorporation at 400 mg/kg feed proved to be an effective growth promoter, enhancing growth performance, metabolites and antioxidant profiles in broiler chicks.

Key words: Antioxidants, Carcass, Chicks, Feed conversion, Growth, Metabolites, Serum.

INTRODUCTION

The global population is expected to reach 9.7 billion by 2050 and 10.4 billion by 2100 (Mohammed and Alshaibani 2025; Mohammed *et al.*, 2025). The escalating global population presents a profound challenge to food security, demanding a significant increase in food production and distribution (Abdullahi *et al.*, 2024). This surge necessitates not only expanded food production but also more efficient and sustainable practices to mitigate environmental impacts (Gamage *et al.*, 2024). Furthermore, shifts in dietary preferences, particularly the increased consumption of animal products linked to increase incomes, further strain on resources (Vermeulen *et al.*, 2020). Therefore, ensuring adequate food availability for a growing population requires innovative solutions, including technological advancements in agriculture, reductions in food waste and equitable distribution systems (Pawlak *et al.*, 2020).

Poultry production is essential for providing protein-rich food to the world's increasing urban populations and is a rapidly expanding area of animal agriculture. Therefore, the incorporation of dietary probiotics in broiler chicken diets has gained significant attention due to its potential impact to enhance poultry production and health (Krysiak *et al.*, 2021; Dong *et al.*, 2024; Ayana and Kamutambuko 2024). Several positive impacts have been confirmed in several studies focused on the inclusion of dietary probiotics in broiler chicken including improvement of feed conversion and nutrient utilization, body weight gain and gut health (Abd El-Hack *et al.*, 2020; Urban *et al.*, 2024; Naeem and Bourassa, 2025).

¹Department of Animal and Fish Production, College of Agriculture and Food Sciences, King Faisal University, P.O. Box 402, Al-Ahsa 31982, KSA.

²Avian Research Center, King Faisal University, P.O. Box 402, Al-Ahsa 31982, KSA.

³Research and Training Station, King Faisal University, P.O. Box 402, Al-Ahsa 31982, KSA.

Corresponding Author: A. Mohammed, Department of Animal and Fish Production, College of Agriculture and Food Sciences, King Faisal University, P.O. Box 402, Al-Ahsa 31982, KSA. Email: aamohammed@kfu.edu.sa

How to cite this article: Alyousef, Y., Shawky, M., Rayan, G., Darrag, H., Najib, H. and Mohammed, A. (2025). Effects of Different Doses of Dia-VTM MB_{PRO} on Growth Performance, Serum Metabolites and Antioxidant Profiles of Broiler Chickens. Indian Journal of Animal Research. 1-7. doi: 10.18805/IJAR.BF-1984.

Submitted: 04-03-2025 **Accepted:** 23-04-2025 **Online:** 03-06-2025

The dietary incorporation *Saccharomyces cerevisiae* and/or essential oils into broiler chicken nutrition has gained significant attention due to their potential to improve poultry health and production through several mechanisms (Heinsohn *et al.*, 2024; Movahedi *et al.*, 2024). Yeast culture contains yeast biomass and fermentation metabolites (Shurson, 2018). Yeast culture nutritional and health care functions to stimulate animal growth are attributed to its fermentation metabolites, including micronutrients and/or pro-vitamins (Brake 1991; Callaway and Martin 1997; Sun

et al., 2020). In addition, essential oils due to their growth-promoting and immunity-boosting properties and antimicrobial and antioxidant activities have gained much interest in the poultry industry (Valdivieso-Ugarte *et al.*, 2019; Movahedi *et al.*, 2024). Essential oil compounds can improve enzymatic activity in the gastrointestinal tract of chickens resulting in better digestion and absorption of nutrients (Su *et al.*, 2021). Consequently, while dietary *S. cerevisiae* and/or essential oils might offer a valuable tool for enhancing growth performance, careful consideration of these combinations is essential for optimal results. Therefore, the aim of the current study was to investigate the impacts of the dietary Dia-V™ MB_{PRO} supplementation (0, 200, 400, or 800 mg/kg feed) on growth performances, carcass traits, serum antioxidant and biochemical profiles of growing broilers.

MATERIALS AND METHODS

The experimental procedures were approved by the ethical committee concerning experimental animal care guidelines of King Faisal University [KFU-REC-2025-MAR-EA252670]. The procedure of the experiment was executed in the experimental animal research station of King Faisal University.

Site of study and experimental design

The current study was carried out from November 2024 to January 2025. A total of one hundred sixty one-day old Ross308 broiler chicks were randomly classified over four groups, each consisting of four replicates/ten birds (Fig 1). The basal diet was formulated following the strain recommendations by Aviagen (2018). These birds were fed a basal diet supplemented with 0, 200, 400, or 800 mg/kg feed of *Saccharomyces cerevisiae* and essential oil compounds, which were commercially obtained (Dia-V™

MB_{PRO}). The Dia-V™ MB_{PRO} additives were directly mixed into the feed during diet preparation to ensure even distribution. The birds were bred in floor pens measuring 16 square (each single pen) feet. The chicks were kept at 33.0°C for the first three days of age, after which the temperature was gradually reduced to 31.0°C for next 48.0 h. Subsequently, the birds were kept at an average of 31.0°C natural ambient temperature and 60.0% relative humidity during the study. The lighting program was maintained at 23 h light to 1h dark for the first 7 days, followed by 18 h light to 6 dark for the remaining of the experiment. The chicks had *ad libitum* access to feed and water.

Growth and carcass traits

At 7, 14, 21, 28 and 35 days of age, body weight gain (Kg) and feed intake (Kg) were recorded on a pen basis and feed conversion ratios were calculated (Abdel-Moneim *et al.*, 2025). On 35 days of age, eight chicks per group representing the groups' average weight were randomly selected for carcass evaluation to ensure unbiased sampling. The selected chicks were fasted for 12:0 h before weighing followed by slaughtering, de-feathering and eviscerating. The eviscerated carcasses and giblets were weighed for all groups. The eviscerated carcasses and giblets were recorded using a digital balance to ensure accuracy and expressed as a percentage of live body weight (Abdel-Moneim *et al.*, 2025).

Blood sample collection and analyses

Blood samples were collected from the brachial vein of twelve birds of each group at the end of experiment. The obtained blood samples were centrifuged to obtain serum for biochemistry and antioxidant analyses (Analyzer Skyla VB1). The analyzed biochemistry parameters include total proteins,




Fig 1: Experimental design of study for Dia-V™ MB_{PRO} additives to chicks.

liver enzymes, urea, creatinine and mineral values. The antioxidant enzymatic activities of serum malondialdehyde (MDA) was measured using a colorimetric method that involves its interaction with 2-thiobarbituric acid (2-TBA) at a wavelength of 532 nm (Mihara and Uchiyama, 1978). The ferric reducing ability (FRAP) test was conducted as described by Benzie and Strain with modification (Benzie and Strain, 1996). The ABTS radical cation decolorization test was conducted as the recommendation of Re *et al.* (1999). The free radical scavenging activity (FRSA) was assessed as outlined by Blois with modification (Blois, 1958). Total thiol (sulfhydryl group, -SH) concentrations (TTL) were quantified as described by Hu with modification (Hu, 1994).

Statistical analysis

Body weight gain (g), feed conversion (kg/kg), dressing and giblets percentage (%), serum antioxidant and metabolites values of control and Dia-VTM MB_{PRO} treated groups were statistically analyzed using SAS program (SAS 2008) through General Linear Model procedure according to the model:

$$Y_{ij} = \mu + T_i + E_{ij}$$

Where,

μ = Mean.

T_i = Effects of Dia-VTM MB_{PRO} levels (200MB_{PRO}, 400 MB_{PRO} and 800 MB_{PRO}).

E_{ij} = Standard error.

To compare the significance among means of control and Dia-VTM MB_{PRO} treated groups, Duncan's multiple range test (1955) was used.

RESULTS AND DISCUSSION

The effects of Dia-VTM MB_{PRO} dietary inclusions (200 MB_{PRO}, 400 MB_{PRO} and 800 MB_{PRO}) to Ross-308 broiler chicks ration on feed intake, live body weight, feed conversion, carcass traits, serum metabolites and antioxidant capacity are presented in (Fig 2, 3 and 4) and (Table 1).

Growth performances

The effects of dietary inclusions (200 MB_{PRO}, 400 MB_{PRO} and 800 MB_{PRO}) on growth performance are presented in (Fig 2, 3 and 4). The dietary inclusions (200 MB_{PRO}, 400 MB_{PRO} and 800 MB_{PRO}) resulted in significantly higher final body weight (g) compared to the control group (0 MB_{PRO}) ($P < 0.03$). The Dia-VTM MB_{PRO} dietary inclusions resulted in significant growth performance represented by improved body weight gain, feed intake, feed conversion ratio, carcass traits (Fig 2, 3 and 4). In general, dietary inclusions resulted in significant improvement in growth performance parameters concerning live body weight, feed conversion and carcass traits. In addition, 400 MB_{PRO} dietary inclusions gave the highest values in body weight and dressing percentage compared to other groups. While, 200 mg/kg had better feed conversion ratio compared to all other groups.

Serum metabolites and antioxidant capacity

The effects of different levels of Dia-VTM MB_{PRO} in broiler diet on serum metabolites and antioxidant capacity to broiler

chickens are presented in (Table 1). The Dia-VTM MB_{PRO} dietary inclusions resulted in significantly higher total protein levels compared to the control group ($P < 0.03$). In 200 MB_{PRO} group, albumin value showed a significantly higher value compared to the control and other Dia-VTM MB_{PRO} groups ($P < 0.04$). There was no statistically significant differences in creatinine level among the control and Dia-VTM MB_{PRO} groups ($P = 0.47$). Groups 400 MB_{PRO} and 800 MB_{PRO} exhibited significantly higher glucose levels compared to the control and 200 MB_{PRO} groups ($P = 0.05$). The Dia-VTM MB_{PRO} inclusions significantly reduced total bilirubin levels compared to 0 MB_{PRO} inclusion ($P < 0.0001$). Regarding to AST, ALT and GGT enzymes, there were no significant differences in AST ($P = 0.54$), ALT ($P = 0.89$) and GGT (0.74) levels among the control and Dia-VTM MB_{PRO} groups. Values of ALP were significantly ($P = 0.01$) decreased in 400 MB_{PRO} and 800 MB_{PRO} groups compared to 200 MB_{PRO} and control ones. Regarding the antioxidant capacity, the FRAP, FRSA, ABTS and TTL values were not significantly different among groups whereas MDA values were significantly ($P = 0.001$) decreased in Dia-VTM MB_{PRO} groups compared to control one.

The results of the current study indicated the highest significant improvement of Dia-VTM MB_{PRO} dietary inclusions to Ross-308 broiler chicks on growth performance, serum metabolites and antioxidant capacity at level of 400 mg/kg. The effect of Dia-VTM MB_{PRO} could be attributed to several factors including improved gut health, boosted immune system and reduction of pathogens (Mazziotta *et al.*, 2023; Chandrasekaran *et al.*, 2024).

Growth performances

The effects of *Saccharomyces cerevisiae* (*S. cerevisiae*) and essential oils dietary inclusions to broiler chicks on growth performances have been explored in several studies (Lin *et al.*, 2023; Okasha *et al.*, 2023). Our results indicated that dietary 400 mg/kg MB_{PRO} inclusion gave the highest body weight compared to all groups. Higher feed conversion ratio was recorded for chicks fed diet containing 200 mg/kg compared to other groups. The negative or positive changes of growth performances over *S. cerevisiae* supplementation to broiler chicks were dependent on the given doses (Haldar *et al.*, 2011; Swamy and Upendra 2013). Okasha *et al.*, (2023) found that 0.25, 0.5 and 0.75% *S. cerevisiae* supplementation decreased body weight gain. On the other, significant increase of broilers BW fed diet added containing 0.1, 0.5, 0.75 and 1.0% *S. cerevisiae* compared to control one (Swamy and Upendra 2013; Okasha *et al.*, 2023).

Additionally, essential oils are characterized by biological useful properties in the diet of broiler chickens reflected in improved growth performance and increase the body's immune response (Ezzat *et al.*, 2016; Giannenas *et al.*, 2018; Puvaca *et al.*, 2015, 2022).

Collectively, dietary *Saccharomyces cerevisiae* and essential oils inclusion might provide comprehensive

benefits to poultry of all ages and types, fostering a healthy gut environment that leads to improved digestion and enhanced growth performance (Lin *et al.*, 2023; Movahedi *et al.*, 2024; Abd El-Ghany 2024). They achieve this by balancing gut bacteria, strengthening the intestinal lining

and stimulating the immune system while reducing harmful inflammation. The *S. cerevisiae* provides valuable proteins, vitamin B-complex and other beneficial factors that have been reported to increase phosphorus availability and utilization in animals (Moore *et al.*, 1994; Medina *et al.*, 2002).

Table 1: Effects of dietary *Saccharomyces cerevisiae* and essential oils (Dia-V™ MB_{PRO}) on plasma biochemistry profiles of broiler chickens at 35 d of age.

Parameters	Control	Treat 1	Treat 2	Treat 3	SEM	P value
Biochemical parameters						
Total protein, g/dl	3.95 ^b	4.12 ^a	4.40 ^a	4.23 ^a	0.075	0.03
Albumin, g/dl	2.13 ^b	2.22 ^b	2.41 ^a	2.24 ^b	0.055	0.04
Creatinine, mg/dl	0.77	0.86	0.83	0.085	0.04	0.47
Glucose, mg/dl	150.25 ^b	148.0 ^b	162.0 ^a	164.25 ^a	2.50	0.05
Total bilirubin, µmol/l	13.20 ^a	11.22 ^b	11.16 ^b	11.60 ^b	0.66 ^b	<0.0001
Aspartate aminotransferase, U/l	66.46	64.60	63.70	63.10	0.45	0.54
Alanine aminotransferase, U/l	58.25	56.33	55.28	54.95	0.45	0.89
γ-glutamyl transferase, U/l	43.0	44.00	45.30	42.0	0.92	0.74
Alkaline phosphatase, U/l	11.86 ^a	11.67 ^a	11.44 ^b	11.41 ^b	0.05	0.01
Antioxidant capacity						
FRAP, µM	820.6	842.0	853.0	886.3	7.22	0.67
FRSA, Inh. %	86.31	86.04	85.37	81.22	0.62	0.33
ABTS, µM	1.710	1.710	1.716	1.716	0.002	0.97
TTL, µM	143.12	138.24	137.64	130.92	1.313	0.05
MDA, mMol/dl	1.625 ^a	1.423 ^b	1.40 ^b	1.236 ^b	0.040	0.001

^{a, b}, Values between groups at the same row with different superscripts significantly differ at P<0.05.

SEM, Standard error.

FRAP, Ferric reducing antioxidant power.

FRSA, Free radical scavenging activity.

ABTS, 2,2'-Azino-bis(3-ethylbenzothiazoline-6-sulfonic acid).

TTL, Total thiol levels.

MDA, Malondialdehyde.




Fig 2: Effect of dietary Dia-V™ MB_{PRO} on final body weight of broiler chickens at 35 d of age.

Furthermore, it has been indicated in several studies that dietary *S. cerevisiae* inclusion reduced disease infections and improved feed efficiency (Line *et al.*, 1997; Santin *et al.*, 2001).

Serum metabolites and antioxidant capacity

The values of serum metabolites and antioxidant capacity (Table 1) were improved in Dia-VTM MB_{PRO} groups compared to control group as in other studies (Paryad and Mahmoudi 2008; Okasha *et al.*, 2023). Total protein and albumin (g/dl) were significantly increased in Dia-VTM MB_{PRO} groups compared to control one. This could be attributed to the significant improvement of feed intake and feed conversion obtained in those groups, which subsequently resulted in increased glucose and creatinine values. Total bilirubin values were decreased ($P=0.0001$) in Dia-VTM MB_{PRO} groups compared to control groups. Both *S. cerevisiae* and

essential oils have shown potential to improve liver health in poultry. By maintaining healthy liver function, these additives may contribute to efficient bilirubin processing and excretion in addition to unchanged liver enzymes (Table 1).

The antioxidant capacities were gradually decreased in Dia-VTM MB_{PRO} groups with increasing doses if compared to the control group except ABTS, μM . *S. cerevisiae* contains components, such as beta-glucans, that can stimulate the poultry's immune system, which enhances antioxidant defenses. Some studies have shown that *S. cerevisiae* can enhance the activity of antioxidant enzymes in poultry. Additionally, the essential oils are rich in bioactive compounds, such as terpenes and phenols, which possess potent antioxidant properties and modulate the activity of antioxidant enzymes (De Sousa *et al.* 2023; Sequeda-Castañeda *et al.*, 2025).




Fig 3: Effect of dietary Dia-VTM MB_{PRO} on feed conversion of broiler chickens at 35 d of age.




Fig 4: Effect of dietary Dia-VTM MB_{PRO} on dressing and giblets percentage of broiler chickens at 35 d of age.

CONCLUSION

It could be concluded that Dia-V™ MB_{PRO} supplement modulates growth performance, serum metabolites and antioxidant capacity. The best dietary supplement was 400 MB_{PRO} compared to 200 MB_{PRO} and 800 MB_{PRO} levels.

ACKNOWLEDGEMENT

The authors want to thank and acknowledge Deanship of Scientific Research, King Faisal University, Saudi Arabia for funding and support (KFU251404).

Conflict of interest

All authors declared that there is no conflict of interest.

REFERENCES

- Amma Z, Reiczigel J, Fébel H, Solti L. Relationship between Milk Yield and Reproductive Parameters on Three Hungarian Dairy Farms. *Veterinary Science*. 11(5): 218.
- Doumas, B.T., Watson, W.A., Biggs, H.G. (1971). Albumin standards and measurement of serum albumin with bromocresol green. *Clinica Chimica Acta*. 31: 87-96.
- Abd El-Ghany, W.A. (2024). Applications of Organic Acids in Poultry Production: An Updated and Comprehensive Review. *Agriculture*. 14(10): 1756.
- Abd El-Hack, M.E., El-Saadony, M.T., Shafi, M.E., Qattan, S.Y.A., Batiha, G.E., Khafaga, A.F., Abdel-Moneim, A.E., Alagawany, M. (2020). Probiotics in poultry feed: A comprehensive review. *Animal Physiology and Animal Nutrition*. <https://doi.org/10.1111/jpn.13454>.
- Abdel-Moneim, A.E., Mesalam, N.M., Yang, B., Elsadek, M.F. (2025). Dietary incorporation of biological curcumin nanoparticles improved growth performance, ileal architecture, antioxidative status, serum lipid profile, and and humoral immune response of heat-stressed broiler chickens. *Poultry Science*. 104(2): 104740.
- Abdullahi, A.M., Kalengyo, R.B. and Warsame, A.A. (2024). The unmet demand of food security in East Africa: Review of the triple challenges of climate change, economic crises, and and conflicts. *Discover Sustainability*. 5: 244.
- AL-Suwaiegh B.S. (2023). Effect of feeding green microalgae and bacillus subtilis on growth performance, blood metabolites and nutrient digestibility of beef bulls in arid subtropics. *Indian Journal of Animal Research*. 57(7): 882-888.
- Aviagen, R. (2018). Ross Broiler Management handbook Aviagen Limited New bridge Midlothian EH28 8SZ. pp. 1-113.
- Ayana, G.U., Kamutambuko, (2024). Probiotics in Disease Management for Sustainable Poultry Production. *Advanced Gut and Microbiome Research*. <https://doi.org/10.1155/2024/4326438>.
- Benzie, I.F.F., Strain J.J. (1996). The ferric reducing ability of plasma (FRAP) as measure of antioxidant power: The FRAP assay. *Analytical Biochemistry*. 239(1): 70-76.
- Blois, M.S. (1958). Antioxidant determinations by the use of a stable free radical. *Nature*. 26: 1199-1200.
- Brake, J. (1991). Research note: Lack of effect of a live yeast culture on broiler breeder and progeny performance. *Poultry Science*. 70: 1037-1039.
- Callaway, E.S., Martin, S.A. (1997). Effects of a *Saccharomyces cerevisiae* culture on ruminal bacteria that utilize lactate and digest cellulose. *Journal Dairy Science*. 80: 2035-2044.
- Chandrasekaran, P., Weiskirchen, S., Weiskirchen, R. (2024). Effects of Probiotics on Gut Microbiota: An overview. *International Journal of Molecular Sciences*. 25(11): 6022.
- De Sousa, D.P., Damasceno, R.O.S., Amorati, R., Elshabrawy, H.A., de Castro, R.D., Bezerra, D.P., Nunes, V.R.V., Gomes, R.C., Lima, T.C. (2023). Essential Oils: Chemistry and Pharmacological Activities. *Biomolecules*, 13(7): 1144.
- Dong, S., Li, L., Hao, F., Fang, Z., Zhong, R., Wu, J., Fang, X. (2024). Improving quality of poultry and its meat products with probiotics, prebiotics and phytoextracts. *Poultry Science*. 103(2): 103287.
- Duncan, D.B. (1955). Multiple range and multiple ftest. *Biometrics*. 11: 1.
- Ezzat Abd El-Hack, M., Alagawany, M., Ragab Farag, M., Tiwari, R., Karthik, K., Dhama, K., Zorriezhahra, J., Adel, M. (2016). Beneficial impacts of thymol essential oil on health and production of animals, fish and poultry: A review. *Journal of Essential Oil Research*. 28: 365-382.
- Gamage, A., Gangahagedara, R., Subasinghe, S., Gamage, J., Guruge, C., Senaratne, S., Randika, T., Rathnayake, C., Hameed, Z., Madhujith, T., Othmane Merah, O. (2024). Advancing sustainability: The impact of emerging technologies in agriculture. *Current Plant Biology*. 40: 100420.
- Giannenas, I., Bonos, E., Skoufos, I., Tzora, A., Stylianaki, I., Lazari, D., Tsinas, A., Christaki, E., Florou-Paneri, P. (2018). Effect of herbal feed additives on performance parameters, intestinal microbiota, intestinal morphology and meat lipid oxidation of broiler chickens. *British Poultry Science*. 59: 545-553.
- Haldar, S., Ghosh, T.K. and Bedford, M.R. (2011). Effects of yeast (*Saccharomyces cerevisiae*) and yeast protein concentrate on production performance of broiler chickens exposed to heat stress and challenged with salmonella enteritidis. *Animal feed science and technology*. 168(1-2): 61-71.
- Heinsohn, Z., Brown, A., Sobotik, E., House, G., Stiewert, A., Chaney, W.E., Kuttappan, V., and Archer, G.S. (2024). Evaluating the effects of feeding a concentrated *saccharomyces cerevisiae* fermentation product on the performance and stress susceptibility of broiler chickens. *Poultry*. 3(1): 57-65.
- Hu, M.L. (1994). Measurement of protein thiol groups and glutathione in plasma. *Methods Enzymology*. 233: 380-385.
- Jin, L.Z., Ho, Y.W., Abdullah, N., Jalaludin, S. (1997). Probiotics in poultry: Modes of action. *World's Poultry Science Journal*. 53: 351-368.
- Krysiak, K., Konkol, D., Korczyński, M. (2021). Overview of the Use of Probiotics in poultry production. *Animals (Basel)*. 11(6): 1620.
- Kumar, R., Sood, U., Gupta, V., Singh, M., Scaria, J., Lal, R. (2020). Recent advancements in the development of modern probiotics for restoring human gut microbiome dysbiosis. *Indian Journal Microbiology*. 60: 12-25.
- Lin, j., Comi, M., Vera, P., Alessandro, A., Qiu, K., Wang, J., Wu, S.-G., Qi, G.-H., Zhang, H.-J. (2023). Effects of *Saccharomyces cerevisiae* hydrolysate on growth performance, immunity function, and and intestinal health in broilers. *Poultry Science*. 102(1): 102237.

- Line, J.E., Bailey, J.S., Cox, N.A., Stern, N.J. (1997). Yeast treatment to reduce *Salmonella* and *Campylobacter* population associated with broiler chickens subjected to transport stress. *Poultry Science*. 76: 1227-1231.
- Mazziotta, C., Tognon, M., Martini, F., Torreggiani, E., Rotondo, J.C. (2023). Probiotics mechanism of action on immune cells and beneficial effects on human health. *Cells*. 12(1): 184.
- Medina, B., Girard, I.D., Jacotot, E., Julliard, V. (2002). Effect of a preparation of *Saccharomyces cerevisiae* on microbial profiles and fermentation patterns in the large intestine of horses fed a high fiber or a high starch diet. *Journal Animal Science*. 80(10): 2600-2609.
- Mihara, M., Uchiyama, M. (1978). Determination of malonaldehyde precursor in tissues by thiobarbituric acid test. *Analytical Biochemistry*. 86(1): 271-278.
- Mohammed A.A., Alshaibani N. (2025). Camel milk yield and composition and its adjuvant potential impacts on health of consumers: A review. *Indian Journal of Animal Research*. 59(2): 177-184.
- Mohammed, A.A., Almuyidi, A., Almarri, H., Alkhalifah, H., Alhmad, A., Alali, H., AlHuwaish, O. and AlKhawaher, M. (2025). Unique characteristics of camel body systems: Adaptation to harsh conditions, productive and reproductive performances: A review. *Indian Journal of Animal Research*. doi: 10.18805/IJAR.BF-1922.
- Moore, B.E., Newman, K.E., Spring, P., Chandler, F.E. (1994). The effect of yeast culture (Yea Sacc 1026) in microbial population's digestion in the cecum and colon of the equine. *Journal Animal Science*. 72: 1.
- Movahedi, F., Nirmal, N., Wang, Jin, H., Grøndal L., Li, L. (2024). Recent advances in essential oils and their nano-formulations for poultry feed. *Journal of Animal Science and Biotechnology*. 15: 110.
- Naeem, M. and Bourassa, D. (2025). Probiotics in poultry: Unlocking productivity through microbiome modulation and gut health. *Microorganisms*. 13(2): 257.
- Okasha, E.G., Ismail, Z.S.H., Hassan, A. (2023). Effects of yeast (*Saccharomyces cerevisiae*) supplementations on the blood parameters and productive performance of broiler chickens. *SVU-International Journal of Agricultural Sciences*. 5(4): 130-140.
- Paryad, A., Mahmoudi, M. (2001). Effect of different levels of supplemental yeast (*Saccharomyces cerevisiae*) on performance, blood constituents and carcass characteristics of broiler chicks. *African Journal of Agricultural Research*. 3(12): 835-842.
- Pawlak, K. and Kołodziejczak, M. (2020). The role of agriculture in ensuring food security in developing countries: Considerations in the context of the problem of sustainable food production. *Sustainability*. 12(13): 5488.
- Puvača, N., Ljubojević, D., Kostadinović, L., Lević, J., Nikolova, N., Mišćević, B., Könyves, T., Lukač, D., Popović, S. (2015). Spices and herbs in broilers nutrition: Hot red pepper (*Capsicum annuum* L.) and its mode of action. *Worlds Poultry Science Journal*. 71: 683-688.
- Puvača, N., Tufarelli, V., Giannenas, I. (2022). Essential oils in broiler chicken production, immunity and meat quality: Review of *Thymus vulgaris*, *Origanum vulgare* and *Rosmarinus officinalis*. *Agriculture*. 12(6): 874.
- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., Rice-Evans, C. (1999). Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine*. 26: 1231-1237.
- Santin, E., Maiorka, A., Macari, M., Grecco, M., Sanchez, J.C., Okada, T.M., Myasaka, A.M. (2001). Performance and intestinal mucosa development of broiler chickens fed diets containing *Saccharomyces cerevisiae* cell wall. *Journal of Applied Poultry Research*. 10: 236-244.
- SAS (2008). SAS user's guide: Basics. Statistical Analysis System Institute, Inc., Cary, NC, USA.
- Sequeda-Castañeda, L.G., Castellanos-Gómez, M.A., Céspedes-Acuña, C.L.A. (2025). Bioactive properties of *Pentacalia vaccinioides* (Kunth) Cuatrec. (Asteraceae) essential oils: Evaluation of antimicrobial and antioxidant activities. *Separations*. 12(1): 9.
- Shurson, G.C. (2018). Yeast and yeast derivatives in feed additives and ingredients: Sources, characteristics, animal responses, and quantification methods. *Animal Feed Science Technology*. 235: 60-76.
- Su, G., Wang, L., Zhou, X., Wu, X., Chen, D., Yu, B., Huang, Z., Luo, Y., Mao, X., Zheng, P., Yu, J., Luo, J., He, J. (2021). Effects of essential oil on growth performance, digestibility, immunity, and and intestinal health in broilers. *Poultry Science*. 100(8): 101242.
- Sun, Z., Wang, T., Demelash, N., Zheng, S., Zhao, W., Chen, X., Zhen, Y., and Qin, G. (2020). Effect of yeast culture (*Saccharomyces cerevisiae*) on broilers: A preliminary Study on the effective components of yeast culture. *Animals*. 10(1): 68.
- Swamy, M.N., Upendra, H.A. (2013). Growth performance, crude protein, ether extract and total ash in the breast muscle of broiler chickens supplemented with probiotics. *International Journal of Science, Environment and Technology*. 2: 1000-1007.
- Urban, J., Kareem, K.Y., Matuszewski, A., Bien, D., Ciborowska, P., Lutostanski, K., Michalczuk, M. (2024). Enhancing broiler chicken health and performance: The impact of phytobiotics on growth, gut microbiota, antioxidants, and and immunity. *Phytochemistry Reviews*. <https://doi.org/10.1007/s11101-024-09994-0>.
- Valdivieso-Ugarte, M., Gomez-Llorente, C., Plaza-Díaz, J., Gil, Á. (2019). Antimicrobial, antioxidant and immunomodulatory properties of essential oils: A systematic review. *Nutrients*. 11(11): 2786.
- Vermeulen, S.J., Park, T., Khoury, C.K., Béné, C. (2020). Changing diets and the transformation of the global food system. *Annals of the New York Academy of Sciences*. 1478(1): 13-17.