



Multidimensional Characterization and Adaptive Differentiation of Leptin Gene in *Eothenomys miletus*: Based on Bioinformatics and System Evolution Analysis

Huairan Wang¹, Wenrong Gao², Wanlong Zhu¹

10.18805/IJAR.BF-2042

ABSTRACT

Background: To investigate the differences between leptin gene in *Eothenomys Miletus* and other 55 mammal species, this study conducted a comparative analysis of multidimensional characteristics and adaptive differentiation of leptin genes.

Methods: In this study, molecular biology experimental strategies have been used to extract and reverse transcription, gene cloning and sequencing of *E. miletus* and bioinformatics evaluation techniques have been used to analyze the sequence and structure, multi-sequence alignment and evolutionary evaluation of *E. miletus* and ecological and distribution statistical strategies had been used to analyze the leptin genes of 56 mammalian species such as *E. miletus*.

Result: Leptin gene in *E. miletus* contained a Sec/SPI signal peptide, with α -helices dominating its structure. A model was constructed using leptin-LepR trimer (8×80.1 . C leptin) as template. It encodes 127 amino acids with high proportions of leucine and serine; its molecular formula is $C_{824}H_{1353}N_{221}O_{252}S_8$ (2658 atoms) with molecular weight of 18644.62 and theoretical isoelectric point of 5.86. The protein is unstable and hydrophilic, yet more hydrophobic than *Tupaia chinensis* and *Homo sapiens*. *E. miletus* was genetically close to *Microtus*, distant from *Lagomorpha* and clusters with temperate *Rodentia*. Its codon preference differed from *Lagomorpha*, with moderate GC content, GC3 content and ENC values. Motifs 1-5 were conserved in its leptin. Functional annotation shows regions related to signal peptide, leptin and extracellular domains. The ω value of ~ 0.5661 suggests evolutionary conservation. These results provide basis for exploring leptin gene evolution in *E. miletus* and mammals, offering perspective on mammalian adaptation to extreme environments and high-altitude protection.

Key words: Adaptation, Dimensional effect, *Eothenomys miletus*, Leptin gene, Mammals.

INTRODUCTION

Leptin, a conserved protein hormone encoded by the obesity gene and primarily secreted by white adipose tissue (Priyadarshini *et al.*, 2015), was identified in humans and mice in 1994 and has since become a key research focus. As a core regulator of energy metabolism, it modulates energy balance and suppresses food intake (Zhang *et al.*, 2009; Sun, 2019), while also participating in immune responses (Wei *et al.*, 2013), infectious diseases and reproduction (Jin *et al.*, 2021; Zhang, 2015). Molecular and genetic studies have explored leptin genes in animals (Pang *et al.*, 2016; Li *et al.*, 2021; Meena *et al.*, 2017), their associations with diseases (Yan *et al.*, 2023; Ma, 2012) and leptin receptor genes (Yan *et al.*, 2011; Li *et al.*, 2011). Its physiological roles, linked to gene structural and functional variations, are critical for thermoregulation and metabolic heat production under environmental temperature stress.

Eothenomys miletus, a Cricetidae rodent, which was endemic to the Hengduan Mountains of Yunnan (Han *et al.*, 2021), it has been studied by our team for adaptive strategies, including physiological ecology, such as correlations between serum leptin levels and physiological parameters under varying conditions (Hou *et al.*, 2020; Yang *et al.*, 2013; Yang *et al.*, 2016) and leptin-mediated regulation of body mass and energy metabolism via hypothalamic neuropeptides (Zhu, 2015). However, research on the

¹School of Life Sciences, Yunnan Normal University, Kunming, 650500, China.

²School of Biological Resources and Food Engineering, Qujing Normal University, Qujing, 655011, China.

Corresponding Author: Wanlong Zhu, School of Life Sciences, Yunnan Normal University, Kunming, 650500, China.

Email: zwl_8307@163.com

ORCIDs: 0009-0004-3687-4081; <https://orcid.org/0000-0001-9521-7613>; 0000-0001-8261-4089.

How to cite this article: Wang, H., Gao, W. and Zhu, W. (2025). Multidimensional Characterization and Adaptive Differentiation of Leptin Gene in *Eothenomys miletus*: Based on Bioinformatics and System Evolution Analysis. *Indian Journal of Animal Research*. 1-11. doi: 10.18805/IJAR.BF-2042.

Submitted: 11-07-2025 **Accepted:** 30-08-2025 **Online:** 18-09-2025

molecular evolutionary mechanisms of its leptin gene remains limited. To clarify the leptin gene's bioinformatics characteristics and evolutionary selection pressure, this study integrated leptin sequences from 55 mammals across five orders (*Rodentia*, *Dermoptera*, *Lagomorpha*, *Scandentia*, *Primates*) - taxa with wide distributions from equatorial tropics to cold zones, subject to strong temperature selection pressure. Recent studies note cold-dwelling *Vulpes lagopus* exhibits leptin receptor sensitivity,

while tropical *Tupaia belangeri* has unique leptin expression regulation (Zhu *et al.*, 2014), indicating temperature-driven adaptive evolution of leptin. However, the universality of such latitude- and temperature-related adaptations across taxa and cross-order evolutionary patterns of leptin genes, remain unclear.

This study analyzed the leptin gene's protein structure and physicochemical properties in *E. miletus* and performed cross-order comparative analyses of leptin sequences from 56 mammals (including *E. miletus*) to reveal the gene's molecular characteristics, adaptive evolutionary signals and cross-order evolutionary patterns in mammals.

MATERIALS AND METHODS

Obtaining the complete sequence of leptin gene from *E. miletus*

From 2024 to June 2025, experiments and follow-up data analysis is conducted at the School of Life Sciences of Yunnan Normal University.

Primers were designed according to the sequence of leptin gene in *E. miletus*. The upstream primer F was 5'-ATGTGCTGGGAGACTCCTGTG-3' and the downstream primer R was 3'-CAGCATCAGCTAAGGTCC-5'. Primers were synthesized by the company. The size of the amplified product was 503bp and the temperature of cleavage was 89.8°C. The steps and operations of extracting and purifying total RNA from hypothalamus in *E. miletus* was details in Zhu's (2015). The whole sequence is transmitted to NCBI (GenBank No: pv838002).

Sequence analysis of leptin gene in *E. miletus*

Signal peptide prediction and protein secondary and tertiary structure analysis

The sign peptide of leptin protein was once estimated by using SignalP-6.0 on-line tool, the secondary shape of leptin protein was once envisioned through SOPMA on line software and the tertiary shape of leptin protein used to be analyzed by means of SWISS-MODEL on line software.

Analysis of physicochemical properties of leptin protein from *E. miletus*

The hydrophilicity and hydrophobicity of leptin protein in *E. miletus*, *Ochotona princeps*, *Tupaia belangeri* and *Homo sapiens* were predicted and compared by using the ProtScale Program in the online software of ExPASy. The physical and chemical properties of rat leptin protein were analyzed by using the ProtParam program of ExPASy online software.

Cross order comparison of leptin gene sequences in 56 species

Collation sequence

Leptin gene sequences from 55 mammalian species available on the NCBI platform, along with the newly obtained sequence from *E. miletus*, were organized (Table 1). These sequences are sorted out by means of techniques such as (Mao *et al.*, 2022; Yang, 1997).

Construction of phylogenetic tree

The systematic tree was constructed by the neighbor-joining (NJ) method and then the optimal model and the maximum likelihood method (ML) were used to construct the systematic tree and the self-expansion test (Bootstrap) was set to set 1000 iterations. Compare tree structures from different methods to ensure accuracy. Beautify the phylogenetic tree using Chiplot online software.

Distribution analysis of 56 species in five temperature zones

From the perspective of ecological and biological distribution, the main distribution of species is stable survival and reproduction, with large population, high density and wide distribution range. If the species are mainly distributed in a certain temperature zone, it is recorded as 1 and other distributions are recorded as 0. With the support of animal geographical distribution, ecological habits, professional literature and database, the main distribution of 56 species in 5 temperature zones were counted by Excel. Chiplot was used to analyze the clustering of 56 species in each temperature zone.

Codon preference and determination of GC content

The number of effective codons (ENC), GC content and third GC content (GC3s) of the sequence were calculated using the online tool EMBOSS Explorer.

Conservative region determination and functional annotation analysis of multiple sequences

MEME online software is used to predict the conservative motifs of multiple sequences, compare the predicted conservative motifs with Pfam, PROSITE and other databases, annotate the functional domains corresponding to the motifs and visualize the conservative regions through TBtools - II.

Select pressure detection

PAML software was used to calculate the ratio of nonsynonymous replacement rate (dN) and synonymous replacement rate (dS) ($\omega = dN/dS$) to measure the selection pressure of leptin gene in the process of evolution. The single ratio (M0) and free ratio (M1) branching models that rely on likelihood ratio test (LRT) in Codeml program are used to detect the overall selection pressure of leptin gene and whether there is heterogeneity of selection pressure in different evolutionary branches of leptin gene.

RESULTS AND DISCUSSION

Bioinformatics analysis of leptin in *E. miletus*

Signal peptide assay

It showed that there was a highly reliable Sec/SPI type signal peptide in the leptin protein of *E. miletus* (Fig 1). The probability distribution of different regions of signal peptide is high and the probability of non-signal peptide region is low, which is consistent with the judgment of signal peptide.

Table 1: Basic information of 56 species of mammals and NCBI entry numbers.

Specie latin name	Order	Family	NCBI Accession No.	GC content	GC3 content	ENC
<i>Cercocebus atys</i>	Primates	Cercopithecidae	XM_012087965.1	54.17	73.81	45.13
<i>Macaca mulatta</i>	Primates	Cercopithecidae	U58492.1	53.97	73.81	45.08
<i>Macaca thibetana thibetana</i>	Primates	Cercopithecidae	XM_050782581.1	54.56	73.16	43.33
<i>Rhinopithecus bieti</i>	Primates	Cercopithecidae	XM_017896214.1	54.37	74.40	44.86
<i>Rhinopithecus roxellana</i>	Primates	Cercopithecidae	XM_010378726.1	54.37	74.40	44.86
<i>Trachypithecus francoisi</i>	Primates	Cercopithecidae	XM_033192942.1	54.17	73.81	44.28
<i>Gorilla gorilla gorilla</i>	Primates	Hominidae	XM_004046161.4	54.76	73.21	45.33
<i>Homo sapiens</i>	Primates	Hominidae	NM_000230.3	54.56	72.02	46.92
<i>Pan paniscus</i>	Primates	Hominidae	XM_003806045.5	54.56	72.62	46.53
<i>Pan troglodytes</i>	Primates	Hominidae	U96450.1	55.56	75.51	42.56
<i>Pongo pygmaeus</i>	Primates	Hominidae	U72873.1	55.56	74.83	42.66
<i>Microcebus murinus</i>	Primates	Cheirogaleidae	XM_012789083.1	56.35	73.81	47.50
<i>Hylobates moloch</i>	Primates	Hylobatidae	XM_032753530.2	54.76	72.62	46.61
<i>Symphalangus syndactylus</i>	Primates	Hylobatidae	XM_055281456.1	54.56	72.02	47.66
<i>Octodon degus</i>	Rodentia	Abrocomidae	XM_004642767.2	48.90	64.07	55.54
<i>Heterocephalus glaber</i>	Rodentia	Abrocomidae	XM_004856402.2	50.40	65.48	46.76
<i>Eothenomys miletus</i>	Rodentia	Cricetidae	PV838002	54.76	71.43	42.54
<i>Alexandromys fortis</i>	Rodentia	Cricetidae	XM_050141585.1	55.16	72.02	46.33
<i>Arvicola amphibius</i>	Rodentia	Cricetidae	XM_038321033.1	55.36	71.43	46.74
<i>Cricetulus griseus</i>	Rodentia	Cricetidae	XM_027391330.2	53.79	67.26	52.32
<i>Meriones unguiculatus</i>	Rodentia	Cricetidae	XM_021659887.2	56.13	75.29	44.02
<i>Mesocricetus auratus</i>	Rodentia	Cricetidae	XM_005078071.4	54.56	69.64	48.70
<i>Microtus ochrogaster</i>	Rodentia	Cricetidae	XM_005365724.2	54.76	70.83	47.53
<i>Microtus oregoni</i>	Rodentia	Cricetidae	XM_041675323.1	53.97	68.45	51.11
<i>Myodes glareolus</i>	Rodentia	Cricetidae	XM_048451023.1	55.16	72.62	42.61
<i>Peromyscus californicus insignis</i>	Rodentia	Cricetidae	XM_052721409.1	56.05	72.63	47.48
<i>Peromyscus leucopus</i>	Rodentia	Cricetidae	XM_028866260.2	54.76	70.83	49.49
<i>Phodopus roborovskii</i>	Rodentia	Cricetidae	XM_051175915.1	54.56	70.83	49.67
<i>Castor canadensis</i>	Rodentia	Castoridae	XM_020182860.1	52.98	69.05	43.56
<i>Chinchilla lanigera</i>	Rodentia	Chinchillidae	XM_005402367.2	53.49	71.26	47.46
<i>Apodemus sylvaticus</i>	Rodentia	muridae	XM_052172451.1	54.37	69.64	42.19
<i>Arvicanthis niloticus</i>	Rodentia	muridae	XM_034518822.1	54.34	68.45	44.19
<i>Mastomys coucha</i>	Rodentia	muridae	XM_031380792.1	53.57	68.45	47.64
<i>Mus caroli</i>	Rodentia	muridae	XM_021165722.1	53.37	67.26	42.54
<i>Mus musculus</i>	Rodentia	muridae	NM_008493.3	53.97	68.45	47.47
<i>Mus pahari</i>	Rodentia	muridae	XM_029535370.1	54.96	69.64	44.45
<i>Onychomys torridus</i>	Rodentia	muridae	XM_036180469.1	55.36	72.02	52.24
<i>Peromyscus eremicus</i>	Rodentia	muridae	XM_059257460.1	56.55	75.00	49.79
<i>Peromyscus maniculatus bairdii</i>	Rodentia	muridae	XM_006979375.3	55.16	72.02	47.33
<i>Psammomys obesus</i>	Rodentia	muridae	XM_055608191.1	56.15	74.40	44.18
<i>Rattus norvegicus</i>	Rodentia	muridae	NM_013076.3	55.95	73.21	45.43
<i>Rattus rattus</i>	Rodentia	muridae	XM_032905243.1	56.15	73.81	44.29
<i>Ictidomys tridecemlineatus</i>	Rodentia	Sciuridae	XM_047562352.3	54.17	66.67	48.81
<i>Marmota marmota marmota</i>	Rodentia	Sciuridae	XM_015488308.2	54.17	66.67	48.81
<i>Marmota monax</i>	Rodentia	Sciuridae	XM_046422457.1	54.37	66.67	49.31
<i>Sciurus carolinensis</i>	Rodentia	Sciuridae	XM_015488308.1	55.56	71.43	49.50
<i>Jaculus jaculus</i>	Rodentia	Dipodidae	XM_004658616.1	54.17	71.43	45.10
<i>Cavia porcellus</i>	Rodentia	Caviidae	XM_003475050.3	50.60	65.66	58.39

Table 1: Continue...

Functional annotation analysis of multiple sequence conservative motifs

Functional annotation was performed on motif 1-11, some conservative motifs had no functional annotation (Fig 9). Leptin was concentrated near the 20-25 amino acids. The region of membrane bound protein predicted to be outside the membrane, in the extracellular region, is expected to be located outside the membrane, occupying the posterior segment of the sequence (about 25-50 positions). Motif 1-3 showed the functional region of the conserved motif of the protein sequence encoded by the *E. miletus* leptin gene sequence.




Fig 4: Three-dimensional structure prediction result of leptin protein in *E. miletus*.

Analysis of selection pressure and positive selection effect

The single ratio model selection pressure analysis (Table 3) shows that the overall $\omega=0.32993<1$ and the leptin gene dN (1.5749) is significantly lower than dS (4.7733), indicating the presence of significant purification (negative) selection. *E. miletus* and *Microtus* gather together. In some branches, synonymous substitutions of rodents decrease, while in others, synonymous substitutions increase. In addition, the average content of GC3 between sequences is 0.7142.

Free ratio model-based branch selection pressure analysis showed a better overall likelihood value (LNL = -5630.47) than the single ratio model (LNL = -5721.88). Likelihood ratio test (LRT) confirmed significant heterogeneity in selection pressure among branches. The overall kappa ($ts/tv = 4.13$) indicated a high transition/transversion ratio, consistent with mammalian gene evolution characteristics. With total nonsynonymous substitution rate (dN = 1.62) lower than synonymous substitution rate (dS = 4.55; dN < dS), the overall selection remained dominated by purifying selection. Under the free ratio model, *Microtus* and *E. miletus* ($\omega = 0.5661$) clustered into one branch. Most species underwent strong purifying selection, while some branches showed significant positive selection.

Sequence characteristics of leptin gene and protein structure of *E. miletus* adapted to their living needs

This study undertook an analysis of the leptin gene in *E. miletus*, predicting that its leptin protein contains a Sec/SPI-type signal peptide with a notably high support probability. Leptin functions by binding to its receptor




Fig 5: Prediction results of leptin affinity/hydrophobicity of *O.*

(LepR), with α -helices playing a pivotal role in receptor recognition. The leptin of *E. miletus* exhibits a high α -helix ratio, which is conducive to a stable signal transduction domain. Research on *Ochotona curzoniae* has demonstrated that elevated α -helix ratios enhance receptor affinity and thermogenesis, facilitating adaptation to extreme cold conditions (Yang *et al.*, 2011). Compared to humans and mice, *E. miletus* leptin displays a higher α -helix ratio, which is essential for energy regulation in adapting to high-altitude, cold environments. Its low random coil ratio indicates a compact structure, thereby reducing energy consumption. These findings, coupled with high sequence consistency with the leptin-LepR trimer template, confirm structural compatibility and enhance receptor binding efficiency in cold environments.

E. miletus leptin (127 amino acids) has the highest leucine proportion, potentially enhancing energy reserve and utilization in alpine environments (Zhang *et al.*, 2009). Its amino acid composition is more "energy-efficient" than human leptin ($\approx 11\%$ leucine), aligning with survival needs in high-altitude, food-deficient habitats. As an unstable hydrophilic protein requiring continuous *in vivo* synthesis,

its theoretical half-life indicates eukaryotic stability, ensuring sustained energy regulation signals.

The hydrophobic region interacts with the phospholipid bilayer in transmembrane structures and maintains the conformational stability of membrane proteins at low temperatures (Tomczak *et al.*, 2002). *E. miletus* exhibits high hydrophobicity, which facilitates protein structural maintenance and receptor signal transduction efficiency in cold environments (Chen *et al.*, 2022). Plateau cold-zone species (*O. princeps* and *E. miletus*) showed strong hydrophobicity, ensuring stability and response speed of energy regulation signals. In contrast, temperate *H. sapiens* and tropical *T. belangeri* balance hydrophilicity and hydrophobicity, likely meeting the energy metabolism and environmental stability requirements.

Multidimensional characteristics of *E. miletus* adapted to living environment

Further evolutionary analysis showed that *E. miletus* and *Microtus* clustered, indicating leptin gene conservation in the same family and genus. Rodentia families have longer branches, with niche differentiation possibly driving leptin




Fig 6: Phylogenetic tree of 56 mammalian leptin genes.

gene differentiation. Primates and hominids cluster closely, reflecting high leptin gene conservation.

In temperature distribution, Primates, *T. belangeri* and Dermoptera are mostly tropical. Ochotonidae and Leporidae differ in differentiation: some species range from tropical to temperate zones, with their leptin genes scattered in temperate clusters, showing environment-adaptive polymorphism to meet energy needs across

temperature zones, reflecting molecular-level "ecological range expansion" (Londrville *et al.*, 2014; Gong *et al.*, 2013). *E. miletus*, mainly temperate, uses leptin gene regulation to adapt to the Hengduan Mountains' extreme, climate-variable, large temperature-difference environment.

Molecular-level analyses of codon preference and conserved motifs further support these conclusions. The leptin gene shows a preference for G/C-ending codons;




Fig 7: Clustering heat map of the distribution of 56 species in each temperature zone.




Fig 8: Results of conserved motif determination of 56 leptin gene sequences.




Fig 9: Functional annotation results of each conserved motif.

Table 3: Reference values of leptin gene results under two branch models.

Model	tree length for dN/dS	lnL	tree length	kappa (ts/tv)	GC3 content
M0	1.6241/4.5475	-5630.468199	7.148581	4.12861	C:0.39517 G:0.32513
M1	1.5749/4.7733	-5721.884113	7.214167	4.11744	C:0.39517 G:0.32513

high GC3 content enhances DNA stability, aiding adaptation to UV-intense tropical environments and cold zones with large temperature fluctuations (Sambrook and Russell, 2001). A lower ENC value indicates stronger codon preference and faster translation (Sharp and Li, 1987); the moderate ENC value of *E. miletus* leptin suggests a trade-off between rapid translation and sequence variation, balancing core energy regulation functions with temperature zone specialization demands. Motifs 1-5, as core functional domains, ensure cross-species energy regulatory basics. The distribution of motifs 1-3 reveals *E. miletus* leptin retains signal peptide-guided secretion, metabolic regulation and extracellular membrane protein functions, linked to adaptation to the Qinghai-Tibet Plateau's high-altitude, cold and hypoxic conditions (Chen *et al.*, 2022; Gong *et al.*, 2022).

Selection pressure analyses showed overall purifying selection on the leptin gene, with highly conserved functions. Some Rodentia branches exhibited higher dS, indicating rapid evolution and sequence divergence, potentially linked to adaptive radiation. Most rodents clustered closely with short branch lengths and low genetic distances. *Microtus* clusters with *E. miletus*, supporting their close phylogenetic relationships. Leptin function is highly conserved among congeneric rodent species, with consistent patterns observed in other taxa.

CONCLUSION

This study elucidates the molecular mechanisms underlying *Eothenomys miletus* adaptation to high-altitude

environments in the Hengduan Mountains. Key findings demonstrate that: Leptin gene evolution reflects long-term selective pressures under plateau conditions, with specific mutations stabilizing energy-regulation signaling pathways while maintaining metabolic flexibility to cope with resource fluctuations. Physiological specialization at the molecular level provides a paradigm for mammalian adaptation to extreme environments, bridging a critical gap between genetic changes and ecosystem-level resilience. These results offer two-fold implications: Theoretically, establishing leptin as a biomarker for high-altitude adaptation, challenging traditional views on metabolic adaptation timelines. Practically, proposing molecular targets for monitoring vulnerable plateau ecosystems, informing conservation strategies under climate change.

ACKNOWLEDGEMENT

This work was supported by the National Natural Scientific Foundation of China (No. 32160254), the Special Basic Cooperative Research Programs of Yunnan Provincial Undergraduate Universities' Association (NO. 202301 BA070001-076).

Ethical approval

All animal procedures were within the rules of Animals Care and Use Committee of School of Life Sciences, Yunnan Normal University. This study was approved by the committee (13-0901-011).

Conflict of interest

The authors declare that there are no conflicts of interest regarding the publication of this article. No funding or sponsorship influenced the design of the study, data collection, analysis, decision to publish, or preparation of the manuscript.

REFERENCES

- Chen, H., Zhang, H., Jia, T., Wang, Z. and Zhu, W. (2022). Roles of leptin on energy balance and thermoregulation in *Eothenomys miletus*. *Frontiers in Physiology*. **13**: 1054107. doi: 10.3389/fphys.2022.1054107.
- Gong, N., Einarsdottir, I.E., Johansson, M. and Björnsson, B.T. (2013). Alternative splice variants of the rainbow trout leptin receptor encode multiple circulating leptin-binding proteins. *Endocrinology*. **154**(7): 2331-2340. doi: 10.1210/en.2012-2082.
- Gong, X.N., Jia, T., Zhang, D., Zhu, W.L. (2022). faster response to high-fat diet in body mass regulation from lower altitude population in *eiothenomys miletus*. *Pakistan Journal of Zoology*. **54**(1): 167-177. doi: 10.17582/journal.pjz/20200211050216.
- Han, C.Y., Jia, T., Yang, J., Wang, Z.K., Zhu, W.L. (2021). Body mass regulation by *Eothenomys miletus* at different areas of the hengduan mountains in winter. *Chinese Journal of Wildlife*. **42**(1): 86-91. doi: 10.3969/j.issn.1000-0127.20
- Hou, D.M., Ren, X.Y., Zhu, W.L., Zhang, H. (2020). Comparative study on phenotypic differences in *Eothenomys miletus* under food restriction and refeeding between xianggelila and jianchuan from hengduan mountain regions. *Indian Journal of Animal Research*. **54**(7): 835-840. doi: 10.18805/ijar.B-1151.
- Jin, M.H., Huang, W.Y., Zhang, M. Y., Zhang, Y.W., Liu, Y., Ding, Z.D. (2021). Leptin and Male Reproduction. *Journal of International Reproductive Health/Family Planning*. **40**(1): 38-43.
- LI, G.H., Chen, Q.Y., Wu, W.D., Zheng, X.B. (2021). Eukaryotic expression and biological characterization of leptin gene in buffalo. *Journal of Henan Agricultural Sciences*. **50**(10): 132-137. doi: 10.15933/j.cnki.1004-3268.2021.10.017.
- Li, L., Ou, Z.G., Wang, L.J., Zhang, H.P., Du, L.X. (2011). Sequencing and analysis of ovine partial leptin peceptor gene and mutation loci. *China Animal Husbandry and Veterinary Medicine*. **38**(8): 107-113.
- Londrville, R.L., Macotela, Y., Duff, R.J., Easterling, M.R., Liu, Q. and Crespi, E.J. (2014). Comparative endocrinology of leptin: Assessing function in a phylogenetic context. *General and Comparative Endocrinology*. **203**: 146-157. doi: 10.1016/j.ygcen.2014.02.002.
- Ma, X.B., Che, X.W., Wu, S., Jin, Y.L., Zhang, N., Guo, J. Q., Xu, W.H. (2012). Relations between serum leptin and leptin receptor Gln223 Arg polymorphism with gastro-esophageal reflux disease. *Journal of Shandong University (Health Sciences)*. **50**(12): 65-69. doi: 10.6040/j.issn.1671-7554.2012.12.013.
- Mao, J.P., Huang, L.W., Hao, J., Liu, T.Y., Huang, S.W. (2022). Phylogenetic and molecular evolution analyses of *DXS* gene in plants. *Journal of Biology*. **39**(2): 23-28. doi: 10.3969/j.issn.2095-1736.2022.02.023.
- Meena, A.S., Bhatt, R.S., Sahoo, A., Kumar, S. (2017). Polymorphism of the exon 3 of leptin gene in Malpura sheep. *Indian Journal of Animal Research*. **51**(3): 469-473. doi: 10.18805/ijar.v0i0f.3783.
- Pang, C.Y., Deng, T.X., Lu, X.R., Zhu, Peng, Duan, A.Q., Liang, X.W. (2016). Single nuclease polymorphism analysis of leptin gene in buffalo. *China Animal Husbandry and Veterinary Medicine*. **43**(9): 2418-2424. doi: 10.16431/j.cnki.1671-7236.2016.09.029.
- Priyadarshini, L., Yadav, A.K., Singh, H.S., Mishra, A., Jain, A.K., Ahirwar, M.K. (2015). Role of leptin in physiology of animal reproduction-A review. *Agricultural Reviews*. **36**(3): 235-240. doi: 10.5958/0976-0741.2015.00027.6.
- Sambrook, J., Russell, D.W. (2001). Molecular Cloning: A Laboratory Manual (3rd ed.). Cold Spring Harbor Laboratory Press.
- Sharp, P.M. and Li, W.H. (1987). The codon adaptation Index-a measure of directional synonymous codon usage bias and its potential applications. *Nucleic acids Research*. **15**(3): 1281-1295. doi: 10.1093/nar/15.3.1281.
- Sun, P. (2019). Effects of hypothalamic VMH leptin116-130 on food and water intake and energy metabolism in rats and its potential mechanism. Doctoral thesis, Qingdao University.
- Tomczak, M.M., Hinch, D.K., Estrada, S.D., Wolkers, W.F., Crowe, L.M., Feeney, R.E., Tablin, F. and Crowe, J.H. (2002). A mechanism for stabilization of membranes at low temperatures by an antifreeze protein. *Biophysical Journal*. **82**(2): 874-881. doi: 10.1016/S0006-3495(02)75449-0.
- Wei, T.F., Tang, G.S., Shen, Q. (2013). The effects of Leptin system On the differentiation, proliferation and function of regulatory T lymphocytes. *International Journal of Immunology*. **36**(2): 97-100, 111. doi: 10.3760/cma.j.issn.1673-4394.2013.02.004.
- Yan, M.L., Zhang, J., Ye, X.M., Li, A.M. (2023). Correlation of leptin, leptin receptor level and gene polymorphism with Kawasaki's disease. *Chinese Journal of Modern Drug Application*. **17**(6): 26-30. doi: 10.14164/j.cnki.cn11-5581/r.2023.06.007.
- Yan, W., Luo, Y.Z., Yang, Q., Cheng, S.R., Hu, J. (2011). Polymorphism of leptin receptor gene (LEPR) in yak (*Bos grunniens*). *Journal of Agricultural Biotechnology*. **19**(2): 323-329. doi: 10.3969/j.issn.1674-7968.2011.02.018.
- Yang, J., Bromage, T. G., Zhao, Q., Xu, B.H., Gao, W.L., Tian, H.F., Tang, H.J., Liu, D.W. and Zhao, X.Q. (2011). Functional evolution of leptin of *Ochotona curzoniae* in adaptive thermogenesis driven by cold environmental stress. *PLoS one*. **6**(6): e19833. doi: 10.1371/journal.pone.0019833.
- Yang, S.C., Zhu, W.L., Zheng, J., Zhang, D., Mu, Y., Wang, Z.K. (2013). Effect of mild and sever food restriction on body mass and thermogenesis in *Eothenomys miletus*. *Sichuan Journal of Zoology*. **32**(4): 508-514. doi: 10.3969/j.issn.1000-7083.2013.04.006.
- Yang, T., Fu, J.H., Chen, J.L., Ye, F.Y., Zuo, M.L., Hou, D.M., Zhu, W.L. (2016). Effect of seasonal simulation on energy metabolism of *Eothenomys miletus*. *Sichuan Journal of Zoology*. **35**(3): 414-420. doi: 10.11984/j.issn.1000-7083.20160013.
- Yang, Z. (1997). PAML: A program package for phylogenetic analysis by maximum likelihood. *Computer applications in the biosciences: CABIOS*. **13**(5): 555-556. doi: 10.1093/bioinformatics/13.5.555.

- Zhang, M., Du, Z. H., Bai, X.J. (2009). Correlation analysis of leptin and receptor gene polymorphism and production performance of the Arctic fox. *Journal of Northeast Agricultural University*. **40(9)**: 75-81. doi: 10.3969/j.issn.1005-9369.2009.09.016.
- Zhang, Q., Qi, Z.T., Ding, Z.S. (2009). Molecular mechanism of leucine and its metabolites improving protein synthesis and the effects of supplementation. *Chemistry of Life*. **29(1)**: 103-107.
- Zhang, Y. (2015). Research advance on the relationship between leptin and reproduction activities of mammal. *China Animal Husbandry and Veterinary Medicine*. **42(7)**: 1836-1841. doi: 10.16431/j.cnki.1671-7236.2015.07.031.
- Zhu, W.L. (2015). Regulation of hypothalamic neuropeptides (NPY, AgRP, POMC, CART) on body weight and energy metabolism in rats. Doctoral thesis, Nanjing Normal University. doi: 10.7666/d. Y2857380.
- Zhu, W.L., Cai, J.H., Zhang, L., Wang, Z.K. (2014). Seasonal changes of body mass, serum leptin levels and hypothalamic neuropeptide express levels in *Tupaia belangeri*. *Journal of Biology*. **31(3)**: 33-37. doi: 10.3969/j.issn.2095-1736.2014.03.033.