



Okra Plant: A Multi-purpose Underutilized Vegetable Crop: A Review

Pavan Gupta, Shamayita Patra

Shri Vaishnav Institute of Textile Technology, Shri Vaishnav Vidyapeeth Vishwavidyalaya, Indore-453 111, Madhya Pradesh, India.

Received: January 2021

Accepted: June 2021

ABSTRACT

The plant kingdom has various species that are underutilised. Proper scientific approach help them to utilised for multiple end applications. Okra is one of such underutilized multi-purpose vegetable crop. Green immature okra fruits are consumed as a vegetable worldwide. Okra seed is a good source of oil and protein. Okra seeds are also used as a substitute for coffee. Okra roots are used for clarification of sugarcane juice. It has medicinal value as well as useful in wastewater treatment. Okra is also a potential source of natural fibers suitable for textile, paper and other engineering applications. The present paper reviewed the okra plant, its cultivation and applications in various form such as food, medicine, textile fiber, paper pulp, biomass, etc. for its introduction as a multi-purpose vegetable crop.

Key words: Agricultural waste, *Kharif* crop, Okra fiber, Okra plant.

The phrase "Food, clothing and shelter" shows the importance of food in human life. The major portion of global food demand is catered by agricultural products such as food grains (cereals and legumes), pulses, oilseeds, fruits and vegetables. Agricultural crops are mostly used for food production. The plant kingdom has various species which can be used for multiple purposes as well (Santulli *et al.* 2014). Few of such purposes are textiles, medical science, paper pulp, biofuel, etc. (Dungani *et al.* 2016; Reddy and Yang, 2005; Sen *et al.* 1987). A few of them are Okra, Banana, Pineapple, Coconut, Cornhusk, Jute, Flax, Maize, etc. (Dungani *et al.* 2016; Kambli *et al.* 2019; Rai *et al.* 2011; Reddy and Yang, 2005; Sen *et al.* 1987).

Okra is one of the underutilized multi-purpose crops in India. India ranks first in the world with 5,849 thousand ton Okra production (Ministry of Agriculture and Farmers Welfare 2017; Ministry of Environment and Forest 2011). It is a vegetable crop. It grows once (Summer) in the hills, twice or thrice (summer, *Kharif* and Late *Kharif*) in the eastern, western and northern Indian plains and almost the year in south India (Ministry of Environment and Forest 2011). Okra is valued for its green immature edible seed pods. Okra fruits are generally consumed as a vegetable in India. It can also be used for coffee, oil and protein as a part of food. Other than food application okra plant can be used for medical, textile, paper pulp and fuel purposes. The present paper reviews the okra plant, its cultivation and applications (food grade and non-food grade) for its introduction as a multi-purpose *Kharif* crop. Also, it advocates farmers for growing okra plant for additional income generation, while preventing production of greenhouse gases during burning (commonly practiced).

Okra plant

Okra (*Abelmoschus esculentus*) is the only vegetable crop which belongs to family Malvaceae. It is one of the oldest cultivated crops (Chanchal *et al.* 2018; Fathima and Balasubramanian, 2006; Martin 1982; Ministry of Environment and Forest 2011; Osawaru *et al.* 2014; Sathish and Eswar, 2013; Ullah *et al.* 2018; Varmudy 2011). It is grown in tropical and sub-tropical regions of the world (Chanchal *et al.* 2018; Jain *et al.* 2012; Makhadmeh and Ereifej, 2004; Martin 1982; Ministry of Environment and Forest 2011; Sen *et al.* 1987). It is an important vegetable crop especially in India, West Africa and Brazil (Joh *et al.* 2015). It is quite popular especially in India due to easy cultivation, drought resistance, dependable yield and adaptability to varying moisture conditions (Kocak *et al.* 2018; Ministry of Environment and Forest 2011; Sathish and Eswar, 2013).

Eight *Abelmoschus* species have been found in India. Out of these *A. esculentus* is the only known cultivated species. *A. moschatus*, a wild species, is also cultivated for its aromatic seeds. Okra is mainly propagated by seeds. It matures within 90-180 days. It is generally a perennial plant with a robust erected stem with variable branches. Okra plant's height can vary from 0.5 to 4.0 m (Agrawal *et al.* 2004; Ministry of Environment and Forest 2011; Moosavi *et al.* 2018; Sarasini and Fiore, 2018).

Okra requires a long, warm and humid cultivation period. It grows easily in hot humid climate. However, it is sensitive to frost and extreme low temperatures. The preferred temperature for its growth and development is 24-28°C. Adaptability with climatic factors helps them to grow once (summer) in the hilly areas, twice or thrice (summer, *Kharif* and late *Kharif*) in the eastern, western and northern Indian

plains and almost throughout the year in south India. It has well-developed tap root system. Therefore, it can grow in a wide variety of soil type i.e. sandy to clay soils. Loose, friable, well-manured loam soil is suitable for its cultivation. The soil with a pH range of 6.0–6.8 is desirable for okra plant (Ministry of Environment and Forest 2011; Moosavi *et al.* 2018; Sharma and Prasad, 2010). In India, the major okra producing states are West Bengal, Bihar, Orissa Andhra Pradesh, Gujarat, Jharkhand, Chhattisgarh, Maharashtra and Madhya Pradesh.

Applications of Okra Plant

Okra as food source

Okra is mainly cultivated for its green non-fibrous immature fruit. Young fruits are harvested before full seed development and mostly consumed as a vegetable in a variety of ways or as alone (Joh *et al.* 2015; Martin 1982; Ministry of Environment and Forest 2011). The mature okra seed has superior nutritional quality. Okra is a rich source of vitamin A, vitamin B, vitamin C, folic acid, amino acids, calcium, zinc, iron, potassium and dietary fibers (Table 1). Protein (30%) and oil (15-19%) contained in the seeds of okra serve as the source of first-rate vegetable protein. The yield of protein and oil from okra is comparable to the yield of the same from soybean (Camciuc *et al.* 1998; Chanchal *et al.* 2018; Joh *et al.* 2015; Martin 1982; Moosavi *et al.* 2018; Sathish and Eswar, 2013). Its ripe seeds are roasted, ground and used as a substitute for coffee in some countries that has a good aroma while lacks the stimulating effect of caffeine. The tender leaves of okra plant are often used as a vegetable in areas where a wide variety of leaves are used in the food (West Africa, Southeast Asia). Its seeds are used in place of legumes in soups. In West Africa the okra fruits are sliced, sun-dried and stored for future use (Martin 1982).

Table 1: Raw okra nutrition value per 100g (Chanchal *et al.*).

Energy	33 kcal
Carbohydrates	7.45 g
Sugars	1.48 g
Dietary Fibers	3.2 g
Fat	0.19 g
Protein	1.93-2 g
Water	90.19 g
Vitamin A	36ig (7%)
Thiamine(B1)	0.2 mg (17%)
Riboflavin(B2)	0.06 mg (5%)
Niacin (B3)	1mg (7%)
Vitamin C	23mg (28%)
Vitamin E	0.27 mg (2%)
Vitamin K	31.3 mg (30%)
Calcium	82 mg (8%)
Iron	0.62 mg (5%)
Magnesium	57 mg (16%)
Potassium	299mg (6%)
Zinc	0.58 mg (6%)

Okra for medicine

Fruits, leaves, root, seed and flower of okra are used in Indian ethnomedicine for treating dysentery, diarrhoea, kidneys catarrhal infections, hair treatment, *etc.* (Chanchal *et al.* 2018). Its medicinal value has also been reported in curing ulcers and relief of hemorrhoids (Chanchal *et al.* 2018; Joh *et al.* 2015; Ministry of Environment and Forest 2011; Roy *et al.* 2014; Sathish and Eswar, 2013). The mucilage and dietary fibers present in okra helps in adjusting blood sugar (Joh *et al.* 2015). Young leaves of okra have shown wound healing properties (Camciuc *et al.* 1998, Chanchal *et al.* 2018). Okra fruit mucilage has shown its suitability as potential alternative for blood plasma and stabilizer for foams and suspensions (Camciuc *et al.* 1998, Chanchal *et al.* 2018). Okra is also known for being identified with high level of antioxidants (Chanchal *et al.* 2018).

Okra as textile fiber

Okra fiber is extracted from the plant waste by retting process. It is a multicellular lignocellulosic fiber like jute. (Alam and Khan, 2007; Chanchal *et al.* 2018; Sen *et al.* 1987; Sen *et al.* 1988). It contains 60-70% α -cellulose, 15-30% hemicellulose, 5-14% lignin, which resembles to other traditional bast fibers (Alam and Khan, 2007). Its physical characteristics are quite comparable with jute fiber (Sen *et al.* 1987).

Okra fiber can be white, light cream or yellow in color. It has good mechanical strength (Fathima and Balasubramanian, 2006). It has immense potential for textile applications in the form of fiber, yarn and fabric such as rope, twine, fishing net, sacking and decorative fabric, carpets, mats, handicraft, paper laminates and fibre reinforced composites (Agrawal *et al.* 2004; Fortunati *et al.* 2013; Isa *et al.* 2014; Khan *et al.* 2009; Khan *et al.* 2014; Potluri *et al.* 2017; Sen *et al.* 1988; Sule *et al.* 2014). Utilization of okra fiber is three decades old, but major research work has started after 2007 (Khan *et al.* 2017; Srinivasababu 2015). In India, the scope for okra fiber is enormous due to high amount of cultivation and availability of huge harvest waste plant is a potential fiber source.

Application of okra fiber in apparels is a challenging area due to its coarseness. It is three times coarser than jute (Rai *et al.* 2012). Coarse yarns can be produced from these fibers when processed in jute machinery along with jute (Sen *et al.* 1988). Studies have shown that okra fiber has good mechanical strength and can be a promising reinforcement material for polymer composites in commercial applications (Fortunati *et al.* 2013; Isa *et al.* 2014; Kalia *et al.* 2011; Khan *et al.* 2014; Moniruzzaman *et al.* 2009; Rosa *et al.* 2010; Srinivasababu *et al.* 2009; Srinivasababu *et al.* 2009; Sule *et al.* 2014). Cellulose nanocrystals (CNC) were extracted from okra fibers. The results have shown some potential for okra fibers for their application in the form of CNC in nanocomposite systems also (Fortunati *et al.* 2013). Diversified products can be made using okra fiber such as cleaning brush, shoe brush, painting

brush and rope (Gogoi *et al.* 2017; Reddy and Reddy, 2016). Nonwovens obtained from okra fiber showed good mechanical properties and thermal conductivity values similar to the traditional fibers (both natural and synthetic) used in industry (Duman *et al.* 2017).

Okra for the paper pulp

1st report about the use of okra fiber in the paper industry was in 1982 (Fathima and Balasubramanian, 2006; Martin 1982; Sen *et al.* 1988). It was used for commercial paper in the Southern USA (Camciuc *et al.* 1998). The 'Mobile Register Journal' was printed on okra paper in 1870 (Camciuc *et al.* 1998). Okra fibers were also used for making handmade papers (Gogoi *et al.* 2017). Okra fiber was also used in paper fiber composites (Agrawal *et al.* 2004). The paper laminates prepared from okra fiber showed better tearing and bending resistance than the laminates made from pure paper sheets. The developed paper fiber laminates can be a potential candidate for the use as a low-cost alternative to high-grade corrugated paper laminates in packaging industry (Agrawal *et al.* 2004).

Okra for purification

The roots and stems of okra are used for clarification of sugarcane juice from which gur or brown sugar is prepared (Ministry of Environment and Forest 2011). The seed powder has been used as a substitute for aluminum salts in water purification (Camciuc *et al.* 1998; Ministry of Environment and Forest 2011). Okra mucilage was found as good coagulant for industrial wastewater treatment (Freitas *et al.* 2015).

Biomass and fuel

At the end of okra plant life cycle, when all the okra fruits are removed from the plant, the remaining green plant can still be cut and the stem can be used as biomass in a power station for the generation of electricity. The thoroughly dried okra stems are also burned as an inexpensive cooking fuel (Martin 1982).

CONCLUSION

Okra is an underutilized multiple-purpose crop. Other than vegetable it is used as coffee substitute, oil and protein source, biomass, fuel, paper pulp production, water purification, medicinal and textile fiber usage. Production of okra can contribute to the sustainable development in the agricultural field which in turn create dual benefit from single crop while preventing production of greenhouse gases during burning (commonly practiced). India is the highest producer of Okra in the world.

REFERENCES

Agrawal, A.K., Behera, B.K. and Abbas, J. (2004). Development of reinforced paper laminates using bhindi fibers. *Indian Journal of Fiber and Textile Research*. 29: 49-56.
 Alam, M.S. and Khan, G.A. (2007). Chemical analysis of okra bast fiber (*Abelmoschus esculentus*) and its physico-

chemical properties. *Journal of Textile and Apparel Technology and Management*. 5:1-9.
 Camciuc, M., Deplagne, M., Vilarem, G. and Gaset, A. (1998). Okra-*Abelmoschus esculentus* L. (Moench.) a crop with economic potential for set aside acreage in France. *Industrial Crops and Products*. 7: 257-264.
 Chanchal, D.K., Alok, S., Kumar, M., Bijauliya, R.K., Rashi, S. and Gupta, S. (2018). A brief review on *Abelmoschus esculentus* linn. Okra. *International Journal of Pharmaceutical Sciences and Research*. 9: 58-66.
 Duman, M.N., Kocak, E.D., Merdan, N. and Mistik, I. (2017). Nonwoven production from agricultural okra wastes and investigation of their thermal conductivities. *IOP Conference Series: Materials Science and Engineering*. 254: 192007.
 Dungani, R., Karina, M., Sulaeman, A., Hermawan, D. and Hadiyane A. (2016). Agricultural waste fibers towards sustainability and advanced utilization: A review. *Asian Journal of Plant Science*. 15: 42-55.
 Fathima, M. and Balasubramanian, A. (2006). Effect of plant growth regulators on the quality of bast fibres in *Abelmoschus esculentus* (Linn.) Moench. *Acta Botanica Croatica*. 65: 101-112.
 Fortunati, E., Puglia, D., Monti, M., Santulli, C., Maniruzzaman, M. and Kenny, J. M. (2013). Cellulose nanocrystals extracted from okra fibers in PVA nanocomposites. *Journal of Applied Polymer Science*. 128: 3220-3230.
 Fortunati, E., Puglia, D., Monti, M., Santulli, C., Maniruzzaman, M., Foresti, M.L., Vazquez, A. and Kenny, J.M. (2013). Okra (*Abelmoschus esculentus*) fibre based PLA composites: mechanical behaviour and biodegradation. *Journal of Polymers and the Environment*. 21: 726-737.
 Freitas, T.K.F.S., Oliveira, V.M., De Souza, M.T.F., Geraldino, H.C.L., Almeida, V.C., Fávoro, S.L. and Garcia, J.C. (2015). Optimization of coagulation-flocculation process for treatment of industrial textile wastewater using okra (*A. esculentus*) mucilage as natural coagulant. *Industrial Crops and Products*. 76: 538-544.
 Gogoi, N., Gogoi, M. and Choudhury, S. (2017). Utilization of agro waste-okra and its potentiality. *Asian Journal of Home Science*. 12: 250-256.
 Isa, M.T., Usman, S., Ameh, A.O., Ajayi, O.A., Omorogbe, O. and Ameuru, S.U. (2014). The effect of fiber treatment on the mechanical and water absorption properties of short okra/glass fibers hybridized epoxy composites. *International Journal of Materials Engineering*. 4: 180-184.
 Jain, N., Jain, R., Jain, V. and Jain, S. (2012). A review on: *Abelmoschus esculentus*. *Pharmacia*. 1: 84-89.
 Joh, A., Anyebe, S.B. and Kwaya, P.V. (2015). Effect of processing methods on selected physico chemical properties of Okra. *International Journal of Engineering Technology and Computer Research*. 3: 118-125.
 Kalia, S., Kaith, B.S. and Kaur, I. eds. (2011). Cellulose fibers: bio-and nano-polymer composites: Green Chemistry and Technology. Springer Science and Business Media.
 Kambli, N.D., Samanta, K.K., Basak, S., Chattopadhyay, S.K., Patil, P.G. and Deshmukh, R.R. (2018). Characterization of the corn husk fibre and improvement in its thermal stability by banana pseudostem sap. *Cellulose*. 25: 5241-5257.

- Khan, G.M.A., Abedin, S.M.A., Choudhury, M.J., Gafur, M.A. and Alam, M.S. (2014). Renewable okra bast fiber reinforced phenol formaldehyde resin composites: mechanical and thermal studies. *Research and Reviews: Journal of Material Sciences*. 2: 32-36.
- Khan, G.M.A., Shaheeruzzaman, M., Rahman, M.H., Razzaque, S.M.A., Islam, Md. S. and Alam, Md. S. (2009). Surface modification of okra bast fiber and its physico-chemical characteristics. *Fibers and Polymers*. 10: 65-70.
- Khan, G.M.A., Yilmaz, N.D. and Yilmaz, K. (2017). Okra fibers: Potential material for green biocomposites." In: *Green Biocomposites*, 261-284. Springer, Cham.
- Kocak, E.D., Merdan, N., Mistik, I. and Sahinbaskan, B.Y. (2018). Fiber extraction from okra plant agricultural wastes, their characterizations and surface modifications by environmental methods. In: *Sustainable Innovations in Textile Fibres*, 53-80. Singapore: Springer.
- Makhadmeh, I.M. and Ereifej, K.I. (2004). Geometric characteristics and chemical composition of okra (*Hibiscus esculentus* L.) grown under semi-arid conditions. *International Journal of Food Properties*. 7: 83-90.
- Martin, F.W. (1982). Okra, potential multiple-purpose crop for the temperate zones and tropics. *Economic Botany*. 36: 340-345.
- Ministry of Agriculture and Farmers Welfare. (2017). Horticultural statistics at a glance 2017. <http://www.indiaenvironmentportal.org.in/content/450238/horticultural-statistics-at-a-glance-2017/>
- Ministry of Environment and Forest. (2011). *Biology of Abelmoschus esculentus* L. (Okra). Accessed July 10, 2018. [http://envfor.nic.in/divisions/csuvr/geac/Biology_of_Okra\[1\].pdf](http://envfor.nic.in/divisions/csuvr/geac/Biology_of_Okra[1].pdf)
- Moniruzzaman, M., Maniruzzaman, M., Gafur, M.A. and Santulli, C. (2009). Lady's finger fibres for possible use as a reinforcement in composite materials. *Journal of Biobased Materials and Bioenergy*. 3: 286-290.
- Moosavi, S.A., Aghaalikhani, M., Ghobadian, B. and Fayyazi, E. (2018). Okra: A potential future bioenergy crop in Iran. *Renewable and Sustainable Energy Reviews*. 93: 517-524.
- Osawaru, M.E., Ogwu, M.C. and Emokpare, A.A. (2014). Preliminary assessment of the microanatomy of okra [*Abelmoschus* (L.) Wood. *Egyptian Academic Journal of Biological Sciences*. 5: 39-54.
- Potluri, R., Paul, K.J. and Prasanthi, P. (2017). Mechanical properties characterization of okra fiber based green composites and hybrid laminates. *Materials Today: Proceedings*. 4: 2893-2902.
- Rai, S., Hossain, F. and Hossain, M. (2011). Studies on the combining ability and heterosis in okra [*Abelmoschus esculentus* (Moench) L.] for bast fibre yield. *Journal of Crop and Weed*. 7: 64-66.
- Rai, S., Hossain, M. and Hossain F. (2012). Evaluation of okra [*Abelmoschus esculentus* (Moench) L.] as bast fibre crop. *Journal of Crop and Weed*. 8: 101-104.
- Reddy, N. and Yang, Y. (2005). Biofibers from agricultural by products for industrial applications. *Trends in Biotechnology*. 23: 22-27.
- Reddy, S.P. and Reddy, A.C. (2016). Tensile and flexural strength of okra fiber reinforced polymer composites. *International Journal of Engineering and Management Research*. 6: 491-495.
- Rosa, D., Maria, I., Kenny, J.M., Puglia, D., Santulli, C. and Sarasini, F. (2010). Morphological, thermal and mechanical characterization of okra (*Abelmoschus esculentus*) fibres as potential reinforcement in polymer composites. *Composites Science and Technology*. 70: 116-122.
- Roy, A., Shrivastava, S.L. and Mandal, S.M. (2014). Functional properties of Okra *Abelmoschus esculentus* L. (Moench): traditional claims and scientific evidences. *Plant Science Today*. 1: 121-130. <http://dx.doi.org/10.14719/pst.2014.1.3.63>.
- Santulli, C., Sarasini, F., Fortunati, E., Puglia, D. and Kenny, J.M. (2014). Okra fibres as potential reinforcement in biocomposites. In: *Biomass and Bioenergy*. 175-190. Springer, Cham.
- Sarasini, F. and Fiore, V. (2018). A systematic literature review on less common natural fibres and their biocomposites. *Journal of Cleaner Production*. doi:10.1016/j.jclepro.2018.05.197.
- Sathish, D. and Eswar, A. (2013). A review on: *Abelmoschus esculentus* (Okra). *International Research Journal of Pharmaceutical and Applied Sciences*. 3: 129-132.
- Sen, K.K., Saha, S.C. and Ray, P.K. (1987). Physical and structural properties of raw and alkali-treated Bhindi (*Hibiscus esculentus* L) fibre. *Indian Journal of Textile Research*. 12: 152-153.
- Sen, K.K., Saha, S.C., Mojumder, P. and Ray, P.K. (1988). Some physical properties of alkali treated bhindi fiber (*Hibiscus esculentus* linn.). *Journal of Applied Polymer Science*. 36: 1601-1605.
- Sharma, R.K. and Prasad, K. (2010). Characterisation of promising okra genotypes on the basis of Principal Component Analysis. *The Society for Advancement of Horticulture*. 12: 71-74.
- Srinivasababu, N. (2015). An overview of okra fibre reinforced polymer composites. In: *IOP Conference Series: Materials Science and Engineering*. 83: 012003. doi:10.1088/1757-899X/83/1/012003.
- Srinivasababu, N. Rao, K.M.M. and Kumar, J.S. (2009). Experimental determination of tensile properties of okra, sisal and banana fiber reinforced polyester composites. *Indian Journal of Science and Technology*. 2: 35-38.
- Srinivasababu, N., Rao, K.M.M. and Kumar, J.S. (2009). Tensile properties characterization of okra woven fiber reinforced polyester composites. *International Journal of Engineering*. 3: 403-412.
- Sule, U., Isa, M.T., Ameh, A.O., Ajayi, O.A. and Omorogbe, O. (2014). Studies on the properties of short Okra/Glass fibers reinforced epoxy hybrid composites. *The International Journal of Science and Technoledge*. 2: 260-265.
- Ullah, S., Kilpeläinen, P., Ilvesniemi, H., Pakkanen, H. and Alén, R. (2018). Chemical characterization of Okra stalk (*Abelmoschus esculentus*) as potential raw material for biorefinery utilization. *Cellulose Chemistry and Technology*. 52.
- Varmudy, V. (2011). Need to boost okra exports. *Facts for You*. 31: 21-23.