



Influence of Vegetable and Fruit Wastes Vermicompost on the Growth and Yield of Black Gram (*Vigna mungo* L.)

Pinky Raihing, A. Vijayalakshmi

10.18805/ag.D-5492

ABSTRACT

Background: Composting is the process of breaking down of organic matter using living organisms which can be used as manure for soil conditioning. Vermicomposting is the rapid decomposition of biowastes using earthworm species which will enhance the growth and yield of plants.

Methods: The vermicomposting of vegetable and fruit wastes was carried out during summer season (May-July) and by using the vermicompost a pot culture experiment was conducted in black gram for three months from August to October, 2019. The treatments consist of combinations of vermicomposted wastes and cowdung along with bioinoculants, Soil served as control. The biometric parameters such as root length, shoot length, fresh weight and dry weight on 15, 35, 55 DAS and yield characters like number of pods/plants, number of seeds/pods, pod length, weight of seed/ pod, pod fresh weight and pod dry weight were recorded and statistically analysed. The present study has been attempted to convert the vegetables and fruit waste into compost as manure and analyse its effect on vegetative growth and yield parameters of Black gram (*Vigna mungo* L.).

Result: The highest value of shoot length, root length, fresh weight and dry weight were noted in fruit waste + cowdung + *Pleurotus eous* (APK1) + *Trichoderma asperelloides* + *Eudrilus eugeniae* on 15, 35 and 55 day after sowing (DAS) and the least was reported in control (only soil). The application of T₈ treatment reported the highest yield characters which were followed by other treatments and control on 15, 35 and 55 DAS. The study concluded that T₈ (F.W + cow dung + *P. eous* + *T. asperelloides* + *Eudrilus eugeniae*) is an effective biocompost for the growth and yield of black gram (*Vigna mungo* L.) which is eco-friendly.

Key words: Black gram, Fruit waste, *Eudrilus eugeniae*, Vermicomposting, Vegetable waste.

INTRODUCTION

Agriculture plays a vital role in Indian economy as most of the population, from rural areas depend on it for their livelihood. Organic farming can address many of these problems, as this system helps to maintain soil productivity and effectively control pest by enhancing natural processes and cycles in harmony with environment (Barik, 2017). Organic waste and the biofertilizers are the alternate sources that meet the nutrient requirement on crops (Faheed and Fattah, 2008). India produces about 3000 million tons of organic waste annually, which can be degraded and can produce eco-friendly manure for sustainable environment (Achsah and Lakshmi, 2013). As disposal of organic waste are important for healthy environment, environment friendly management and disposal of wastes has become global priority (Sakthivigneswari and Vijayalakshmi, 2017). Vermicomposting is a cheap and low-cost technological process of composting where species of earthworm is introduced to produce a better end product. Vermicompost application promoted the plant growth and also protects the crops from the pest and diseases (Sinha *et al.* 2010). Black gram is a bean that grows mainly in south Asia and it is an annual herb. The plant is erect and hairy. It is healthy pulse which is rich in vitamins, minerals. It is also the store-house to many nutrients especially calcium, potassium, iron, magnesium, copper, manganese. The seed of black gram is used for cooling and astringent properties. The aim of the experiment was biocomposting of vegetable and fruit wastes

Department of Botany, Avinashilingam Institute for Home Science and Higher Education for Women, Coimbatore-641 043, Tamil Nadu, India.

Corresponding Author: Pinky Raihing, Avinashilingam Institute for Home Science and Higher Education for Women, Coimbatore-641 043, Tamil Nadu, India.

Email: raihingpinky17@gmail.com

How to cite this article: Raihing, P. and Vijayalakshmi, A. (2022). Influence of Vegetable and Fruit Wastes Vermicompost on the Growth and Yield of Black Gram (*Vigna mungo* L.). Agricultural Science Digest. DOI: 10.18805/ag.D-5492.

Submitted: 22-09-2021 **Accepted:** 26-11-2021 **Online:** 11-01-2022

into biomanure and to study the effect on growth and yield of black gram (*Vigna mungo* L.).

MATERIALS AND METHODS

The composting and vermicomposting were carried out during summer season (May-July) and the pot culture experiment was conducted for three months from August to October during 2019 at Avinashilingam Institute for Home Science and Higher Education for Women, Coimbatore, Tamil Nadu (11.0196°N, 76.9504°E).

Collection of waste

The vegetable wastes were collected from local vegetables market situated near the Institute. The fruit wastes were

collected from nearby fruit sellers. Thus, collected wastes were cut into small pieces, sun dried and stored in polythene bags.

Assembling of seed

The seeds of black gram (*Vigna mungo* L.) were obtained from Tamil Nadu Agriculture University, Coimbatore, Tamil Nadu and were subjected to pot culture studies with three replications.

Preparation of compost pit

Eight compost pits were dug with a dimension of 1.5 feet length and 4 square feet width. The compost pit were designated as C₁- C₈ (C denotes as compost).

Collection of earthworm

The earthworm species African earthworm (*Eudrilus eugeniae*) was collected from ICAR KVK (Krishi Vigyan Kendra) Coimbatore, Tamil Nadu.

Preparation of vermicompost

About 60×30×15 cm size plastic tray was purchased from the market and small holes were made at the bottom of the trays to facilitate draining of excess water. Vermibeds were prepared by combination of pre-decomposed vegetable and fruit wastes and cow dung in 2:1 ratio. (2.0 kg of pre-decomposed waste and 1 kg of dried cow dung in the trays).

The experimental treatments

A pot culture experiment was carried out with sandy loam clay soil and vermicompost prepared from vegetable and fruit waste. Initially, the pebbles present in the soil were removed and 7.0 kg of sandy loam clay soil was filled in each pot. Ten seeds were sown and among the germinated ones, five healthy plants were maintained in pot for the study. The different treatments for pot culture are given below.

Treatments

C	Control (Only soil)
T ₁	Vegetable wastes + Cow dung + <i>Eudrilus eugeniae</i> (5 t/ha).
T ₂	Vegetable wastes + Cow dung + <i>Pleurotus eous</i> (APK1) + <i>Eudrilus eugeniae</i> (5 t/ha).
T ₃	Vegetable wastes + Cow dung + <i>Trichoderma asperelloides</i> + <i>Eudrilus eugeniae</i> (5 t/ha).
T ₄	Vegetable wastes + Cow dung + <i>P. eous</i> (APK1) + <i>T. asperelloides</i> + <i>Eudrilus eugeniae</i> (5 t/ha).
T ₅	Fruit wastes + Cow dung + <i>Eudrilus eugeniae</i> (5 t/ha).
T ₆	Fruit wastes + Cow dung + <i>Pleurotus eous</i> (APK1) + <i>Eudrilus eugeniae</i> (5 t/ha).
T ₇	Fruit wastes + Cow dung + <i>Trichoderma asperelloides</i> + <i>Eudrilus eugeniae</i> (5 t/ha).
T ₈	Fruit wastes + Cow dung + <i>P. eous</i> (APK1) + <i>T. asperelloides</i> + <i>Eudrilus eugeniae</i> (5 t/ha).

Statistical analysis

The biometric data obtained on 15, 35, 55 DAS and 75 DAS were analyzed statistically using one way or two-way anova.

RESULTS AND DISCUSSION

Biometric character

The present study results showed a positive effect of compost treatments on the test crop and significantly maximum increase was observed in T₈ treated plants during 15, 35 and 55 DAS.

Shoot length and root length (Table 1)

A significant increase in shoot length was observed T₈ (56.9, 62.1 and 71.5 cm) followed by T₄ (44.1, 56.2 and 62.5 cm) on 15, 35 and 55 DAS when compared to Control (21.8, 29.7 and 38.5 cm). The highest root length was reported in T₈ (14.5, 17.2 and 18.5 cm) treatment followed by T₄ (14.1, 16.3 and 16.8 cm) on 15, 35 and 55 DAS.

The results were in agreement with the finding of Badar and Qureshi, (2014), who reported that increased root length of 28.0 cm on 30th day and 29.66 cm on 60th day in sunflower plants in *Trichoderma hamatum* inoculated rice husk (5 kg) as compared to the control.

The finding is in conformity with Kamble *et al.* (2016) that, application of T₃- 100% NPK+ poultry manure @ 5 t ha⁻¹, recorded maximum plant height (37.30 cm), and yield attributing characters such as pod length (12.83 cm), pod diameter (0.93 cm), number of pods plant⁻¹ (42.18). The lowest value of growth characters was recorded in treatment T₁₆ - FYM @ 25 t ha⁻¹ in French bean.

According to Coulibaly *et al.* (2019) the tallest plant (70.19 cm) was measured with the compost from pig waste, followed by the chicken compost (51.4 cm) and the shortest was noted with the control (22.45 cm) on 45 days in maize.

The results in par with Sutar *et al.* (2019) who recorded that application of jeevamrutha @1000l ha⁻¹ has higher plant height (65.60 cm), number of branches (8.89), number of leaves (26.50), leaf area (1039.56 cm²) and yield attributing characters like number of pods per plant (20.57), pod length (19.8 cm), number of seeds per pod (15.58), seed weight per plant (9.53 g) and test weight (11.82 g) when compared to control in cowpea.

Similar results were observed by Pinky and Vijayalakshmi, (2020), who noted significant increase in shoot length and root length in T₄ treatment - F. W + cow dung + *P. eous* + *T. asperelloides* + *Eudrilus eugeniae* (5 t/ha) where C- Control (plain soil) on 15, 35 and 55 DAS in black gram.

The results coincide with Silpa and Vijayalakshmi, (2020) that, shoot length showed significant increase in T₄ (67.933 cm, 112.133 cm and 119.300 cm) on 25, 35 and 45 DAS when compared with the control (13.933 cm, 35.233 cm and 50.233 cm) and the root length is noted maximum in T₄ (18.133 cm, 26.867 cm and 30.100 cm) when compared to control (5.7 cm, 10.767 cm and 14.733 cm) on 25, 35 and 45 DAS in cowpea.

Fresh weight and dry weight

A significant increase in fresh weight content (Table 2) was observed in T₈ (1.910, 1.967 and 2.063 gm) treatment

followed by other treatment and control (0.405, 0.700 and 0.821 gm) treatment on 15, 35 and 55 DAS. A maximum dry weight content was noted in T_8 (0.465, 0.492 and 0.665 gm) treatment which was followed by T_4 (0.390, 0.406 and 0.629 gm).

The present finding coincides with the result of Mehdizadeh *et al.* (2013) who reported that the application of municipal waste compost (20 t/ha) significantly enhanced the fresh weight (shoots-195.2 gm, roots-62.3 gm) and dry weight (shoots-51.2 gm, roots-13.5 gm) of tomato crop.

Furthermore, the present finding was in conformity with that of Sivakumar and Karthikeyan, (2016), who concluded that the increase in fresh weight (6.6 g) and dry weight (2.2 g) was due to the application of vermicomposted weed plants waste using *Eudrilus eugeniae* in brinjal plant.

The results are in par with Sharma *et al.* (2019) who observed the fresh and dry weight of plants were maximum in T_7 (3.44 and 1.40 g) treatment followed by T_6 (2.43 and 0.35 g) and T_1 (1.76 and 0.15 g) in *Spinacia oleracea*.

Similar results were observed by Silpa and Vijayalakshmi, (2020) who noted a significant increase in fresh weight content in T_4 (7.195 g, 8.186 g and 12.555 g) when compared to control (1.053 g, 1.077 g and 1.856 g) and highest dry weight was registered in T_4 treatment

(1.277g, 1.465 g, and 2.254 g) over control (0.494 g, 0.604 g and 0.819 g) on 25, 35 and 45 DAS in cowpea.

The study reported by Al-Sabbagh *et al.* (2020) stated the maximum fresh weight of Chinese kale in Market compost (0.276 gm) grown plants followed by Ecodrum composter (0.271 gm), when compared to control (0.133 gm) respectively.

Yield parameters (Fig 1)

The maximum number of pods/plants was observed in T_8 (28.00) treatment followed by T_4 (24.00) when compared to control (8.00). The pod length and number of seeds/pods were significantly increased in T_8 (4.30 cm and 7.00) treatment followed by T_4 (4.00 cm and 6.00) treatment and control. The fresh and dry weight of the pods were increased significantly in T_8 (0.795 gm and 0.502 gm) followed by other treatment and control (0.410 gm and 0.215 gm). The weight of the seed showed maximum in T_8 (0.372 gm) and followed by T_4 (0.357 gm) treatment when compared to other treatment and control (0.139 gm).

A Similar work was done by Hyder *et al.* (2015) who confirmed highest number of fruit (97) and diameter of the fruit (6.75 cm) in tomato plant due to the application of vermicompost (2 t/ha).

Table 1: Vegetative parameters of black gram influence by vegetable and fruit wastes vermicompost.

Treatment	Shoot length (cm)			Root length (cm)		
	15	35	55	15	35	55
C	21.8	29.7	38.5	4.6	6.2	6.9
T_1	28.9	30.0	43.2	6.5	6.7	7.3
T_2	30.2	32.1	50.3	7.5	10.3	12.3
T_3	36.9	44.5	48.0	9.3	14.5	14.9
T_4	44.1	56.2	62.5	14.1	16.3	16.8
T_5	31.5	34.2	46.2	8.0	9.2	9.8
T_6	31.8	46.4	55.4	8.2	10.5	14.2
T_7	40.2	48.0	64.2	11.2	15.2	16.3
T_8	56.9	62.1	71.5	14.5	17.2	18.5
SEd		2.96265			0.32923	
Cd (p<0.05)		5.93977			0.66008	

Table 2: Effect of vegetable and fruit wastes biocompost on vegetative parameters of black gram.

Treatment	Fresh weight (gm)			Dry weight (gm)		
	15	35	55	15	35	55
C	0.405	0.700	0.821	0.075	0.092	0.117
T_1	0.496	0.736	0.974	0.106	0.146	0.286
T_2	0.643	1.144	1.341	0.139	0.270	0.390
T_3	0.955	1.224	1.580	0.287	0.314	0.456
T_4	1.241	1.694	1.885	0.390	0.406	0.629
T_5	0.736	1.049	1.168	0.164	0.230	0.295
T_6	0.964	1.332	1.653	0.235	0.315	0.436
T_7	0.972	1.569	1.792	0.287	0.394	0.467
T_8	1.910	1.967	2.063	0.465	0.492	0.665
SEd		0.03366			0.03285	
Cd(p<0.05)		0.06749			0.06585	

The results are in conformity with Joshi *et al.* (2016), that application of T_2 - RDF (20-40-0 NPK kg ha⁻¹) recorded significantly higher number of pods plant⁻¹ (79.60) and number of seeds pod⁻¹ (13.45) in cowpea.

The results coincide with Sajjan *et al.* (2020) who reported significant increase in seed yield such as number of pods per plant (117.19), number of seeds per pod (4.25) and pod length (5.66 cm) was reported in *rabi* season crop in dolichos bean [*Lablab purpureus* (L.) Sweet].

The results are in accordance with Ropo *et al.* (2020) who noted that poultry manure have the highest number of fruits per plant (8.75), heaviest fruit per plant (25.30 g), highest fruit length (6.60 cm) and fruit diameter (1.93 cm), which were significantly different from all other treatments. The results coincide with Silpa and Vijayalakshmi, (2021) who observed that number of pods/plant (21), length of pod (16.50 cm), number of seeds/pod (20), weight of the seed/pod (1.68 g), pod fresh weight (5.711 g) and pod dry weight




Fig 1: Yield attributing parameters of Black gram (*Vigna mungo* L.).

a) Number of pods/plants b) No. of seeds/pod c) Pod length (cm) d) Weight of seed (g) e) Pod fresh weight (g) and Pod dry weight (g).

(2.398 g) in T₈ - Raw Jack fruit peel + 10 g *Pleurotus eous* + 10 g *Pleurotus florida* + *Eudrilus eugeniae* 5t/ha⁻¹ treatment when compared to control in *Vigna unguiculata* (L.) Walp.

CONCLUSION

Vermicompost contains macro and micro nutrients that help in growth and development of the plant. The present study indicated that application of the treatment (T₈ Vegetable waste + cow dung + *P. eous* (APK1) + *T. asperelloides* + *Eudrilus eugeniae*) showed highest growth and yield/plant in black gram. This composting can be done in large scale since it is cost effective and eco-friendly which is not harmful to the environment.

ACKNOWLEDGEMENT

We would like to thank Avinashilingam Institute for Home Science and Higher Education for Women and Ministry of Tribal Affairs for providing facilities and fund to carry out the study.

REFERENCE

- Achsaah, R.S. and Prabha, M.L. (2013). Potential of vermicompost produced from banana waste (*Musa paradisiaca*) on the growth parameters of *Solanum lycopersicum*. International Journal of Chemtech Research. 5: 2141-2153.
- Al-Sabbagh, T., Tareq Madouh, Craig, A.M. and Sugumaran, K. (2020). Influence of dead sheep compost material using aerobic technique on the growth of leafy vegetables in kuwait under greenhouse conditions. Journal of Agriculture and Horticulture Research. 3(2): 31-37.
- Badar, R. and Qureshi, S.A. (2014). Composted rice husk improves the growth and biochemical parameters of sunflower plants. Journal of Botany. Article ID 427648, 6 pages.
- Barik, A.K. (2017). Organic farming in India: Present status, challenges and technological breakthrough. In 3rd Conference on Bio-resource and Stress Management International. (pp. 101-110).
- Coulibaly, S.S., Kouassi, K.I., Koffi, K.K. and Zoro, B.I.A. (2019). Effect of compost from different animal manures on maize (*Zea mays*) growth. Journal of Experimental Biology and Agricultural Sciences. 7(2): 178-185.
- Faheed, F.A. and Fattah, Z.A. (2008). Effect of chlorella vulgaris as biofertilizers on growth parameters and metabolic aspects of lettuce plant. Journal of Agriculture and Social Science. 4: 165-169.
- Hyder, S.I., Farooq, M., Sultan, T., Ali, A., Ali, M., Kiani, M.Z., Ahmad, S. and Tabssam, T. (2015). Optimizing yield and nutrients content in tomato by vermicompost application under greenhouse conditions. Natural Resources. 6: 457-464.
- Joshi, D., Gediya, K.M., Patel, J.S., Birari, M.M. and Gupta, S. (2016). Effect of organic manures on growth and yield of summer cowpea [*Vigna unguiculata* (L.) Walp] under middle Gujarat conditions. Agricultural Science Digest-A Research Journal. 36(2): 134-137.
- Kamble, M.Y., Kalalbandi, B.M., Kadam, A.R. and Rohidas, S.B. (2016). Effect of organic and inorganic fertilizers on growth, green pod yield and economics of french bean (*Phaseolus vulgaris* L.) cv. HPR-35. Legume Research-An International Journal. 39(1): 110-113.
- Mehdizadeh, M., Darbandi, E.I., Naseri-Rad, H. and Tobeh, A. (2013). Growth and yield of tomato (*Lycopersicon esculentum* Mill.) as influenced by different organic fertilizers. International Journal of Agronomy and Plant Production. 4(4): 734-738.
- Pinky Raihing and Vijayalakshmi, A. (2020). Biocomposting of fruit wastes into manure and its effects on the growth, chlorophyll content and yield parameters of black gram (*Vigna mungo* L.). Proteus Journal. 11(10): 70-83.
- Ropo, A.D., Ojochegbe, Y.D., Amina, I.V. and Ugbojo-Ide, O.J. (2020). Effects of organic and inorganic fertilizers on agromorphological traits of tomato [*Solanum lycopersicum* (L.) Moench]. GSC Biological and Pharmaceutical Sciences. 11(2): 314-319.
- Sajjan, A.S., Malabasari, T.A. and Shashidhar, T.R. (2020). Seed and yield parameters as influenced by season and plant growth regulators in dolichos bean [*Lablab purpureus* (L.) Sweet]. Legume Research: An International Journal, 43(6): 856-860.
- Sakthivigneswari, G. and Vijayalakshmi, A. (2017). Biocomposting technology and microbial populations of agroindustrial waste using *pleurotus sajor-caju* and earthworm (*eudrilus eugeniae*). International Journal of Pharma and Bio Sciences. 8(2): 679-685.
- Sharma, A., Chetani, R., Batra, N.G., Saxena, A. and Trivedi, P. (2019). Effect of Organic and Inorganic Fertilizers on the Performance of (*Spinacia oleracea* L.) Journal of Phytological Research. 32(1 and 2): 17-22.
- Silpa, M. and Vijayalakshmi, A. (2020). Effect of cocoa shell biocompost to improve the growth parameters of cowpea [*Vigna unguiculata* (L.) Walp]. International Journal of Current Advanced Research. 09(01): 21114-21118.
- Silpa, M. and Vijayalakshmi, A. (2021). Efficiency of biocomposted agroindustrial wastes and their response in the growth and yield of [*Vigna unguiculata* (L.) Walp]. Agricultural Science Digest - A Research Journal. doi:10.18805/ag.d-5321.
- Sinha, J., Biswas, C.K., Ghosh, A. Saha, A. (2010). Efficacy of Vermicompost against fertilizers on *Cicer* and *Pisum* and on population diversity of N₂ fixing bacteria. Journal of Environmental Biology. 31: 287-292.
- Sivakumar, P. and Karthikeyan, R. (2016). Bioconversion of vermicomposted weed plants waste using *Eudrilus eugeniae* Kinberg for promoting the growth of brinjal (*Solanum melongena*). International Journal of Development Research. 6: 8020-8023.
- Sutar, R., Sujith, G.M. and Devakumar, N. (2019). Growth and yield of Cowpea [*Vigna unguiculata* (L.) Walp] as influenced by jeevamrutha and panchagavya application. Legume Research-An International Journal. 42(6): 824-828.