



Effect of Breed, Season and Stage of Lactation on Different Milk Parameters at Organized Farm

G. Mishra, S.C. Goswami, A.K. Jhirwal, S. Paliwal

10.18805/ajdfr.DR-1937

ABSTRACT

Background: The current study had been conducted to understand the impact of breed as well as season and stage of lactation on various milk constituents viz. freezing point, SNF, pH, lactose, protein, fat and electric conductivity of Sahiwal and Kankrej cows maintained at LRS "Livestock Research Station", College of Veterinary and Animal Science, RAJUVAS, Bikaner, Rajasthan.

Methods: Milk samples from 45 each Sahiwal and Kankrej cows were collected and analyzed for milk composition parameters. Data have been classified as per breed along with season and stage of lactation.

Result: The analysis of variance for breed demonstrated highly significant impacts of fat and SNF whereas non-significant effect was found on pH, freezing point, lactose and protein. The analysis of variance demonstrated highly significant ($P < 0.01$) impact of season on fat, SNF, freezing point and significant effect on protein whereas non-significant effect on lactose, pH and Electric conductivity in Kankrej cattle however highly significant ($P < 0.01$) effect was found on fat in Sahiwal cattle. The analysis of variance presented significant ($P < 0.05$) effect of lactation stage on fat and Protein in Sahiwal cattle whereas on fat and SNF in Kankrej cattle.

Key words: Breed, Kankrej, Milk composition, Sahiwal, Season, Stage of lactation.

INTRODUCTION

The nutritional composition of cow milk comprises 3.5% protein, 4.5% fat, 8.5% solid not fat, 4.6% lactose and freezing point -0.55°C . Study on milk composition along with functional properties of the indigenous cow milk are of the considerable importance to milk consumers, manufacturers as well as dairy farmers. Dairy farmers, manufacturers and consumers all depend on milk components and composition for ensuring the quality of their raw milk along with milk products and overall health and well-being (Reis *et al.*, 2013).

These properties of milk might be impacted by various factors including: seasonal variation (Bernabucci *et al.*, 2002), storage of milk (Hanna *et al.*, 2004), nutritional level, breeds of animals, parity of cows, milk somatic cell (Ramos *et al.*, 2015; Singh, 2019) lactation stage as well as farm's geographic location (Alyaqoubi *et al.*, 2014).

Breed is a genetic factor which affects the composition, quantity and quality of milk significantly. Milk characteristics may show changes in a year as a result of the influence of the seasons on animals. Among the main causes of seasonal variations could be due to dietary factors, in particular the characteristics of feeds, as well as housing factors, feeding methods and climatic conditions of the region. A few conditions, like cows confined indoors and given silage in the winter, then kept on pasture as well as fed new grass in the spring and summer, have a key impact in the dietary aspects. The season could be considered the main factor that has affected the fluctuations in milk characteristics throughout as feed, climatic variables changes. According to Bernabucci *et al.* (2015), milk fat and protein concentrations are lower in summer, higher in winter; intermediate in spring milk.

Department of Livestock Production Management, College of Veterinary and Animal Sciences, Bikaner-334 001, Rajasthan, India.

Corresponding Author: G. Mishra, Subject Matter Specialist (Animal Husbandry), Krishi Vigyan Kendra, Gudamalani, Agriculture University, Jodhpur-342 304, Rajasthan, India.

Email: geeteshmishra09@gmail.com

How to cite this article: Mishra, G., Goswami, S.C., Jhirwal, A.K. and Paliwal, S. (2022). Effect of Breed, Season and Stage of Lactation on Different Milk Parameters at Organized Farm. Asian Journal of Dairy and Food Research. DOI: 10.18805/ajdfr.DR-1937.

Submitted: 11-04-2022

Accepted: 23-06-2022

Online: 15-07-2022

Stage of lactation also shows variable impact upon milk constituents. During the height of lactation, high-yielding cows require more energy than they can get from their food. This is due to a decrease in hunger following parturition, as well as an increase in metabolic activities, which has the effect of necessitating a high number of substrates for the synthesis of milk ingredients (Józwik *et al.*, 2012). Fat percentages and milk protein percentages are influenced by the lactation stage in a similar way. Just after freshening, colostrum contains the maximum concentration of fat and protein in milk. During the first 25 to 50 days after calving, levels drop to their lowest point and they reach their maximum at 250 days, when milk production begins to decline (Heinrichs *et al.*, 2016). In this study we included genetic factor viz. breed and non genetic factors season and stage of lactation into consideration and analysed the effect of these factors upon different milk constituents.

MATERIALS AND METHODS

The current research was performed at Livestock Research Station, Collage of Veterinary and Animal Science, Bikaner (RAJUVAS), Rajasthan, India, during 2019-20 on Kankrej along with Sahiwal cattle that were kept in loose housing system. Every animal on farm fed with seasonal dry fodder, green fodder and compound concentrate mixture in accordance with the farm's typical feeding schedule. 45 total milk samples were collected from 45 healthy, lactating Kankrej and Sahiwal cows. Clean sampling bottles were used to collect about 50 ml of aseptic milk sample after discarding the first 2-3 streaks of foremilk. After collecting all samples, they have been immediately taken to the laboratory for additional analysis. On the same day, all milk samples were processed. According to normal protocol, all samples were tested using milk auto analyzer whereas pH and electric conductivity were measured by pH meter and EC meter. The seasons were classified viz. Summer, Winter, Rainy. In the present study, effect of breed, season and stage of lactation were taken into consideration. ANOVA (One-way analysis of variance) has been utilized for evaluating the data whereas Duncan's multiple range test has been utilized for comparing the means. In order to do any statistical analysis, SPSS software were utilized (version 20.0).

RESULTS AND DISCUSSION

Milk composition is the primary index for assessment of quality; mainly milk fat content is utilized as quality test (Zelalem, 2010). Milk's solids content determines both its nutritional and commercial value. Nutritional value increases as the solids level increases and more milk products may be produced (Pandy and Voskull, 2011). Protein content is a major criterion for evaluating the quality of milk and is used to determine payments to dairy farmers in many countries when other factors such as fat and solids content are taken into account the product's non-fat content (FAO, 2004). Breed, species, parity (Sarkar *et al.*, 2006; Mushtaq *et al.*, 2012; Singh, 2019), BCS (Mushtaq *et al.*, 2012; Singh, 2019), dietary composition, season, lameness, heat stress, behavior, locality udder health, along with lactation stage all are the factor that influence the Milk composition (Sarkar *et al.*, 2006; Kayastha *et al.*, 2008; Arora and Bhojak, 2013).

Effect of breed on milk composition

The mean \pm SE values for effect of breed on fat, protein, lactose, SNF, freezing point, pH and EC are presented in Table 1. The outcomes of the analysis of variance revealed that breed had significant impact on fat storage. According to Adesina, (2012) breed had significant effect on fat percent. Fat content in milk on a daily basis was also obtained to be significantly higher ($p < 0.05$) in synthetic cows (Ethiopian Ogaden cattle, Jersey \times Horro crosses and Holstein Friesian \times Jersey \times Horro crosses) whereas lowest in Holstein Frisian, according to a study by Kebede, 2018. Analysis of variance revealed breed's significant impact on SNF. The significant difference of SNF in our study has been in agreement with Bobbo *et al.*, (2014), Falta *et al.*, (2014) and Kedzierska-Matysek *et al.*, (2011). As per the analysis of variance, there was no significant influence of breed on protein, which is in agreement with Adesina, (2012). African cattle breeds like Bonsmara, Afrikaner, Tuli, Nguni and Boran, were analysed by Myburgh *et al.* (2012) and found non significant variation in protein content. According to Ivanov *et al.*, (2017), Bendelja *et al.*, (2011) and Pintic *et al.*, (2007), the influence of breed on milk protein content was shown to be non-significant. The analysis of variance results presented that breed lactose content has not been significantly affected by breed. The similar findings were also observed by Kebede, (2018), Adesina, (2012), Falta *et al.*, (2014) and Singh, (2019). Contradicted finding and significant differences in lactose content among different breeds reported by Myburgh *et al.*, (2012). Analysis of variance results demonstrate that breed had no effect on the freezing point. Same results were observed by Kedzierska-Matysek *et al.*, (2011) and Brzozowski and Zdziarski (2006) for freezing point. Breed had no significant impact on pH, according to the analysis of variance. As per Singh, (2019) stated non significant impacts of breed on pH among Sahiwal, Kankrej and Rathi. Contrary to our findings Bobbo *et al.*, (2014) found significant difference in the milk pH in the different cattle breeds. There is variations in electric conductivity among breeds as electrical conductivity in milk is influenced by the udder health status of the cow. Electrical conductivity of milk may be a potential trait in a breeding program. Electrical conductivity was also found a low

Table 1: Means \pm SE milk composition from Sahiwal and Kankrej breeds.

Parameters	Sahiwal	Kankrej	Level of significance
Fat (g/100 g)	3.46 \pm 0.03 ^a	4.37 \pm 0.06 ^b	**
Protein (g/100 g)	3.51 \pm 0.04	3.57 \pm 0.03	NS
Lactose (g/100 g)	4.45 \pm 0.07	4.64 \pm 0.08	NS
SNF (g/100 g)	8.73 \pm 0.08 ^a	9.2 \pm 0.07 ^b	**
Freezing point	-0.553 \pm 0.005	-0.560 \pm 0.002	NS
pH	6.33 \pm 0.03	6.31 \pm 0.03	NS
EC	3.50 \pm 0.07	3.60 \pm 0.06	NS

a, b- differences significant at $P < 0.05$.

** $P < 0.01$.

heritable trait (Povinelli *et al.*, 2005) in Italian brown cattle but still effect of genetic parameters on electric conductivity needs to be researched in indigenous cattle.

Effect of season on milk composition

The Mean \pm SE values for effect of season on fat, protein, lactose, SNF, freezing point, pH and EC are given in Table 2 along with Table 3. The analysis of variance has significant impact of season on SNF, fat and protein whereas non significant effect on lactose, pH and electric conductivity in Kankrej and Sahiwal cattle. The significant effect of season was found on freezing point in Kankrej cattle whereas non significant was observed in Sahiwal cattle. The post hoc DMRT revealed higher values of Fat, protein, lactose and SNF in winter season than rainy and summer season in Kankrej and Sahiwal cattle. Similar result was reported by Parmar *et al.*, (2020) researched significant effect of season on protein along with fat. Sahu *et al.*, (2018) found that season had significantly ($p < 0.01$) affect protein, solid not fat, fat along with lactose percentage except total solids in Kosali cow. Kabil *et al.*, (2015) observed higher fat and protein during winter season in cow's milk in Egypt.

These variations in season *viz.* Rainy, winter and summer might be due to that it is well established fact that diet and photo period (Parmar *et al.*, 2020) affect milk composition in dairy cattle. During later part of rainy season and first half of winter season cattle offered abundant amount of green fodder and regional grasses and cows are allowed to graze outdoor and this will affect milk composition *viz.* fat. Feeding patterns and milk composition are influenced

by photoperiod in dairy cattle (Parmar *et al.*, 2020). Photo period refers to day length or day light received by an animal which influences series of hormonal changes which ultimately affects the feeding behaviour and milk composition. Therefore combination of these factors affects milk composition.

Effect of stage of lactation on milk composition

The Mean \pm SE values for effect of stage of lactation on fat, protein, lactose, SNF, freezing point, pH and EC are presented in Table 4 along with Table 5. Fat in Sahiwal and Kankrej cattle were shown to have a significant impact on lactation stage, but lactose, EC and freezing point were found to have no effect. The significant effect of lactation stage was found on protein in Sahiwal cattle and significant effect was observed on SNF in Kankrej cattle. The post hoc DMRT revealed higher values of SNF, lactose, protein and fat, in late lactation stage in Sahiwal and Kankrej cattle. Same results have been seen by Parmar *et al.*, (2020) who reported higher protein and fat in late lactation stage. Józwick *et al.*, (2012) observed higher values of fat, protein and lactose in late lactation stage (Days in milk-200) in Polish Holstein-Friesian White along with Black dairy cows. An examination of the late stages of lactation in Holstein Friesian cross cows by Shibru *et al.*, (2019) found no significant differences in fat percentage but a substantial difference in protein %. Gajbhiye *et al.*, (2019) reported increased fat, SNF along with protein in later lactation stages in Gir cows. Shuiep *et al.*, (2016) stated significant impacts of lactation stage on SNF, fat as well as lactose and reported higher fat

Table 2: Means \pm SE different milk composition of kankrej breed in different seasons.

Parameters	Rainy	Winter	Summer	Level of significance
Fat (g/100 g)	3.92 ^a \pm 0.06	4.64 ^b \pm 0.05	4.56 ^b \pm 0.05	**
Protein (g/100 g)	3.47 ^a \pm 0.05	3.67 ^b \pm 0.07	3.56 ^{ab} \pm 0.04	*
Lactose (g/100 g)	4.59 \pm 0.04	4.69 \pm 0.02	4.64 \pm 0.04	NS
SNF (g/100 g)	8.92 ^a \pm 0.08	9.46 ^b \pm 0.12	9.31 ^b \pm 0.14	**
Freezing point	-0.555 ^a \pm 0.002	-0.568 ^b \pm 0.003	-0.557 ^a \pm 0.004	**
pH	6.41 \pm 0.03	6.2 \pm 0.05	6.3 \pm 0.06	NS
EC	3.64 \pm 0.05	3.60 \pm 0.06	3.56 \pm 0.07	NS

a, b, c - Differences significant at $P < 0.05$.

* $P < 0.05$; ** $P < 0.01$.

Table 3: Means \pm SE milk composition of sahiwal breed from different seasons.

Parameters	Rainy	Winter	Summer	Level of significance
Fat (g/100 g)	3.34 ^a \pm 0.04	3.52 ^b \pm 0.05	3.53 ^b \pm 0.03	**
Protein (g/100 g)	3.44 ^a \pm 0.05	3.62 ^b \pm 0.06	3.48 ^a \pm 0.07	*
actose (g/100 g)	4.6 \pm 0.12	4.46 \pm 0.12	4.28 \pm 0.14	NS
SNF (g/100 g)	8.55 ^a \pm 0.1	8.9 ^b \pm 0.09	8.69 ^a \pm 0.17	*
Freezing point	-0.559 \pm 0.01	-0.557 \pm 0.006	-0.544 \pm 0.003	NS
pH	6.38 \pm 0.03	6.29 \pm 0.06	6.32 \pm 0.05	NS
EC	3.49 \pm 0.04	3.45 \pm 0.06	3.56 \pm 0.06	NS

a, b, c - Differences significant at $P < 0.05$.

* $P < 0.05$.

Table 4: Effect of stage of lactation on Means \pm SE of milk composition of Sahiwal cattle.

Parameters	First stage	Second stage	Third stage	Level of significance
Fat (g/100 g)	3.36 ^a \pm 0.04	3.45 ^{ab} \pm 0.04	3.56 ^b \pm 0.05	**
Protein (g/100 g)	3.38 ^a \pm 0.07	3.48 ^a \pm 0.04	3.65 ^b \pm 0.05	**
Lactose (g/100 g)	4.43 \pm 0.12	4.36 \pm 0.16	4.54 \pm 0.12	NS
SNF (g/100 g)	8.54 \pm 0.12	8.74 \pm 0.15	8.92 \pm 0.12	NS
Freezing point	-0.566 \pm 0.008	-0.556 \pm 0.008	-0.538 \pm 0.008	NS
pH	6.38 \pm 0.06	6.31 \pm 0.03	6.26 \pm 0.06	NS
EC	3.49 \pm 0.06	3.51 \pm 0.04	3.53 \pm 0.05	NS

a, b, c - differences significant at P<0.05.

**P<0.01.

Table 5: Effect of stage of lactation on Means \pm SE of milk composition of Kankrej cattle.

Parameters	First stage	Second stage	Third stage	Level of significance
Fat (g/100 g)	4.18 ^a \pm 0.12	4.39 ^a \pm 0.09	4.54 ^b \pm 0.08	*
Protein (g/100 g)	3.48 \pm 0.06	3.54 \pm 0.05	3.66 \pm 0.05	NS
Lactose (g/100 g)	4.62 \pm 0.06	4.65 \pm 0.04	4.67 \pm 0.04	NS
SNF (g/100 g)	8.95 ^a \pm 0.09	9.20 ^a \pm 0.11	9.53 ^b \pm 0.13	**
Freezing point	-0.556 \pm 0.003	-0.558 \pm 0.002	-0.560 \pm 0.003	NS
PH	6.38 \pm 0.05	6.32 \pm 0.04	6.24 \pm 0.05	NS
EC	3.59 \pm 0.04	3.55 \pm 0.06	3.66 \pm 0.06	NS

a, b, c - differences significant at P<0.05.

**P<0.01; *P<0.05.

in late stage of lactation. The variations during various lactation stages in fat might be due to increases in energy deficit, which is always the case in early lactation, but mainly increases at the end of lactation (Vanbergue *et al.*, 2017). Water drawn into milk influences the quantity of lactose synthesis. Lactose and water secretion rates are essentially consistent during lactation (Pollott, 2004). Milk SNF, which is mostly made of minerals, proteins, along with lactose, has a larger amount of additional nonfat ingredients in the late lactation stages (Gurmessia and Melaku, 2012). However, increase in fat and protein in late lactation stage might be due change in milk composition as it reduces partially due to milk reduction in volume at late lactation (Auldish *et al.*, 1998; Hickey *et al.*, 2006).

CONCLUSION

The milk composition is significantly affected by breed factor. Selective breeding could be an effective tool to improve milk quality. The season has a considerable impact on milk fat in Kankrej and Sahiwal cattle, these changes are mostly due to the variation in feed quality in accordance with the season. Stage of lactation is an important physiological parameter divided the lactation period according to milk yield *viz.* early, mid and late lactation according to which milk components *viz.* protein, lactose and fat varies as the lactation proceeds. Milk fat variation is an essential parameter for dairy farmer fetching premium prices. Dairy farmer can improve their income by understanding these variations and tuning the feeding programme and housing practices for improvement.

Conflict of interest: None.

REFERENCES

- Adesina, K. (2012). Effect of breed on the composition of cow milk under traditional management practices in Ado-Ekiti, Nigeria. *J. Appl. Sci. Environ. Manage.* 16: 55-59.
- Alyaqoubi, S., Aminah Abdullah, A., Samudi, M., Abdullah1, N., Addai, Z.A. and Al-ghazali, M. (2014). Effect of different factors on goat Milk Antioxidant Activity. *International Journal of Chem.Tech.Research.* 6(5): 3091-3196.
- Arora, R. and Bhojak, N. (2013). Physiochemical and environmental factors responsible for change in milk composition of milking animal. *The International Journal of Engineering and Science.* 2(1): 275-277.
- Auldish, M.J., Walsh, B.J. and Thomson, N.A. (1998). Seasonal and locational influences on bovine milk composition in New Zealand. *Journal of Dairy Research.* 65: 401- 411.
- Bendelja, D., Prpić, Z., Mikulec, N., Ivkić, Z., Havranek, J. and Antunac, N. (2011). Milk urea concentration in Holstein and Simmental cows. *Mljekarstvo.* 61.
- Bernabucci, U., Lacetera, N., Ronchi, B. and Nardone, A. (2002). Effects of hot season on milk protein fractions in dairy cows. *Anim. Res.* 51: 25-33.
- Bernabucci U., Basiricò L., Morera P., Dipasquale D., Vitali A., Piccioli, Cappelli F. and Calamari L. (2015). Effect of summer season on milk protein fractions in Holstein cows. *J Dairy Sci.* 98(3): 1815-27.
- Bobbo, T., Cecchinato, A., Cipolat-Gotet, C., Stocco, G. and Bittante, G. (2014). Effect of breed and dairy system on milk composition and udder health traits in multi-breed dairy herds. *Acta Agraria Kaposváriensis.* 18 (1): 81-88.
- Brzozowski, P. and Zdziarski, K. (2006). Influence of genotype, age, lactation stage and daily milk performance of Black and White cows on the freezing point of milk. *Polish Journal of Veterinary Sciences.* 62: 93-95.

- FAO (Food and Agricultural Organization) (2004). Milk Producer Group Resource Book: A Practical Guide to Assist Milk producer groups, by Jurjen Draaijer. Rome. Available at <http://www.karmay.org/redirect/strred.asp?docId=21604>. Accessed 21 January 2015.
- Falta, D., Adamski, M., Cejna, V., Hanus, O., Lategan, F., Kupczynski, R., Chladek, G., Filipcik, R. and Machal, L. (2014). The effect of air temperature and breed on bovine milk composition and its processing quality. *Bulg. J. Agric. Sci.* 20(1): 215-219.
- Gajbhiye, P.U., Ahlawat, A.R., Sharma, H.A. and Parikh, S.S. (2019). Effect of stage, season and parity of lactation on milk composition in gir cattle. *Int. J. Curr. Microbiol. App. Sci.* 8(03): 2419-2425.
- Gurmessa, J. and Melaku, A. (2012). Effect of lactation stage, pregnancy, parity and age on yield and major components of raw milk in bred cross Holstein Friesian cows. *World Journal of Dairy and Food Sciences.* 7(2): 146-149.
- Hanna, N., Ahmed, K., Anwar, M., Petrova, A., Hiatt, M. and Hegyi, T. (2004). Effect of storage on breast milk antioxidant activity. *Arch. Dis. Child Fetal Neonatal Ed.* 89:F518-F520.
- Heinrichs, J., Jones, C.M. and Bailey, K. (2016). Milk Components: Understanding Milk Fat and Protein Variation in Your Dairy Herd. Penn State Extension. Available online at <https://extension.psu.edu/media/wysiwyg/media/assets/lanternfly.png>.
- Hickey, D.K., Kilcawley, K.N., Beresford, T.P., Sheehan, E.M. and M.G. Wilkinson. (2006). The influence of a seasonal milk supply on the biochemical and sensory properties of Cheddar cheese. *Int. Dairy J.* 16: 679-690. <https://doi.org/10.1016/j.idairyj.2005.10.017>.
- Ivanov, G., Bilgucu, E., Balabanova, T.B., Ivanova, I.V. and Uzatici, A. (2017). Effect of animal breed, season and milk production scale on somatic cell count and composition of cow milk. *Bulgarian Journal of Agricultural Science.* 23(6): 1047-1052.
- Józwik, A., N. Strzalkowska, E. Bagnicka, W. Grzybek, J. Krzyzewski, E. Polawska, A. Kolataj, J.O. Horbańczuk (2012). Relationship between milk yield, stage of lactation and some blood serum metabolic parameters of dairy cows. *Czech J. Anim. Sci.* 57(8): 353-360.
- Kabil, Osama and Ibrahim Abdou, Ekbal and Elbarbary, Hend and Ali, Mahdy. (2015). Effect of seasonal variation on chemical composition of Cow's milk. *Benha Vet Med J.* 28: 150-154. [10.21608/bvmj.2015.32728](https://doi.org/10.21608/bvmj.2015.32728).
- Kayastha, R.B., Zaman G. and Goswami, R.N. (2008). Factors affecting the milk constituents of native cattle of Assam. *Indian J. Anim. Res.* 42(4): 270-272.
- Kebede E. (2018). Effect of cattle breed on milk composition in the same management conditions. *Ethiop. J. Agric. Sci.* 28(2): 53-63.
- Kedzierska-Matyssek, M., Litwinczuk, Z., Florek, M. and Barłowska, J. (2011). The effects of breed and other factors on the composition and freezing point of cow's milk in Poland. *International Journal of Dairy Technology.* 64, 3: 336-342.
- Mushtaq, M.S., Qureshi, S., Khan, G., Habib, Z.A., Swati and Rahman S.U. (2012). Body condition score as a marker of milk yield and composition in dairy animals. *The Journal of Animal and Plant Sciences.* 22(3): Page: 169-173.
- Myburgh, J., Osthoff, G., Hugo, A., De Wit, M., Nel, K. and Fourie, D. (2012). Comparison of the milk composition of free-ranging indigenous African cattle breeds. *South African Journal of Animal Science.* 42: 1-14.
- Pandy, G.S. and Voskuil, G.C.J. (2011). Manual on milk safety, quality and hygiene. Golden Valley Agricultural Trust. pp 1-50.
- Parmar, P., Lopez, V., Nicolas, Tobin, John, Murphy, Eoin, McDonagh, Arleen, Crowley, Shane, Kelly, Alan, Shalloo and Laurence. (2020). The effect of compositional changes due to seasonal variation on milk density and the determination of season-based density conversion factors for use in the dairy industry. *Foods.* 9. 10.3390/foods9081004.
- Pintia, N., Poljak, F., Daki, A., Blažek, D., Jelen T. and Pintia, V. (2007). Quantitative indicators of milk quality and nutritional status of Simmental and Holstein cows in the Kalnik piedmont region. *Krmiva.* 49 (2): 79-88.
- Pollott, G.E. (2004). Deconstructing milk yield and composition during lactation using biologically based lactation models. *Journal Dairy Science.* 87: 2375-2387.
- Povinelli, M., Gallo, L., Carnier, P., Marcomin, D., Zotto, D.R. and Cassandro, M. (2005). Genetic aspects of milk electrical conductivity in Italian Brown cattle. *Ital. J. Anim. Sci.* 4 (SUPPL. 3):169-171.
- Ramos, T.M., Costa, F.F., Pinto, I.S.B., Pinto, S.M. and Abreu, L.R. (2015). Effect of Somatic Cell Count on Bovine Milk Protein Fractions. *J. Anal. Bioanal. Tech.* 6(5) 1-7.
- Reis, C.B.M., Barreiro, J.R., Mestieri, L., Porcionato, M.A.F. and Santos, M.V. (2013). Effect of somatic cell count and mastitis pathogen on milk composition in Gir cows. *BMC Veterinary Research.* 9(67): 1-7.
- Sahu, J., Bhonsle, D., Mishra, S., Khune, N.V. and Chaturvedani, A.K. (2018). Factors affecting the milk composition of Kosali cow. *Int. J. Curr. Microbiol. App. Sci.* 7(8): 3795-3801.
- Sarkar, U., Gupta, A.K., Sarkar, V., Mohanty, T.K., Raina, V.S. and Prasad, S. (2006). Factors affecting test day milk yield and milk composition in dairy animals. *Journal of Dairying, Foods and Home Sciences.* 25(2): 129-132.
- Shibru, D., Tamir, B., Kasa, F. and Goshu, G. (2019). Effect of Season, Parity, exotic Gene Level and Lactation Stage on Milk Yield and Composition of Holstein Friesian Crosses in Central Highlands of Ethiopia. *Eur. Exp. Biol.* 9(4):15.
- Shuiep, S.E., Eltaher, H.A. and El Zubeir, I.E.M. (2016). Effect of Stage of lactation and order of parity on milk composition and daily milk yield among local and crossbred cows in south darfur state, Sudan. *SUST Journal of Agricultural and Veterinary Sciences.* 17(2): 86-99.
- Singh, N. (2019). Study on quality and bioactive components in milk of various indigenous cattle breeds in hot arid region of Rajasthan. Ph.D. Thesis, RAJUVAS, Bikaner.
- Vanbergue, E., Delaby, L., Peyraud, J.L., Colette, S., Gallard, Y. and Hurtaud, C. (2017). Effects of breed, feeding system and lactation stage on milk fat characteristics and spontaneous lipolysis in dairy cows. *Journal of Dairy Science.* 100(6): 4623-4636.
- Zelalem, Y. (2010). Quality factors that affect Ethiopian milk business: Experiences from selected dairy potential areas. Netherlands Development Organization, Addis Ababa, Ethiopia.