



Phenotypic Extended-spectrum Beta-lactamase Producing *Escherichia coli* among Chickens, Cage Swabs and Wastewaters from Poultry in Indonesia

Sheila Marty Yanestria¹, Wiwiek Tyasningsih², Mustofa Helmi Effendi³,
Aswin Rafif Khairullah⁴, Emmanuel Nnabuike Ugbo⁵

10.18805/ajdfr.DRF-473

ABSTRACT

Background: The aim of the research is to detect Extended-Spectrum Beta-Lactamase (ESBL) producing *Escherichia coli* among chickens, cage swabs and wastewaters from poultry in Indonesia.

Methods: A total of 426 samples were collected, consisting of 200 chicken small intestine contents, 149 cage swabs and 77 wastewater. Isolation and identification of *Escherichia coli* were carried out on samples, followed by antimicrobial resistance testing using the Kirby-bauer diffusion test method. Phenotypic ESBL confirmation testing uses the double disc synergy test (DDST).

Result: Of the 264 *Escherichia coli* isolates, 19.3% (n = 51) were confirmed ESBL producers, out of which 58.8% (n = 30) were obtained from ayam and 41.2% (n = 21) from poultry environments. Out of a total of 21 *Escherichia coli* isolates from poultry environments, 80.9% (n = 17) were from cage swabs and 19.1% (n = 4) from wastewaters. Based on these results, it can be seen that chickens can be considered as a transmission medium for ESBL-producing *Escherichia coli* and play a role in its spread into the environment which can ultimately pose a risk to public health. Coordinated and appropriate action is needed to reduce its impact now and in the future.

Key words: Chicken, Environment, ESBL, *Escherichia coli*, Human health.

INTRODUCTION

Escherichia coli is a commensal microorganism of the gut microbiota of healthy humans and animals, as well as an important opportunistic pathogen, which may be involved in many types of infections (Brower *et al.*, 2017; Debbarma *et al.*, 2019). The plasticity of its genome led to the evolution of this organism and made it a pathogenic strain that can cause important diseases and public health syndromes in humans and animals (Fortuna, 2022; Li *et al.*, 2020). Pathogenic *Escherichia coli* is divided into two groups depending on the location of the disease: extraintestinal pathogenic *Escherichia coli* (ExPEC) and intestinal pathogenic *Escherichia coli* (InPEC) (Pokharel *et al.*, 2023). ExPEC strains are associated primarily with neonatal meningitis and urinary tract infections in adults (Poolman and Wacker, 2016). InPEC strains are associated with diarrheal disease (Rojas-Lopez *et al.*, 2018). ExPEC is the etiological agent of colibacillosis in chickens, which is zoonotic and causes various infections in humans (Mellata, 2013). Studies show that poultry, including chickens, can be a source of transmission of *Escherichia coli* to humans (Stromberg *et al.*, 2017). Human diseases associated with *Escherichia coli* pose a large economic burden due to medical costs and lost productivity (Mitchell *et al.*, 2015).

The most frequent cause of both community-acquired and hospital-acquired extraintestinal infections, such as bloodstream, kidney and urinary tract infections (UTI), is extraintestinal pathogenic *Escherichia coli* (ExPEC). In the United States, extraintestinal infections are thought to cause

¹Department of Veterinary Public Health, Universitas Wijaya Kusuma Surabaya, Jl. Dukuh Kupang XXV No.54, Dukuh Kupang, Dukuh Pakis, Surabaya 60225, East Java, Indonesia.

²Division of Veterinary Microbiology, Faculty of Veterinary Medicine, Universitas Airlangga, Jl. Dr. Ir. H. Soekarno, Kampus C Mulyorejo, Surabaya 60115, East Java, Indonesia.

³Division of Veterinary Public Health, Faculty of Veterinary Medicine, Universitas Airlangga, Jl. Dr. Ir. H. Soekarno, Kampus C Mulyorejo, Surabaya 60115, East Java, Indonesia.

⁴Research Center for Veterinary Science, National Research and Innovation Agency (BRIN), Jl. Raya Bogor Km. 46 Cibinong, Bogor 16911, West Java, Indonesia.

⁵Department of Applied Microbiology, Faculty of Science, Ebonyi State University, Abakaliki 480211, Nigeria.

Corresponding Author: Wiwiek Tyasningsih, Division of Veterinary Microbiology, Faculty of Veterinary Medicine, Universitas Airlangga. Jl. Dr. Ir. H. Soekarno, Kampus C Mulyorejo, Surabaya 60115, East Java, Indonesia. Email: witya_kh@yahoo.com

How to cite this article: Yanestria, S.M., Tyasningsih, W., Effendi, M.H., Khairullah, A.R. and Ugbo, E.N. (2025). Phenotypic Extended-spectrum Beta-lactamase Producing *Escherichia coli* among Chickens, Cage Swabs and Wastewaters from Poultry in Indonesia. Asian Journal of Dairy and Food Research. doi: 1-8. doi: 10.18805/ajdfr.DRF-473.

Submitted: 19-10-2024 **Accepted:** 12-05-2025 **Online:** 28-07-2025

more than 7 million doctor visits, 1 million ER visits and 100,000 hospitalizations annually. According to estimates, the annual expenses of these illnesses in the US range

from \$1 billion to \$1.6 billion (including indirect expenditures) (Manges, 2016). ExPEC infections are a significant source of financial losses for the poultry industry and are crucial to human health. Infections with ExPEC in humans and birds cost the US economy more than \$4 billion annually (Mellata, 2013; Stromberg *et al.*, 2017). Health workers and poultry workers should be educated that people around poultry are a high-risk group for fecal transmission of *Escherichia coli* and should pay attention to biosecurity and good management practices on their farms (Aworh *et al.*, 2021).

Antibiotics are widely used medically in disease prevention (e.g., prophylaxis and metaphylaxis), treatment and growth promotion (Manyi-Loh *et al.*, 2018). It is estimated that two-thirds of global antimicrobials are consumed in the livestock sector (Hosain *et al.*, 2021). Based on a report on antimicrobial agents intended for use in animals published in 2021, 26% of 160 countries analyzed in 2019 still used antibiotics as growth promoters in animal production (Lekagul *et al.*, 2023). Several studies show that the widespread use of antibiotics in poultry farming contributes to an increase in antimicrobial resistance (AMR) (Hedman *et al.*, 2020).

The beta-lactamase enzyme is the biggest cause of antimicrobial resistance, especially in *Escherichia coli* (Wardhana *et al.*, 2021). Dierikx *et al.* (2013) stated that *Escherichia coli* from chicken farms has experienced a high prevalence of resistance to several antibiotics because it can produce Extended Spectrum Beta Lactamase (ESBL). ESBL is an enzyme that can hydrolyze penicillin, cephalosporin and monobactam antibiotics and causes resistance to all of these antibiotics (Akpaka *et al.*, 2021). ESBL-producing *Escherichia coli* which causes multidrug resistance (MDR) often causes infections that are difficult to treat and can be transmitted quickly through food of animal origin so that the impact on health is even wider (Brendecke *et al.*, 2022). Currently, ESBL-producing bacterial infections have become widespread and are often found in humans (Yousefipour *et al.*, 2019). The extensive use of antimicrobials in poultry farming exposes humans to antimicrobial-resistant bacteria through direct and indirect pathways (Hedman *et al.*, 2020). These include exposure through direct contact with livestock or contaminated food products, indirect gene transfer across bacterial species and widespread release of antimicrobial-resistant pathogens into the environment (Brower *et al.*, 2017). Chickens with *Escherichia coli* in their bodies have the potential to spread resistance genes in the environment which can affect human health (Aworh *et al.*, 2020). A health threat will occur if the *Escherichia coli* bacteria that contaminate the environment are ESBL-producing bacteria (Islam *et al.*, 2023). A report conducted by Larsson and Flach (2022) shows that traces of antibiotics, bacteria that are resistant to these antibiotics and genes that cause resistance in bacteria can spread in the environment.

The relationship between livestock and the emergence of ESBL in humans and the transmission of ESBL-producing bacteria must receive special attention, plus the literature regarding the spread of ESBL-producing organisms in Indonesia is still very limited (Widodo *et al.*, 2020). Based on the description above, it is necessary to carry out research regarding the prevalence of extended-spectrum beta-lactamase (ESBL) producing *Escherichia coli* among chickens, cage swabs and wastewaters from poultry in Indonesia, which can then be used in determining efforts to control, spread and prevent epidemics of producing bacteria. ESBL, as well as increasing public awareness.

MATERIALS AND METHODS

Ethical clearance

Animal ethical approval was obtained *via* the ethical clearance committee of the Faculty of Veterinary Medicine, Universitas Wijaya Kusuma Surabaya, Indonesia (ethics no: 86-KKE/2022).

Study area and sample collection

Samples were taken from broiler chicken farms in Lekok, Grati and Rejoso Districts, Pasuruan Regency, Indonesia. Sampling was carried out from March 2023 to August 2023. Samples consisted of broiler chicken intestine contents, cage swabs and wastewater. Sampling of chicken intestinal contents was carried out by slaughtering broiler chickens, then abdominal dissection was carried out to take the contents of the small intestine. Samples were placed in sterile plastic which had been prepared to prevent microbial contamination from the environment. The coop swab sample was taken from a swab on the wall of the chicken coop using a sterile cotton swab, then the sample was put into a tube containing sterile Buffered Peptone Water (HiMedia, India) during transportation. Wastewater samples are taken from rivers or ditches closest to the farm in the amount of 50 ml and then put into sterile plastic bottles for transportation. Samples are stored in a cool box while traveling to the laboratory for research. A total of 426 samples were collected in this study, consisting of 200 chicken small intestine contents, 149 cage swabs and 77 wastewaters.

Microbiological analysis

The *Escherichia coli* enrichment stage was carried out by placing the sample in a test tube containing Buffered Peptone Water (HiMedia, India). Samples were incubated at 37°C for 24 hours. *Escherichia coli* was isolated using selective Eosin Methylene Blue Agar (EMBA) media (Oxoid, England) and incubated for 20-24 hours at 37°C. Next, purification was carried out by streaking a single colony of *Escherichia coli* with a sterile loop from the primary isolate to the new EMBA media. Identification of *Escherichia coli* is carried out using two methods, namely Gram staining and biochemical tests. The biochemical tests in this study

consisted of Indol-Motility, Methyl Red, Voges Proskauer, Citrate and Triple Sugar Iron Agar (Effendi *et al.*, 2021; Wibisono *et al.*, 2021).

Antibiotic susceptibility testing

Antimicrobial susceptibility testing was by Clinical and Laboratory Standards Institute standard procedures using the Kirby-Bauer Diffusion Test method on Mueller Hinton Agar (MHA) media (HiMedia, India. The disk disc (Oxoid, England) used in this study consisted of ciprofloxacin 5 µg, nalidixic acid 30 µg, enrofloxacin 5 µg, erythromycin 15 µg, tetracycline 30 µg, streptomycin 10 µg, gentamicin 10 µg, ampicillin 10 µg, aztreonam 30 µg and amoxicillin-clavulanate 30 µg. The *Escherichia coli* isolate was added to physiological NaCl according to the equation 0.5 Mc. Farland (1.5×10^8 CFU/ml). The suspension that has been made is then spread evenly over the surface of the MHA media using a sterile swab. The antibiotic disc is placed on the surface of the MHA media that has been swabbed at a distance of 25-30 mm and incubated at $35^\circ\text{C} \pm 2^\circ\text{C}$ for 16-18 hours under aerobic conditions. Interpretation of inhibition results by measuring the diameter of the inhibition zone formed by Clinical and Laboratory Standards Institute 2018 (Musa *et al.*, 2020).

ESBL confirmation testing using double disc synergy test (DDST)

Double disc synergy test (DDST) testing was carried out on isolates that were positive for multidrug resistance (MDR). DDST is carried out by placing a combination of cephalosporin antibiotic disks and clavulanic acid (a beta-lactamase enzyme resistance inhibitor) on the surface of MHA media in which *Escherichia coli* bacteria have been planted. DDST in this study used the antibiotic disk amoxicillin-clavulanate 30 µg, ceftazidime 30 µg, aztreonam 30 µg and cefotaxime 30 µg. MHA medium was incubated at 37°C for 24 hours. The formation of a keyhole inhibition zone or a picture of synergy between cephalosporin antibiotic disks and clavulanic acid indicates that the bacteria are positive for producing Extended Spectrum Beta Lactamase (Prayudi *et al.*, 2023).

RESULTS AND DISCUSSION

Overall, 426 samples consisting of chicken small intestine contents (n = 200), cage swabs (n = 149) and wastewater (n = 77) were collected from 40 broiler chicken farms. The overall prevalence of *Escherichia coli* from all sources was 60.6% (n = 264) out of which 58.3% (n = 154) were obtained from chickens and 41.7% (n=110) from poultry environments. Out of a total of 110, *Escherichia coli* isolates from poultry environments, 60.9% (n = 67) were from cage swabs and 39.1% (n = 43) from wastewater. The sample that has high prevalence of *Escherichia coli* was chicken small intestine contents.

The results of the antimicrobial susceptibility test against *Escherichia coli* showed that there was high resistance to enrofloxacin (87.1%), erythromycin (78.8%), ampicillin (75%), nalidixic acid (67.8%) and tetracycline (62.1%). To a lesser extent, *Escherichia coli* was resistant to streptomycin (58.7%), ciprofloxacin (42.1%) and gentamycin (41.3%). Very low resistance rates were seen with the antibiotics aztreonam (14.4%) and amoxicillin-clavulanate (7.2%) (Table 1).

Of 264 *Escherichia coli* isolates, 78.4% (n = 207) were multidrug resistant (MDR). Out of these, 60.9% (n = 126) were chicken isolates, while 39.1% (n = 81) were poultry environmental isolates. Out of a total of 81, *Escherichia coli* isolates from poultry environments, 70.4% (n = 57) were from cage swabs and 29.6% (n = 24) from wastewater.

Of the 264 *Escherichia coli* isolates, 19.3% (n = 51) were confirmed ESBL producers by phenotypic characterization (Fig 1). The prevalence of ESBL producing *Escherichia coli* from chickens was 58.8% (n = 30), while from poultry environments it was 41.2% (n = 21).

Based on the research results, seven types of multidrug resistance patterns were obtained in ESBL-producing *Escherichia coli*. The combination "FQ-AM-M-TE-BL" was the most frequent at 41.2%, followed by the combination "FQ-AM-M-BL" at 23.5%. The results showed that the majority of ESBL-producing *Escherichia coli* had multidrug resistance to 5 antimicrobial classes (AMC) as much as 41.2% (21/51), the remainder had multidrug

Table 1: Antimicrobial resistance profile of *Escherichia coli* isolates from chickens and poultry environments.

Class of antibiotics	Antibiotics	Resistance inhibition zone (mm)	Chickens (n = 154)	Cage swabs (n = 67)	Wastewater (n = 43)
Fluoroquinolone	Ciprofloxacin (CIP)	≤15	65 (42.2%)	34 (50.8%)	12 (27.9%)
	Enrofloxacin (ENR)	≤24	138 (89.6%)	56 (83.58%)	36 (83.7%)
	Nalidixic acid (NA)	≤13	115 (74.7%)	49 (73.1%)	15 (34.8%)
Macrolide	Erythromycin (E)	≤13	107 (69.5%)	59 (88.1%)	42 (97.7%)
Aminoglycoside	Streptomycin (S)	≤11	78 (50.6%)	45 (67.2%)	32 (74.4%)
	Gentamycin (CN)	≤12	75 (48.7%)	25 (37.3%)	9 (20.9%)
Tetracycline	Tetracycline (TE)	≤11	95 (61.7%)	48 (71.6%)	21 (48.8%)
Beta-Lactam	Ampicillin (AMP)	≤13	124 (80.5%)	57 (85.1%)	16 (37.2%)
	Aztreonam (ATM)	≤17	15 (9.7%)	19 (28.4%)	4 (9.3%)
	Amoxicillin-clavulanate (AC)	≤13	8 (5.2%)	8 (11.9%)	3 (7%)

resistance to 4 AMC as much as 39.2% (20/51) and 3 AMC as much as 19.6% (10/51), can be seen in Table 2.

The results showed that *Escherichia coli* had high resistance to enrofloxacin (87.1%), erythromycin (78.8%), ampicillin (75%), nalidixic acid (67.8%) and tetracycline (62.1%). These results are the same as the results of other studies, from different regions in Indonesia, which showed that *Escherichia coli* from broilers in Sukabumi had high resistance to enrofloxacin (72%), ampicillin (100%), nalidixic acid (100%), erythromycin (92 %) and tetracycline (88%) (Hardiati *et al.*, 2021). The high resistance to enrofloxacin, ampicillin, erythromycin and tetracycline is clearly related to the high use of these antibiotics in the poultry farming sector in Indonesia (Efendi *et al.*, 2022). The use of enrofloxacin is the most popular used in poultry in Indonesia at around 60%, followed in sequence by tetracycline, erythromycin and ampicillin (Grabowski *et al.*, 2022).

This research shows that there are 19.5% ESBL-producing *Escherichia coli* (30/154) in isolates originating from chickens. This incidence is slightly lower than previous research in Indonesia which stated that there were 28.75% ESBL-producing *Escherichia coli* in broilers in the Blitar area (Wibisono *et al.*, 2021). The prevalence in this study is much lower compared to the incidence in several other Asian countries, Thailand at 70.5% (Tansawai *et al.*, 2018), India 87% (Brower *et al.*, 2017), dan Philippines at 60.26% (Gundran *et al.*, 2019). The low incidence could be because in Indonesia there is a ban on the use of antibiotic growth promoters (AGP) in the livestock sector which became effective in January 2018 referring to Minister of Agriculture Regulation No. 14/2017. In the past, almost all feed factories added antibiotics as feed additives to commercial feed, this is what caused the high incidence of antibiotic resistance in chicken farms (Chattopadhyay, 2014). However, the discovery of ESBL-producing *Escherichia coli* in chickens is a cause for concern considering that it can attack humans through contaminated chicken carcasses and can spread to the environment surrounding the poultry (Da Silva *et al.*, 2023).

The phenotypic prevalence of ESBL-producing *Escherichia coli* was 19.3%. The prevalence of ESBL

producing *Escherichia coli* from chickens was 58.8% (n = 30), while from poultry environments it was 41.2% (n = 21). In this study, ESBL-producing *Escherichia coli* was not only found in chickens but also in the environment around the cage (cage swabs and wastewater), this proves that there is a spread of ESBL-producing *Escherichia coli* from chickens to the surrounding environment. This incident is strengthened by the fact that ESBL-producing *Escherichia coli* from chickens has the same multidrug resistance pattern as ESBL-producing *Escherichia coli* from cage swabs and wastewater (Patricio *et al.*, 2022). The "FQ-AM-M-TE-BL" pattern is the most common in chickens as well as in cage swabs and wastewater.

The research results of Kwoji *et al.* (2019) shows that the transfer of resistance genes in *Escherichia coli* is quite high in the poultry farming environment. Poultry farming environments often have a high concentration of microorganisms, thus triggering the creation of poor environmental quality and benefiting the spread of ESBL-producing *Escherichia coli* because the ESBL coding gene in the plasmid can be transferred to other bacteria (Yanestria *et al.*, 2022; Widodo *et al.*, 2022). Antibiotic resistance bacteria (ARB) and antibiotic-resistant genes (ARG) from destroyed ARB can spread in the environment (Courti *et al.*, 2022; Tyasningsih *et al.*, 2022). Unwise use




Fig 1: *Escherichia coli* was confirmed as ESBL using the DDST method.

Table 2: Data on multidrug resistance of ESBL-producing *Escherichia coli*.

Number of AMC	Multidrug resistance pattern	<i>Escherichia coli</i> (n =51)			Total
		Chicken	Cage swabs	Wastewaters	
5	FQ-AM-M-TE-BL	8	11	2	21
4	FQ-AM-M-BL	9	2	1	12
4	FQ-M-TE-BL	5	2	0	7
4	AM-M-TE-BL	0	0	1	1
3	FQ-M-BL	4	1	0	5
3	FQ-TE-BL	1	1	0	2
3	FQ-AM-BL	3	0	0	3
Total		30	17	4	51

Note: FQ: Fluoroquinolone; AM: Aminoglycosides; M: Macrolides; TE: Tetracycline; BL: Beta-lactam; AMC: Antimicrobial class.

of antibiotics in animals exacerbates the spread of ARBs and ARGs. Humans and animals excrete ARBs and ARGs through urine and feces into the environment (Newton *et al.*, 2015). Potential sources of AMR in poultry production consist of air, dust, soil, feed and rodents or other animals. From these sources, AMR bacteria and genes can be transmitted indirectly to the environment (Martínez-Álvarez *et al.*, 2022; Effendi *et al.*, 2018). Several studies have shown the role of air as a means of dissemination and survival both inside and outside the farm and poultry waste is considered the main vector for the spread of AMR bacteria (Vounba *et al.*, 2019; Hedman *et al.*, 2020).

Large amounts of antibiotic residues are also excreted from the animal's body (Arsène *et al.*, 2022). It is estimated that at least 70% of the total antibiotics consumed are excreted unchanged (Kümmerer, 2009). The released antibiotic, after being reduced by metabolism, has a non-killing concentration, known as a sub-inhibitory concentration (Vasilchenko and Rogozhin, 2019; Ramandinianto *et al.*, 2020). These antibiotics are mixed intensively with pathogenic bacteria in wastewater, more intense contact occurs in non-killing concentrations, the bacteria absorb the antibiotics and begin to adapt to them, eventually developing antibiotic resistance (Rizzo *et al.*, 2013; Riwu *et al.*, 2020). This statement is also reinforced by other literature from Meng *et al.* (2022) which states that waste from animals given antibiotics contains antibiotic residues and resistant bacteria. Other literature states that horizontal plasmid-mediated transfer of Antibiotic Resistance Genes (ARGs) is recognized as the most dominant way of spreading ARGs in humans, animals and the environment. Antibiotic selective pressure is considered to be one of the important contributors to promoting the spread of antibiotic resistance through horizontal gene transfer (HGT) (Liu *et al.*, 2020). ESBLs can also be spread *via* HGT, as suggested by the study of Franz *et al.* (2015) that *Escherichia coli*-producing ESBL is at risk of spreading ARGs, especially in susceptible individuals.

Various genetic mechanisms are involved in the acquisition and spread of AMR (Aminov, 2021). The *Escherichia coli* mobilome includes a variety of mobile and mobilizable genetic elements, including plasmids, transposons, insertion sequences and integrons (intl), the latter of which is known to be involved in the spread of antibiotic resistance, especially among Gram-negative bacteria (Gillings, 2014; Pradika *et al.*, 2019; Durairajan *et al.*, 2021). Integrons are genetic structures that contain AMR genes in their variable regions (as gene cassettes) and have been detected in poultry farms in various studies (Kalantari *et al.*, 2021; Perez-Etayo *et al.*, 2018). Another thing to worry about is that the ESBL-producing *Escherichia coli* in this study is multidrug-resistant (MDR), even resistant to 5 classes of antibiotics. An isolate is defined as multidrug-resistant if it is insensitive to one or more antimicrobials in ≥ 3 different antimicrobial classes (Varga *et al.*, 2019). ESBL-producing *Escherichia coli* in livestock animals has

become a public health concern in recent years (Ansharieta *et al.*, 2021). Infection in humans can cause treatment failure, prolonged illness and increase the death rate, especially in patients with septicemia and urinary tract infections (Rodroo *et al.*, 2021).

Based on the results of this research, it can be seen that poultry can be considered as a transmission medium for MDR bacteria, one of which is that ESBL plays a role in the spread of MDR into the environment which can ultimately pose a risk to public health. Antimicrobial resistance is a global and growing problem that requires a solution through a "One Health" approach (Velazquez-Meza *et al.*, 2022; Yunita *et al.*, 2020). Coordinated and appropriate action is needed to mitigate its impact now and in the future, as well as measures to ensure the economic viability of the poultry sector along with public health security.

CONCLUSION

This study shows that chickens can be considered as a transmission medium for ESBL-producing *Escherichia coli* and play a role in its spread into the environment which can ultimately pose a risk to public health. It is important to increase public awareness of hygiene practices in poultry and chicken slaughtering, as well as during food processing. Control of antimicrobial use needs to be improved to reduce the spread of ESBL-producing *Escherichia coli*.

ACKNOWLEDGEMENT

The author would like to thank the Institute for Research and Community Service (LPPM) Universitas Wijaya Kusuma Surabaya and the Faculty of Veterinary Medicine Universitas Wijaya Kusuma Surabaya for providing the UWKS Research Fund 2023.

Disclaimers

The views and conclusions expressed in this article are solely those of the authors and do not necessarily represent the views of their affiliated institutions. The authors are responsible for the accuracy and completeness of the information provided but do not accept any liability for any direct or indirect losses resulting from the use of this content.

Informed consent

All animal procedures for experiments were approved by the Committee of Experimental Animal care and handling techniques were approved by the University of Animal Care Committee.

Conflict of interest

The authors declare that there are no conflicts of interest regarding the publication of this article. No funding or sponsorship influenced the design of the study, data collection, analysis, decision to publish or preparation of the manuscript.

REFERENCES

- Akpaka, P.E., Vaillant, A., Wilson, C., Jayaratne, P. (2021). Extended spectrum beta-lactamase (ESBL) produced by gram-negative bacteria in trinidad and tobago. *International Journal of Microbiology*. **2021(1)**: 5582755. doi: 10.1155/2021/5582755.
- Aminov, R. (2021). Acquisition and spread of antimicrobial resistance: A tet(X) case study. *International Journal of Molecular Sciences*. **22**: 3905. doi: 10.3390/ijms22083905.
- Ansharieta, R., Ramandinianto, S.C., Effendi, M.H., Plumeriastuti, H. (2021). Molecular identification of *bla*CTX-M and *bla*TEM genes encoding extended-spectrum β -lactamase (ESBL) producing *Escherichia coli* isolated from raw cow's milk in East Java, Indonesia. *Biodiversitas*. **22(4)**: 1600-1605. DOI: 10.13057/biodiv/d220402.
- Arsène, M.M.J., Davares, A.K.L., Viktorovna, P.I. andreevna, S.L., Sarra, S., Khelifi, I., Sergueievna, D.M. (2022). The public health issue of antibiotic residues in food and feed: Causes, consequences and potential solutions. *Veterinary World*. **15(3)**: 662-671. DOI: 10.14202/vetworld.2022.662-671.
- Aworh, M.K., Kwaga, J., Okolocha, E., Harden, L., Hull, D., Hendriksen, R.S., Thakur, S. (2020). Extended-spectrum β -lactamase-producing *Escherichia coli* among humans, chickens and poultry environments in Abuja, Nigeria. *One Health Outlook*. **2(1)**: 8. doi: 10.1186/s42522-020-00014-7.
- Aworh, M.K., Kwaga, P.K., Hendriksen, R.S., Okolocha, C., Thakur, S. (2021). *Escherichia coli* isolated from humans, chickens and poultry environments. *Antimicrobial Resistance and Infection Control*. **10(1)**: 58. doi: 10.1186/s13756-021-00930-x.
- Brendecke, J., Homeier-Bachmann, T., Schmitz Ornés, A., Guenther, S., Heiden, S.E., Schwabe, M., Eger, E., Schaufler, K. (2022). Multidrug-resistant high-risk *Escherichia coli* and *Klebsiella pneumoniae* clonal lineages occur in black-headed gulls from two conservation islands in Germany. *Antibiotics (Basel)*. **11(10)**: 1357. doi: 10.3390/antibiotics11101357.
- Brower, C.H., Mandal, S., Hayer, S., Sran, M., Zehra, A., Patel, S.J., Kaur, R., Chatterjee, L., Mishra, S., Das, B.R., Singh, P., Singh, R., Gill, J.P.S., Laxminarayan, R. (2017). The prevalence of extended-spectrum beta-lactamase-producing multidrug-resistance *Escherichia coli* in poultry chickens and variation according to farming practices in Punjab, India. *Environmental Health Perspectives*. **125(7)**: 077015. doi: 10.1289/EHP292.
- Chattopadhyay, M.K. (2014). Use of antibiotics as feed additives: a burning question. *Frontiers in Microbiology*. **5(1)**: 334. doi: 10.3389/fmicb.2014.00334.
- Courti, I., Muja, C., Maho, T., Saint, F.P., Guillot, P. (2022). Degradation of bacterial antibiotic resistance genes during exposure to non-thermal atmospheric pressure plasma. *Antibiotics (Basel)*. **11(6)**: 747. doi: 10.3390/antibiotics11060747.
- Da Silva, F.B., Ferreira, M.R.A., Sobrinho, I.D.S.J., Dias, M., Rodrigues, R.R., Moreira, C.N. (2023). Occurrence of ESBL-producing avian pathogenic *Escherichia coli* (APEC) isolates in spiced chicken meat in Goiás, Brazil. *Letters in Applied Microbiology*. **76(2)**: ovac070. doi: 10.1093/lambio/ovac 070.
- Debbarma, S., Bora, D.P., Hazarika, R.A., Tamuly, S., Barua, A.G., Das, P., Goswami, C., Joute, J.R., Sinha, S. (2019). Molecular characterization of *Escherichia coli* isolates from food animals. *Indian Journal of Animal Research*. **54(8)**: 985-993. doi: 10.18805/ijar.B-3869.
- Dierikx, C., Goot, J., Fabri, T., Essen-Zandbergen, Smith, H., Mevius, D. (2013). Extended-spectrum- β -lactamase-and AmpC- β -lactamase-producing *Escherichia coli* in Dutch broilers and broiler farmers. *Journal of Antimicrobial Chemotherapy*. **68(1)**: 60-67. DOI: 10.1093/jac/dks349.
- Durairajan R., Murugan M., Karthik K., Porteen K. (2021). Farmer's stance on antibiotic resistance to *E. coli* and extended spectrum - β -lactamase producing (ESBL) *E. coli* isolated from poultry droppings. *Asian Journal of Dairy and Food Research*. **40(1)**: 88-93. doi: 10.18805/ajdfr.DR-1574.
- Efendi, R., Sudarnika, E., Wibawan, I.W.T., Purnawarman, T. (2022) An assessment of knowledge and attitude toward antibiotic misuse by small-scale broiler farmers in Bogor, West Java, Indonesia. *Veterinary World*. **15(3)**: 707-713. doi: 10.14202/vetworld.2022.707-713.
- Effendi, M.H., Harijani, N., Yanestria, S.M., Hastutiek, P. (2018). Identification of shiga toxin-producing *Escherichia coli* in raw milk samples from dairy cows in Surabaya, Indonesia. *Philippine Journal of Veterinary Medicine*. **55(SI)**: 109-114.
- Effendi, M.H., Tyasningsih, W., Yurianti, Y.A., Rahmahani, J., Harijani, N., Plumeriastuti, H. (2021). Presence of multidrug resistance (MDR) and extended-spectrum beta-lactamase (ESBL) of *Escherichia coli* isolated from cloacal swab of broilers in several wet markets in Surabaya, Indonesia. *Biodiversitas*. **22(1)**: 304-310. doi: 10.13057/biodiv/d220137.
- Fortuna, M.A. (2002). The phenotypic plasticity of an evolving digital organism. *Royal Society Open Science*. **9**: 220852. doi: 10.1098/rsos.220852.
- Franz, E., Veenman, C., van Hoek, A.H.A.M., Husman, A.R., Blaak, H. (2015). Pathogenic *Escherichia coli* producing extended-spectrum β -lactamases isolated from surface water and wastewater. *Scientific Reports*. **5(1)**: 1-9. doi: 10.1038/srep14372.
- Gillings, M.R. (2014). Integrins: Past, present and future. *Microbiology and Molecular Biology Reviews*. **78(2)**: 257-277. doi: 10.1128/MMBR.00056-13.
- Grabowski, L., Gaffke, L., Pierzynowska, K., Cyske, Z., Choszcz, M., Węgrzyn, G., Węgrzyn, A. (2022). Enrofloxacin-the ruthless killer of eukaryotic cells or the last hope in the fight against bacterial infections. *International Journal of Molecular Sciences*. **23(7)**: 3648. doi: 10.3390/ijms2307 3648.
- Gundran, R.S., Cardenio, P.A., Villanueva, M.A., Sison, F.B., Benogno, C.C., Kreausakon, K., Pichpol, D., Punyapornwithaya, V. (2019). Prevalence and distribution of *bla*CTX-M, *bla*SHV, *bla*TEM genes in extended- spectrum β -lactamase-producing *E. coli* isolates from broiler farms in the Philippines. *BMC Veterinary Research*. **15(1)**: 227. doi: 10.1186/s12917-019-1975-9.
- Hardiati, A., Safika, S., Wibawan, I.W.T., Indrawati, A., Pasaribu, F.H. (2021). Isolation and detection of antibiotics resistance genes of *Escherichia coli* from broiler farms in Sukabumi, Indonesia. *Journal of Advanced Veterinary and Animal Research*. **8(1)**: 84-90. doi: 10.5455/javar.2021.h489.

- Hedman, H.D., Vasco, K.A., Zhang, L. (2020). A review of antimicrobial resistance in poultry farming within low-resource settings. *Animals (Basel)*. **10(8)**: 1264. doi: 10.3390/ani10081264.
- Hosain, M.Z., Kabir, S.M.L., Kamal, M.M. (2021). Antimicrobial uses for livestock production in developing countries. *Veterinary World*. **14(1)**: 210-221. doi: 10.14202/vetworld.2021.210-221.
- Islam, M.S., Rahman, A.M.M.T., Hassan, J., Rahman, M.T. (2023). Extended-spectrum beta-lactamase in *Escherichia coli* isolated from humans, animals and environments in Bangladesh: A One Health perspective systematic review and meta-analysis. *One Health*. **16(1)**: 100526. doi: 10.1016/j.onehlt.2023.100526.
- Kalantari, M., Sharifiyazdi, H., Asasi, K., Abdi-Hachesoo, B. (2021). High incidence of multidrug resistance and class 1 and 2 integrons in *Escherichia coli* isolated from broiler chickens in South of Iran. *Veterinary Research Forum*. **12(1)**: 101-107. DOI: 10.30466/vrf.2019.96366.2309.
- Kümmerer, K. (2009). Antibiotics in the aquatic environment-A review-part I. *Chemosphere*. **75(4)**: 417-434. doi: 10.1016/j.chemosphere.2008.11.086.
- Kwoji, I.D., Musa, J.A., Daniel, N., Mohzo, D.L., Bitrus, A.A., Ojo, A.A., Ezema, K.U. (2019). Extended-spectrum β -lactamase-producing *Escherichia coli* in chickens from small-scale (backyard) poultry farms in Maiduguri, Nigeria. *International Journal of One Health*. **5(1)**: 26-30. DOI:10.14202/IJOH.2019.26-30.
- Larsson, D.G.J., Flach, C.F. (2022). Antibiotic resistance in the environment. *Nature Reviews Microbiology*. **20(1)**: 257-269. doi: 10.1038/s41579-021-00649-x.
- Lekagul, A., Kirivan, S., Tansakul, N., Krisanaphan, C., Srinha, J., Laoprasert, T., Kaewkhankhaeng, W., Tangcharoensathien, V. (2023). Antimicrobial consumption in food-producing animals in Thailand between 2017 and 2019: The analysis of national importation and production data. *PLoS One*. **18(4)**: e0283819. DOI: 10.1371/journal.pone.0283819.
- Li, J., Fan, Q., Mao, C., Jin, M., Yi, L., Wang, Y. (2020). Effect of biofilm formation on the *Escherichia coli* drug resistance of isolates from pigs in central China. *Indian Journal of Animal Research*. **55(6)**: 697-703. doi: 10.18805/IJAR.B-1304.
- Liu, Y., Tong, Z., Shi, J., Jia, Y., Yang, K., Wang, Z. (2020). Correlation between exogenous compounds and the horizontal transfer of plasmid-borne antibiotic resistance genes. *Microorganisms*. **8(8)**: 1211. doi: 10.3390/microorganisms8081211.
- Manges, A.R. (2016). *Escherichia coli* and urinary tract infections: The role of poultry-meat. *Clinical Microbiology and Infection*. **22(2)**: 122-129. doi: 10.1016/j.cmi.2015.11.010.
- Manyi-Loh, C., Mamphweli, S., Meyer, E., Okoh, A. (2018). Antibiotic use in agriculture and its consequential resistance in environmental sources: Potential public health implications. *Molecules*. **23(4)**: 795. DOI: 10.3390/molecules23040795.
- Martínez-Álvarez, S., Sanz, S., Olarte, C., Hidalgo-Sanz, R., Carvalho, I., Fernández-Fernández, R., Campaña-Burguet, A., Latorre-Fernández, J., Zarazaga, M., Torres, C. (2022). Antimicrobial resistance in *Escherichia coli* from the broiler farm environment, with detection of SHV-12-producing isolates. *Antibiotics (Basel)*. **11(4)**: 444. doi: 10.3390/antibiotics11040444.
- Mellata, M. (2013). Human and avian extraintestinal pathogenic *Escherichia coli*: Infections, zoonotic risks and antibiotic resistance trends. *Foodborne Pathogens and Disease*. **10(11)**: 916-932. doi: 10.1089/fpd.2013.1533.
- Meng, M., Li, Y., Yao, H. (2022). Plasmid-mediated transfer of antibiotic resistance genes in soil. *Antibiotics (Basel)*. **11(4)**: 525. doi: 10.3390/antibiotics11040525.
- Mitchell, N.M., Johnson, J.R., Johnston, B., Curtiss, R., Mellata, M. (2014). Zoonotic potential of *Escherichia coli* isolates from retail chicken meat products and eggs. *Applied and Environmental Microbiology*. **81(3)**: 1177-1187. DOI: 10.1128/AEM.03524-14.
- Musa, L., Proietti, P.C., Branciar, R., Menchetti, L., Bellucci, S., Ranucci, D., Marenzoni, M.L., Franciosini, M.P. (2020). Antimicrobial susceptibility of *Escherichia coli* and ESBL-Producing *Escherichia coli* diffusion in conventional, organic and antibiotic-free meat chicken at slaughter. *Animals*. **10(7)**: 1215. doi: 10.3390/ani10071215.
- Newton, R.J., McLellan, S.L., Dila, D.K., Vineis, J.H., Morrison, H.G., Eren, A.M., Sogin, M.L. (2015). Sewage reflects the microbiomes of human populations. *MBio*. **6(2)**: e02574. doi: 10.1128/mBio.02574-14.
- Patricio, T.C.D.C., Farias, B.O., Santiago, G.S., Souza, V.R.S., Pimenta, R.L., de Oliveira, C.C., Coelho, I.S., de Souza, M.M.S., Coelho, S.M.O. (2022). Production of extended-spectrum beta-lactamases in *Escherichia coli* isolated from poultry in Rio de Janeiro, Brazil. *Brazilian Journal of Veterinary Medicine*. **44(4)**: e001722. doi: 10.29374/2527-2179.bjvm001722.
- Pérez-Etayo, L., Berzosa, M., González, D., Vitas, A.I. (2018). Prevalence of integrons and insertion sequences in ESBL-producing *E. coli* isolated from different sources in Navarra, Spain. *International Journal of Environmental Research and Public Health*. **15(10)**: 2308. doi: 10.3390/ijerph15102308.
- Pokharel, P., Dhakal, S., Dozois, C.M. (2023). The diversity of *Escherichia coli* pathotypes and vaccination strategies against this versatile bacterial pathogen. *Microorganisms*. **11(2)**: 344. doi: 10.3390/microorganisms11020344.
- Poolman, J.T., Wacker, M. (2016). Extraintestinal pathogenic *Escherichia coli*, a common human pathogen: Challenges for vaccine development and progress in the field. *The Journal of Infectious Diseases*. **213(1)**: 6-13. doi: 10.1093/infdis/jiv429.
- Pradika, A.Y., Chusniati, S., Pumama, M.T.E., Effendi, M.H., Yudhana, A., Wibawati, P.A. (2019). Total test of *Escherichia coli* on fresh cow milk at dairy farmer cooperative (KPSP) karyo ngremboko prwoharjo banyuwangi. *Jurnal Medik Veteriner*. **2(1)**: 1-6. doi: 10.20473/jmv.vol2.iss1.2019.1-6.
- Prayudi, S.K.A., Effendi, M.H., Lukiswanto, B.S., Az Zahra, R.L., Benjamin, M.I., Kurniawan, S.C., Khairullah, A.R., Silaen, O.S.M., Lisnanti, E.F., Baihaqi, Z.A., Widodo, A., Riwu, K.H.P. (2023). Detection of genes on *Escherichia coli* producing extended spectrum β -lactamase isolated from the small intestine of ducks in traditional markets Surabaya City, Indonesia. *Journal of Advanced Veterinary Research*. **13(8)**: 1600-1608.
- Ramandinianto, S.C., Khairullah, A.R., Effendi, M.H., Hestiana, E.P. (2020). Profile of multidrug resistance (MDR) and methicillin resistant *staphylococcus aureus* (MRSA) on dairy farms in east java province, Indonesia. *Indian Journal of Forensic Medicine and Toxicology*. **14(4)**: 3439-3445. doi: 10.37506/ijfvt.v14i4.12157.

- Riwu, K.H.P., Effendi, M.H., Rantam, F.A. (2020). A review of extended spectrum β -lactamase (ESBL) producing *klebsiella pneumoniae* and multidrug resistant (MDR) on companion animals. *Systematic Reviews in Pharmacy*. **11(7)**: 270-277. doi: 10.31838/srp.2020.7.43.
- Rizzo, L., Manaia, C., Merlin, C., Schwartz, T., Dagot, C., Ploy, M., Michael, I., Fatta-Kassinos, D. (2013). Urban wastewater treatment plants as hotspots for antibiotic resistant bacteria and genes spread into environment: A review. *Science of The Total Environment*. **447(1)**: 345-360.
- Rodroo, J., Intanon, M., Kreausukon, K., Kongkaew, A., Bender, J., Awainanont, N. (2021). Occurance of extended-spectrum beta-bactamase producing *E. coli* in broiler farm workers and the farm environment in Chiang Mai-Lumphun, Thiland. *Veterinary Integrative Sciences*. **19(1)**: 23-35. doi: 10.12982/VIS.2021.002.
- Rojas-Lopez, M., Monterio, R., Pizza, M., Desvaux, M., Rosini, R. (2018). Intestinal pathogenic *Escherichia coli*: Insight for vaccine development. *Frontiers in Microbiology*. **9(1)**: 440. doi: 10.3389/fmicb.2018.00440.
- Stromberg, Z.R., Johnson, J.R., Fairbrother, J.M., Kilbourne, J., Van Goor, A., Curtiss Rd, R., Mellata, M. (2017). Evaluation of *Escherichia coli* isolates from healthy chickens to determine their potential risk to poultry and human health. *PLoS One*. **12(7)**: e0180599. DOI: 10.1371/journal.pone.0180599.
- Tansawai, U., Sanguansersri, D., Na-udom, A., Walsh, T.R., Niumsup, P.R. Occurrence of extended spectrum β -lactamase and AmpC genes among multidrug-resistant *Escherichia coli* and emergence of ST131 from poultry meat in Thailand. *Food Control*. **84(1)**: 159-164. doi: 10.1016/j.foodcont.2017.07.028.
- Tyasningsih, W., Ramandinianto, S.C., Ansharieta, R., Witaningrum, A.M., Permatasari, D.A., Wardhana, D.K., Effendi, M.H., Ugbo, E.N. (2022). Prevalence and antibiotic resistance of *Staphylococcus aureus* and *Escherichia coli* isolated from raw milk in east java, Indonesia. *Veterinary World*. **15(8)**: 2021-2028. DOI: 10.14202/vetworld.2022.2021-2028.
- Varga, C., Guerin, M.T., Brash, M.L., Siavic, D., Boerlin, P., Susta, L. (2019). Antimicrobial resistance in *Campylobacter jejuni* and *Campylobacter coli* isolated from small poultry flocks in Ontario, Canada: A two year surveillance study. *PLoS One*. **14(8)**: e0221429. DOI: 10.1371/journal.pone.0221429.
- Vasilchenko, A.S., Rogozhin, E.A. (2019). Sub-inhibitory effects of antimicrobial peptides. *Frontiers in Microbiology*. **10(1)**: 1160. DOI: 10.3389/fmicb.2019.01160.
- Velazquez-Meza, M.E., Galarde-López, M., Carrillo-Quiróz, B., Alpuche-Aranda, C.M. (2022). Antimicrobial resistance: One Health Approach. *Veterinary World*. **15(3)**: 743-749. DOI: 10.14202/vetworld.2022.743-749.
- Vounba, P., Arsenault, J., Bada-Alambédji, R., Fairbrother, J.M. (2019). Prevalence of antimicrobial resistance and potential pathogenicity and possible spread of third generation cephalosporin resistance, in *Escherichia coli* isolated from healthy chicken farms in the region of Dakar, Senegal. *PLoS One*. **14(3)**: e0214304. DOI: 10.1371/journal.pone.0214304.
- Wardhana, D.K., Safitri, D.A., Anisa, S., Harijani, N., Estoe pangestie, S., Maghfiroh, L. (2021). Prevalence of extended-spectrum beta-lactamases in producing *Escherichia coli* in beef sold in traditional markets in Surabaya, Indonesia. *Biodiversitas*. **22(5)**: 2789-2793. DOI: 10.13057/biodiv/d220542.
- Wibisono, F.J., Sumiarto, B., Untari, T., Effendi, M.H., Permatasari, D.A., Witaningrum, A.M. (2021). Molecular identification of CTX Gene of extended spectrum beta-lactamase (ESBL) producing *Escherichia coli* on layer chicken in Blitar, Indonesia. *Journal of Animal and Plant Sciences*. **31(4)**: 954-959. DOI: 10.36899/JAPS.2021.4.0289.
- Widodo, A., Effendi, M.H., Khairullah, A.R. (2020). Extended-spectrum beta-lactamase (ESBL)-producing *Escherichia coli* from livestock. *Systematic Reviews in Pharmacy*. **11(7)**: 382-392. DOI: 10.31838/srp.2020.7.57.
- Widodo, A., Lamid, M., Effendi, M.H., Khairullah, A.R., Riwu, K.H.P., Yustinasari, L.R., Kurniawan, S.C., Ansori, A.N.M., Silaen, O.S.M, Dameanti, F.N.A.E.P. (2022). Antibiotic sensitivity profile of multidrug-resistant (MDR) *Escherichia coli* isolated from dairy cow's milk in Probolinggo, Indonesia. *Biodiversitas*. **23(10)**: 4971-4976. DOI: 10.13057/biodiv/d231002.
- Yanestria, S.M., Dameanti, F.N.A.E.P., Musayannah, B.G., Pratama, J.W.A., Witaningrum, A.M., Effendi, M.H., Ugbo, E.N. (2022). Antibiotic resistance pattern of extended-spectrum β -lactamase (ESBL) producing *Escherichia coli* isolated from broiler farm environment in Pasuruan district, Indonesia. *Biodiversitas*. **23(9)**: 4460-4465. DOI: 10.13057/biodiv/d230911.
- Yousefipour, M., Rasoulinejad, M., Hadadi, A., Esmailpour, N., Abdollahi, A., Jafari, S., Khorsand, A. (2019). Bacteria producing extended spectrum β -lactamases (ESBLs) in hospitalized patients: Prevalence, antimicrobial resistance pattern and its main determinants. *Iranian Journal of Pathology*. **14(1)**: 61-67. DOI: 10.30699/IJP.14.1.61.
- Yunita, M.N., Effendi, M.H., Rahmiani, R.P., Arifah, S., Yanestria, S.M. (2020). Identification of *spa* gene for strain typing of methicillin resistant *Staphylococcus aureus* (MRSA) isolated from nasal swab of dogs. *Biochemical and Cellular Archives*. **20(1)**: 2999-3004. DOI: 10.35124/bca.2020.20.S1.2999.