



Assessment of Dietary Habits and Sociodemographic Profiles in the Rabat-Salé-Kénitra Region

Nada Belmejdoub¹, Imane Fadel², Sarah Drider¹,
Hefdhallah Alaizari¹, Milouda Chebabe³, Mohammed Ouhssine¹

10.18805/ajdfr.DRF-498

ABSTRACT

Background: A healthy and varied diet plays a fundamental role in supporting balanced physical development, maintaining overall health, and mitigating the risk of chronic diseases. Understanding dietary habits is essential for designing effective interventions and promoting nutritional well-being. This study investigates the dietary habits of individuals in the Rabat-Salé-Kénitra region, with a focus on sociodemographic and anthropometric characteristics, meal consumption patterns and food choices. The findings aim to highlight gender-specific trends and general dietary behaviors within the population.

Methods: A prospective cross-sectional observational study design was employed to evaluate dietary habits. Data were collected using a structured questionnaire in two phases: the first phase focused on sociodemographic and anthropometric information, while the second captured detailed dietary habits. A total of 1,012 participants were recruited, comprising 60.4% females and 39.6% males. Statistical analyses were conducted using SPSS software (version 22) and Chi-square tests were applied to identify significant relationships between variables. Body Mass Index (BMI) categories were calculated and dietary habits were analyzed by gender.

Result: The study population was predominantly female (60.4%) and included a higher proportion of younger individuals, with 45.4% aged 18-30 years. Most participants (85.9%) had university-level education and civil servants represented the largest professional group (52.9%). Females were overrepresented among unemployed participants, while married individuals comprised the majority (54.8%) of family statuses. BMI analysis indicated that overweight and obesity were more prevalent among females, whereas males dominated the underweight category ($p = 0.0005$). Regarding dietary habits, 53.6% of participants consumed four meals daily, with 80.7% consuming bread and pastries daily and 60.2% consuming fruits and vegetables 4-5 times weekly. Frequent consumption of meat and poultry was observed, with females showing higher intake levels. No significant gender differences were found in fish/seafood, cereals, or dairy product consumption and snack and sweet product consumption patterns were consistent across genders.

Key words: Body mass index (Bmi), Dietary habits, Gender differences, Rabat-salé-kénitra region, Sociodemographic characteristics.

INTRODUCTION

Adapted and balanced nutrition that meets the needs of the body accompanied by regular physical activity helps maintain good health (WHO) (Organization, 2003).

Food plays a central role in societal concerns, with its strong link to health long established. National programs like the PNNS promote healthy diets and physical activity (Hercberg 2011; Lecerf, 2018). Epidemiological studies confirm the relationship between diet and health globally (O'Neil *et al.*, 2014). However, food choices are shaped by economic, cultural, and social factors (Fieldhouse, 2013). Industrialization and globalization have improved living standards but led to unhealthy lifestyles, disproportionately affecting disadvantaged groups and increasing chronic diseases like obesity, diabetes, and cardiovascular conditions (Organization, 2003; Gowshall and Taylor-Robinson, 2018). Non-communicable diseases (NCDs) account for 71% of global deaths, including 15 million premature deaths annually (Perico and Remuzzi, 2012). Nutrition knowledge significantly influences dietary decisions and prevents NCDs (Penzavecchia *et al.*, 2022; Albrahim *et al.*, 2023). Adopting healthy lifestyles and balanced diets can mitigate NCD risks and improve global health (Passi, 2017; Pandey *et al.*, 2021).

¹Laboratory of Natural Resources and Sustainable Development, Faculty of Science, Ibn Tofail University, Morocco.

²Laboratory of Plants, Animals Productions and Agro-Industry, Faculty of Science, Ibn Tofail University, Kenitra, Morocco.

³Laboratory of Health Sciences and Technologies, Department of Science, Higher institute of Health Sciences, Hassan First University, Settat, Morocco.

Corresponding Author: Hefdhallah Alaizari, Laboratory of Natural Resources and Sustainable Development, Faculty of Science, Ibn Tofail University, Morocco. Email: alaizari2@gmail.com
ORCID: <https://orcid.org/0009-0004-8042-5259>, <https://orcid.org/0009-0007-3224-0403>, <https://orcid.org/0009-0003-3335-1544>, <https://orcid.org/0000-0002-9530-1759>, <https://orcid.org/0000-0003-2080-7955>, <https://orcid.org/0009-0001-2748-2623>

How to cite this article: Belmejdoub, N., Fadel, I., Drider, S., Alaizari, H., Chebabe, M. and Ouhssine, M. (2025). Assessment of Dietary Habits and Sociodemographic Profiles in the Rabat-Salé-Kénitra Region. *Asian Journal of Dairy and Food Research*. 1-9. doi: 10.18805/ajdfr.DRF-498.

Submitted: 13-01-2025 **Accepted:** 21-07-2025 **Online:** 20-08-2025

Also Comprehensive knowledge of nutrition is crucial for individuals to be able to understand and meet their nutritional requirements, allowing them to make

informed food choices (Truman *et al.* 2017; Bany-yasin *et al.* 2023).

In Morocco, demographic transition and epidemiological transition are generally associated with changes in lifestyle, including eating habits and physical activity levels (Aboussaleh and Sbaibi 2015). This transition results in a significant increase in the prevalence of non-communicable diseases (NCDs), including cancer, diabetes, cardiovascular disease, chronic respiratory disease and chronic kidney failure. NCDs are the leading cause of death, estimated at 80% according to the WHO, with an estimated 12% premature mortality. This statistic exceeds the global average of 70% of deaths attributed to NCDs, placing Morocco among the countries of the Eastern Mediterranean region with a high mortality due to these diseases (Africa and Asia, 2018). In the same context, malnutrition is a major public health concern, manifested by a relative or absolute deficit or excess of one or more essential nutrients (WHO) (Azandjeme *et al.*, 2020). Malnutrition from nutrient deficiency, particularly micronutrients such as iron and vitamin A, poses a significant public health challenge, especially in developing countries (Peterson *et al.*, 2016).

Consequently, micronutrient deficiencies represent a significant public health issue, with serious consequences, particularly for vulnerable groups such as pregnant and lactating women, adolescent girls, infants and young children (Reyes Matos *et al.*, 2020). Therefore severe and persistent nutritional deficiency can lead to deficiency disorders, such as goiter (hypertrophy of the thyroid gland due to lack of iodine), rickets (bone deformity due to vitamin D deficiency), or pernicious anemia (a form of anemia accompanied by neurological disorders due to vitamin B12 deficiency) (Knobel 2016). Globally, iodine deficiency affects more than 700 million people, largely in the least developed

countries (WHO/UNICEF/International Council for the Fight against Iodine Deficiency Disorders. Progress towards the elimination of iodine deficiency disorders (Organization 2014). In addition, more than 2 billion people suffer from iron deficiency anemia (WHO/UNICEF/UN University. Evaluation, prevention and control of iron deficiency anemia: a guide for program managers (Denno and Paul., 2017).

Vitamin A deficiency remains the leading cause of preventable and unjustified blindness in children, as well as an increased risk factor for premature infant mortality due to infectious diseases; it affects approximately 250 million children under five years of age in a subclinical form (WHO/UNICEF) (West and Darnton-Hill, 2008). So diet and nutrition play an essential role in maintaining good health throughout life. Their influence as key factors in the development of chronic non-communicable diseases is well established which places them at the forefront of prevention (Di Renzo *et al.*, 2015).

The study examines key concerns regarding dietary habits in the Rabat-Salé-Kénitra region. It highlights a dietary imbalance, with a carbohydrate-dominant diet and insufficient protein and vitamin intake. Health implications include the potential impact of these patterns on BMI and overall health, given low consumption of essential food groups like fish and dairy.

MATERIALS AND METHODS

Study area

Rabat-Salé-Kénitra is a region located in the northwestern part of Morocco. The region is situated between the longitude 5°37'W and 7°34'W and latitude 33°08'N and 35°03'N (Fig 1). It is one of the twelve regions of Morocco since the territorial division of 2015 with total area of 18




Fig 1: Study area.

194 km², it is supervised by several regions of Morocco: it shares its limits in the north with the region of Tanger-Tetouan-Al Hoceima, in the east with the region of Fès-Meknès, in the south with the regions of BeniMellal-Khénifra and Casablanca-Settat and in the west, it is bordered by the Atlantic Ocean, the number of municipalities is 114 including 23 urban and 91 rural. Based on 2014 Census of Population data, the region had a total population of 4,580,866. (El Hachlafi *et al.*, 2020; Berred *et al.*, 2022; Loulad *et al.*, 2023).

Study samples

The study included participants aged 18+ from the Rabat-Salé-Kénitra region, using a two-part questionnaire. The first part collected socio-demographic data, while the second focused on food supplement (FS) consumption, including frequency, duration, reasons, components, forms, and purchase locations. Non-consumers were excluded. Conducted as a cross-sectional prospective study, data collection spanned February 2021 to January 2022 via an anonymous Google Drive questionnaire, and within a subsidiary of DAXEN Morocco (NaturAloé) located in the province of Kénitra over a period of 11 months. Statistical analysis was performed using SPSS 22, with results presented in graphs and an Excel database.

RESULTS AND DISCUSSION

Sociodemographic and anthropometric characteristics

The sociodemographic and anthropometric characteristics of the participants are analyzed with a focus on gender, age, education level, employment status, and family situation.

Table 1 presents the age distribution by gender among 1,012 participants (60.4% females, 39.6% males). Participants were categorized into four age groups: 18-30 years, 31-40 years, 41-50 years, and over 50 years. The largest group was 18-30 years, comprising 45.4% of participants (283 females and 176 males), followed by the 31-40 group with 37.1% (194 females and 181 males). The 41-50 group accounted for 9.3% (64 females and 30 males), while the over 50 group, the smallest cohort, made up 8.3% (70 females and 14 males). Females consistently outnumbered males across all age groups, with the gender gap particularly pronounced among participants over 50 years old. The predominance of younger participants, especially in the 18-30 group, suggests a demographic bias, and male participation declined notably in older age groups. The Chi-square test ($\chi^2 = 32.863$, $p = 0.0005$) confirmed a statistically significant difference in age distribution between genders.

Table 1 summarizes data from 1,012 participants (60.4% females, 39.6% males) distributed across age, education, profession, and family status. Age groups included 18-30 (45.4%), 31-40 (37.1%), 41-50 (9.3%), and over 50 (8.3%) years, with females consistently outnumbering males, particularly in the over-50 category. The Chi-square test ($\chi^2 = 32.863$, $p = 0.0005$) revealed

significant gender differences in age distribution. Educational attainment showed most participants (85.9%) had university education, followed by secondary (12.8%) and primary education (1.3%). A Chi-square test ($p = 0.379$) indicated no significant gender differences in education levels. Professionally, civil servants dominated (52.9%), followed by self-employed (20.7%), students (14.3%), and unemployed (12.2%). The unemployed group was exclusively female. Gender differences in professional status were significant ($p = 0.0005$). Regarding family status, singles constituted 41.3%, married participants 54.8%, and divorced individuals 3.9%. The Chi-square test ($p = 0.982$) found no significant gender differences in family status.

BMI group

Table 2 showed the percentage of the different BMI groups distributed by the sexes, providing a clear overview of the body composition of the study population. Out of a total of 1,012 participants, 60.4% were females (611 participants) and 39.6% were males (401 participants). Participants were grouped into six BMI categories: underweight, normal weight, overweight, and three obesity classes.

The underweight category ($BMI < 18.5$) accounts for 5.2% of the total participants (52 individuals), with 18 females (1.8%) and 34 males (3.4%), showing higher representation among males. The normal weight category ($18.5 \leq BMI \leq 24.9$) is the largest, comprising 38.9% of the participants (394 individuals), with 203 females (20.1%) and 191 males (18.9%), indicating near-equal gender distribution. In the overweight category ($25 \leq BMI \leq 29.9$), 30.6% of the participants (310 individuals) were included, with a higher proportion of 207 females (20.5%) compared to 103 males (10.2%). The Obesity Class I ($30 \leq BMI \leq 34.9$) represents 15.6% of participants (158 individuals), consisting of 98 females (9.7%) and 60 males (5.9%). In Obesity Class II ($35 \leq BMI \leq 39.9$), 7.5% of participants (76 individuals) are represented, including 66 females (6.5%) and 10 males (1.0%), with a strong dominance of females. Finally, Obesity Class III ($BMI > 40$) is the smallest category, accounting for 2.2% of participants (22 individuals), with 19 females (1.9%) and only 3 males (0.3%).

The Chi-square test produced a P-value of 0.0005, indicating a statistically significant relationship between gender and BMI distribution at the 0.05 level.

Frequency of food consumption in both gender

Table 3 examines the eating habits of 1,012 participants (60.4% women, 39.6% men). Substantial meals were most common on weekends, with Saturday (31.6%) and Sunday (27.8%) preferred, particularly among women (Saturday: 19.4%, Sunday: 16.1%) compared to men (Saturday: 12.3%, Sunday: 11.7%). Weekdays like Tuesday (2.0%) and Thursday (1.3%) showed lower proportions. The Chi-square test ($p = 0.163$) found no significant gender differences in meal timing.

Regarding meal frequency, 53.6% ate four meals daily, followed by three (19.6%) and two (18.5%) meals, with no

significant gender variation ($p = 0.053$). Daily bread or pastry consumption was reported by 80.7%, equally across genders ($p > 0.05$). Fruits and vegetables were consumed 4–5 times weekly by 60.2% and daily by 23.0%, with no gender differences ($p = 0.543$), this result differs slightly from that reported in a study conducted in the Arwal district of Bihar, where 87.01% of participants stated that they consumed green vegetables every day (Krishnasree *et al.* 2018; Sharma 2022). However, meat and poultry consumption (4–5 times weekly) showed significant differences ($p = 0.0005$),

with women consuming more (31.6% vs. 13.9%). Other categories showed no significant differences. Overall, eating habits were largely similar between genders, with minor variations in specific categories.

This aligns with Morocco's cereal-centric food culture, where bread, often paired with Tagine, replaces traditional grains like barley. Meanwhile, couscous consumption declines, while pasta rises (TOBBA Tassabih, 2023). The Moroccan Nutritional Transition study highlighted the fact that the Moroccan diet has changed considerably, with an

Table 1: Distribution of Sociodemographic and anthropometric characteristics.

			Gender		Total	
			Female	Male		
Age	18-30 years	Effective	283	176	459	
		% of total	28.0%	17.4%	45.4%	
	31-40 years	Effective	194	181	375	
		% of total	19.2%	17.9%	37.1%	
	41-50 years	Effective	64	30	94	
		% of total	6.3%	3.0%	9.3%	
	over 50 years	Effective	70	14	84	
		% of total	6.9%	1.4%	8.3%	
			Gender	Total	Chi-square test	
			Female	Male		P value
Education levels	Primary	Effective	6	7	13	1.940
		% of total	0.6%	0.7%	1.3%	P=0.379
	Secondary	Effective	74	56	130	
		% of total	7.3%	5.5%	12.8%	
	University	Effective	531	338	869	
		% of total	52.5%	33.4%	85.9%	
			Gender	Total	Tests du	
			Female	Male		Khi-deux
Professional status	Student	Effective	82	63	145	96.132
		% of total	8.1%	6.2%	14.3%	P=0.0005
	Civil servant	Effective	304	231	535	
		% of total	30.0%	22.8%	52.9%	
	Self-employed	Effective	102	107	209	
		% of total	10.1%	10.6%	20.7%	
	Unemployed	Effective	123	0	123	
		% of total	12.2%	0.0%	12.2%	
			Gender	Total	Tests du	
			Woman	Man		Khi-deux
Family status	Single	Effective	253	165	418	0.036
		% of total	25.0%	16.3%	41.3%	P=0.982
	Married	Effective	335	220	555	
		% of total	33.1%	21.7%	54.8%	
	Divorced	Effective	23	16	39	
		% of total	2.3%	1.6%	3.9%	
Total	Effective	611	401	1012		
	% of total	60.4%	39.6%	100.0%		

increase in the consumption of animal products, while that of cereals and sugar has remained relatively high, underlining the specificity of Moroccan eating habits (Benjelloun, 2002).

Fruits and vegetables are consumed 4-5 times weekly by 83.2% of Moroccans, but daily intake averages one fruit and 2.5 vegetable servings, below the recommended 400-500 g for cardiovascular health (Organization, 2003). Meat and poultry consumption, mainly 2-5 times weekly, reflects dietary preferences like mutton, despite rising concerns about antibiotic resistance and contamination in poultry products (Fazza *et al.*, 2023; Cohen *et al.*, 2007). Morocco's nutritional transition shows increased animal product intake alongside under-nutrition in children and growing adult obesity (Benjelloun, 2002). Compared to France, where meat ranks second, Moroccans prioritize cereals and vegetables (Sharma, 2021). A study conducted in Hyderabad shows that chicken is the preferred meat, while mutton is less favored, and pork and beef are consumed occasionally due to taboos and religious beliefs, with average consumption levels falling below recommended dietary guidelines (Ankur Ojha *et al.*, 2015; Krishnasree *et al.*, 2018; Sharma, 2022).

Traditional Moroccan dairy products like *lben*, *smen*, and *jben* are vital in rural diets and increasingly marketed informally (Cuvelier *et al.*, 2004). Goat's milk is primarily processed into cheese due to sensory preferences (EFFICACE 2021). These dairy products may protect against colorectal cancer (Kaoutar *et al.* 2019). Among legumes, peas, lentils, white beans and broad beans are the most common. A study analyzed these products for energy, water, protein, fat, total carbohydrates, total ash and 15 minerals, revealing that the energy value of Moroccan foods is higher than that reported in the FAO table for Africa (FAO *et al.*, 2017). Morocco is the leading North African country in terms of cereal production. However, little of this biomass is used for energy production in the

country, representing a major challenge due to heavy dependence on energy imports, limited water resources and a rapidly growing population (Mahdavi *et al.*, 2023). One study explored how the cognitive representation of a food as a "snack" or "meal" influences eating behavior. Foods such as potato chips, crackers, cookies and nuts are seen as snacks, while soups, burritos, pizzas and pancakes are seen as meals. (Wadhera and Capaldi 2012). Dans notre étude la majorité des individus de l'échantillon étudié (67,1%) consomment les aliments type snack (Pizza, Paninis, Hamburgers, Tacos...) d'une fréquence faible qui est d'une seule fois par semaine ce qui est compatible aux habitudes des français, selon (Le snacking en France) 57% fréquentent au moins une fois dans la semaine (Forbes *et al.*, 2016). (including weekends). In contrast to America, almost three quarters of survey respondents (73%) reported consuming snacks at least once a day. Specifically, 34% reported snacking once a day, 25% twice, and 14% three or more times, demonstrating that snacking is an integral part of the daily diet of many in the American population (Wadhera and Capaldi 2012). One study found that high consumption of fast food and industrial baked goods may be associated with an increased risk of depression. The authors recommend limiting consumption of these foods in favor of healthier options (Gougeon, 2014).

This consumption is similar to that observed in a study conducted among 100 subjects in Hisar city of Haryana state, where the intake of pulses was found to be adequate at 69% and 57% of the recommended intake, respectively (Budhwar and Jood, 2010). Snacking, primarily weekly, aligns with French trends (Forbes *et al.*, 2016) but contrasts with frequent American habits (Wadhera and Capaldi, 2012). High fast-food consumption may increase depression risk, highlighting the need for healthier choices (Gougeon, 2014).

Table 2: Distribution of BMI groups by gender.

			Gender		Total	Chi-square test
			Female	Male		
Groupe BMI	BMI <18.5	Effective	18	34	52	61.279 P=0.0005
		% of total	1.8%	3.4%	5.2%	
	18.5< BMI >24.9	Effective	203	191	394	
		% of total	20.1%	18.9%	38.9%	
	25< BMI >29.9	Effective	207	103	310	
		% of total	20.5%	10.2%	30.6%	
	30< BMI >34.9	Effective	98	60	158	
		% of total	9.7%	5.9%	15.6%	
	35< BMI >39.9	Effective	66	10	76	
		% of total	6.5%	1.0%	7.5%	
	BMI >40	Effective	19	3	22	
		% of total	1.9%	0.3%	2.2%	
Total	Effective	611	401	1012		
	% of total	60.4%	39.6%	100.0%		

Table 3: Frequency of food consumption in both gender.

			Sexe		Total	Chi-square test
			Female	Male		
Meal consequences	Monday	Effective	21	28	49	10.488 P=0.163
		% of total	2.1%	2.8%	4.8%	
	Thursday	Effective	11	9	20	
		% of total	1.1%	0.9%	2.0%	
	Wednesday	Effective	12	5	17	
		% of total	1.2%	0.5%	1.7%	
	Tuesday	Effective	7	6	13	
		% of total	0.7%	0.6%	1.3%	
	Friday	Effective	105	60	165	
		% of total	10.4%	5.9%	16.3%	
	Saturday	Effective	196	124	320	
		% of total	19.4%	12.3%	31.6%	
	Sunday	Effective	163	118	281	
		% of total	16.1%	11.7%	27.8%	
	Aucun	Effective	96	51	147	
		% of total	9.5%	5.0%	14.5%	
Number of meals per day	1	Effective	18	10	28	9.354 P=0.053
		% of total	1.8%	1.0%	2.8%	
	2	Effective	118	69	187	
		% of total	11.7%	6.8%	18.5%	
	3	Effective	112	86	198	
		% of total	11.1%	8.5%	19.6%	
	4	Effective	338	204	542	
		% of total	33.4%	20.2%	53.6%	
	5	Effective	25	32	57	
		% of total	2.5%	3.2%	5.6%	
Consumption of bread/wholemeal bread and pastries	Never	Effective	14	7	21	0.609 P= 0.894
		%	2.3%	1.7%	2.1%	
	2-3 times per week	Effective	47	34	81	
		%	7.7%	8.5%	8.0%	
	4-5 times per week	Effective	55	38	93	
		%	9.0%	9.5%	9.2%	
	Dialy	Effective	495	322	817	
		%	81.0%	80.3%	80.7%	
Consumption of Fruit and vegetable	Never	Effective	20	9	29	2.145 P=0.543
		%	2.0%	0.9%	2.9%	
	2-3 times per week	Effective	85	56	141	
		%	8.4%	5.5%	13.9%	
	4-5 times per week	Effective	373	236	609	
		%	36.9%	23.3%	60.2%	
	Dialy	Effective	133	100	233	
		%	13.1%	9.9%	23.0%	
Consumption of meat and poultry	Never	Effective	40	46	86	32.480 P= 0.0005
		%	4.0%	4.5%	8.5%	
	2-3 times per week	Effective	51	54	105	
		%	5.0%	5.3%	10.4%	
	4-5 times per week	Effective	192	152	344	
		%	19.0%	15.0%	34.0%	
	Dialy	Effective	320	141	461	

Table 3: Continue....

Table 3: Continue....

		%	31.6%	13.9%	45.6%	
	Never	Effective	8	8	16	
		%	0.8%	0.8%	1.6%	
Consumption of	Never	Effective	26	13	39	0.676
fish and seafood		%	2.6%	1.3%	3.9%	P=0.713
	1 times per week	Effective	503	333	836	
		%	49.7%	32.9%	82.6%	
	2-3 times per week	Effective	82	55	137	
		%	8.1%	5.4%	13.5%	
	Never	Effective	55	27	82	7.079
		%	5.4%	2.7%	8.1%	P=0.132
Consumption of and	2-3 times per week	Effective	146	108	254	
Cereals (pasta, rice,		%	14.4%	10.7%	25.1%	
etc.) legumes (white	4-5 times per week	Effective	295	194	489	
beans, chickpeas, lentils,		%	29.2%	19.2%	48,3%	
	Dialy	Effective	23	6	29	
		%	2.3%	0.6%	2.9%	
	Never	Effective	92	66	158	
		%	9.1%	6.5%	15.6%	
	Never	Effective	24	17	41	1.825
		%	2.4%	1.7%	4.1%	P=0.768
Produits sucrés	2-3 times per week	Effective	15	12	27	
(chocolats, confiseries,		%	1.5%	1.2%	2.7%	
gâteaux, biscuits	4-5 times per week	Effective	43	28	71	
sucrés, pâtisseries)		%	4.2%	2.8%	7.0%	
	Dialy	Effective	233	166	399	
		%	23.0%	16.4%	39.4%	
	Never	Effective	296	178	474	
		%	29.2%	17.6%	46.8%	
	Never	Effective	17	19	36	3.107
		%	1.7%	1.9%	3.6%	P= 0.540
Consumption of	2-3 times per week	Effective	411	268	679	
Snack foods (Pizza,		%	40.6%	26.5%	67.1%	
Paninis, Hamburgers,	4-5 times per week	Effective	143	90	233	
Tacos, etc.)		%	14.1%	8.9%	23.0%	
	Dialy	Effective	38	22	60	
		%	3.8%	2.2%	5.9%	
	Never	Effective	2	2	4	
		%	0.2%	0.2%	0.4%	

CONCLUSION

This study analyzed dietary habits in Rabat-Salé-Kénitra, revealing a young, educated population. Bread was the most consumed food, followed by fruits, vegetables, and meat. Dairy was moderately consumed, while fish, legumes, and snacks were less frequent. No significant gender differences were found, highlighting a diet leaning towards carbohydrates over proteins.

Disclaimers

The views and conclusions expressed in this article are solely those of the authors and do not necessarily represent

the views of their affiliated institutions. The authors are responsible for the accuracy and completeness of the information provided, but do not accept any liability for any direct or indirect losses resulting from the use of this content.

Conflict of interest

The authors declare that there are no conflicts of interest regarding the publication of this article.

REFERENCES

Aboussaleh, Y. and Sbaibi, R. (2015). Etat Nutritionnel de la population du Nord-Ouest du Maroc. *Antropo*. **33**: 13-20.

- Africa, N.A. and Asia, S.E. (2018). Global leishmaniasis surveillance, 2017–2018 and first report on 5 additional indicators. *Global Health*. **3(40)**: 530-540.
- Albrahim, T., Alsalaan, R., Alhusan, S. I., Alrasheedi, K.R., Aldosari, H.A. and Albargi, H.M. (2023). Exploring the association between nutritional knowledge and fruit and vegetable consumption among young adults: A cross-sectional study. *Current Research in Nutrition and Food Science Journal*. **11(2)**: 617-626.
- Ankur Ojha, A.O., Akriti Sharma, A.S., Manvesh Sihag, M.S. and Seema Ojha, S.O. (2015). Food packaging-materials and sustainability-a review. <https://www.cabidigitallibrary.org/doi/full/10.5555/20163111311>.
- Azandjeme, C.S., Alihonou, F., Sossa, C.J., Gbatcho, U. and Gounongbe, F. (2020). Factors associated with the nutritional status of schoolchildren in the main city of Benin Republic, sub-Saharan Africa. *Int Arch Public Health Community Med*. **4**: 036. <https://pdfs.semanticscholar.org/60a7/99e8531732a805a0b328d6627a9d1e9a1829.pdf>
- Bany-yasin, H., Elmor, A.A., Ebrahim, B.K., Ahmed, A.A.M., Alarachi, M.R., Abedalqader, L., Amer, R., Alyousef, A.M.S., Alhajeh, Y.F., Alyoussef, A., Eid, H.A.M.A., Elsayed, M.M., Desouky, E.D.E., Salem, H.K. and Salem, M.R. (2023). Exploration of the nutrition knowledge among general population: Multi-national study in Arab countries. *BMC Public Health*. **23(1)**: 1178. <https://doi.org/10.1186/s12889-023-15791-9>.
- Benjelloun, S. (2002). Nutrition transition in Morocco. *Public Health Nutrition*. **5(1a)**: 135-140.
- Berred, K., Youssi, M.E. and Berred, S. (2022). Inventorying rabat-salé-kénitra region's geological heritage within central morocco: A useful tool for developing regional geotourism activity. *Geoheritage*. **14(3)**: 91. <https://doi.org/10.1007/s12371-022-00723-7>.
- Budhwar, S. and Jood, S. (2010). Nutritional Status of Subjects of Hisar City Suffering from Non-Insulin Dependent Diabetes Mellitus. *Journal of Dairying, Foods and Home Sciences*. **29(1)**: 25-31.
- Cohen, N., Ennaji, H., Bouchrif, B., Hassar, M. and Karib, H. (2007). Comparative study of microbiological quality of raw poultry meat at various seasons and for different slaughtering processes in Casablanca (Morocco). *Journal of Applied Poultry Research*. **16(4)**: 502-508.
- Cuvelier, C., Cabaraux, J.F., Dufrasne, I., Hornick, J.L. and Istasse, L. (2004). Acides gras: Nomenclature et sources alimentaires. *Annales de Médecine Vétérinaire*. **148(3)**.
- Denno, D. M. and Paul, S. L. (2017). Child health and survival in a changing world. *Pediatric Clinics*. **64(4)**: 735-754.
- Di Renzo, L., Colica, C., Carraro, A., Cenci Goga, B., Marsella, L.T., Botta, R., Colombo, M.L., Gratteri, S., Chang, T.F.M., Drolli, M., Sarlo, F. and De Lorenzo, A. (2015). Food safety and nutritional quality for the prevention of non communicable diseases: The Nutrient, hazard Analysis and Critical Control Point process (NACCP). *Journal of Translational Medicine*. **13(1)**: 128. <https://doi.org/10.1186/s12967-015-0484-2>.
- Efficace, A.S.E. (2021). Compléments Alimentaires Et Vitalité Sexuelle: La Phytothérapie Est-Elle Une.
- El Hachlafi, N., Chebat, A., Bencheikh, R.S. and Fikri-Benbrahim, K. (2020). Ethnopharmacological study of medicinal plants used for chronic diseases treatment in Rabat-Sale-Kenitra region (Morocco). *Ethnobotany Research and Applications*. **20**: 1–23.
- FAO, F., OMS, P. and Unicef. (2017). L'État de la sécurité alimentaire et de la nutrition dans le monde 2017. Renforcer La Résilience Pour Favoriser La Paix et La Sécurité Alimentaire.
- Fazza O, Favard Ennachachibi M, Ennassiri H, Hmyene A (2023) Antibiotic susceptibility of β -Glucuronidase-positive *Escherichia coli* Isolated from poultry products in morocco. *Int J Food Sci*. **2**: 55-68.
- Fieldhouse, P. (2013). Food and nutrition: Customs and culture. Springer.
- Forbes, S. L., Kahiya, E. and Balderstone, C. (2016). Analysis of snack food purchasing and consumption behavior. *Journal of Food Products Marketing*. **22(1)**: 65-88. <https://doi.org/10.1080/10454446.2014.949992>.
- Gougeon, L. (2014). Nutritional predictors of depression in a cohort of community-dwelling elderly Canadians: NuAge cohort.
- Gowshall, M. and Taylor-Robinson, S. (2018). The increasing prevalence of non-communicable diseases in low-middle income countries: The view from Malawi. *International Journal of General Medicine*. **11**: 255-264. <https://doi.org/10.2147/IJGM.S157987>.
- Hercberg, S. (2011). Le Programme National Nutrition Santé (PNNS): Un vrai programme de santé publique. *Cahiers de Nutrition et de Diététique*. **46(2)**: S5–S10.
- Kaoutar, K., Hilali, M.K. and Loukid, M. (2019). Alimentation et comportements alimentaires chez les adolescents scolarisés de la ville de Marrakech (Maroc). *Antropo*. 42.
- Knobel, M. (2016). Etiopathology, clinical features and treatment of diffuse and multinodular nontoxic goiters. *Journal of Endocrinological Investigation*. **39(4)**: 357-373. <https://doi.org/10.1007/s40618-015-0391-7>.
- Krishnasree, V., Nethra, P.V., Dheeksha, J., Madumitha, M.S., Vidyaswari, R. and Lakshya, P. (2018). A pilot study on assessing the sustainability of food safety and hygienic practices in street food handling system. *Asian Journal of Dairy and Food Research*. **37(4)**: 321-325. doi: 10.18805/ajdfr.DR-1381.
- Lecerf, J.-M. (2018). Bilan du Programme National Nutrition Santé (PNNS) 3, et perspectives pour le suivant. *Médecine Des Maladies Métaboliques*. **12(1)**: 46-48.
- Loulad, S., Nguyen, T.T., Simou, M.R., Rhinane, H. and Buerkert, A. (2023). Monitoring rural-urban transformation in the coastal region of Rabat-Sale-Kenitra, Morocco. *Plos One*. **18(8)**: e0290829.
- Mahdavi, M., Jurado, F., Vera, D. and Schmitt, K. (2023). Potential of Moroccan Agriculture Sector for Energy Generation from Barley Straws. 2023 IEEE IAS Global Conference on Emerging Technologies (GlobConET). 1-6.
- O'Neil, A., Quirk, S.E., Housden, S., Brennan, S. L., Williams, L.J., Pasco, J.A., Berk, M. and Jacka, F.N. (2014). Relationship between diet and mental health in children and adolescents: A Systematic Review. *American Journal of Public Health*. **104(10)**: e31-e42.

- Organization, W.H. (2003). Régime alimentaire, nutrition et prévention des maladies chroniques: Rapport d'une consultation OMS/FAO d'experts. Organisation mondiale de la Santé.
- Organization, W.H. (2014). Salt reduction and iodine fortification strategies in public health: Report of a joint technical meeting convened by the World Health Organization and The George Institute for Global Health in collaboration with the International Council for the Control of Iodine Deficiency Disorders Global Network, Sydney, Australia, March 2013.
- Pandey, S., Budhathoki, M. and Yadav, D.K. (2021). Psychosocial determinants of vegetable intake among Nepalese young adults: An exploratory survey. *Frontiers in Nutrition*. **8**: 688059.
- Passi, S. J. (2017). Prevention of non-communicable diseases by balanced nutrition: Population-specific effective public health approaches in developing countries. *Current Diabetes Reviews*. **13(5)**: 461-476.
- Penzavecchia, C., Todisco, P., Muzzioli, L., Poli, A., Marangoni, F., Poggiogalle, E., Giusti, A. M., Lenzi, A., Pinto, A. and Donini, L.M. (2022). The influence of front-of-pack nutritional labels on eating and purchasing behaviors: A narrative review of the literature. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity*. **27(8)**: 3037-3051. <https://doi.org/10.1007/s40519-022-01507-2>.
- Perico, N. and Remuzzi, G. (2012). Chronic kidney disease: A research and public health priority. *Nephrology Dialysis Transplantation*. **27(suppl_3)**, iii19–iii26.
- Peterson, L.A., Cheskin, L.J., Furtado, M., Papas, K., Schweitzer, M.A., Magnuson, T.H. and Steele, K. E. (2016). Malnutrition in bariatric surgery candidates: Multiple micronutrient deficiencies prior to surgery. *Obesity Surgery*. **26(4)**: 833-838. <https://doi.org/10.1007/s11695-015-1844-y>.
- Reyes Matos, U., Mesenburg, M.A. and Victora, C.G. (2020). Socioeconomic inequalities in the prevalence of underweight, overweight and obesity among women aged 20–49 in low-and middle-income countries. *International Journal of Obesity*. **44(3)**: 609-616.
- Sharma, P. (2021). Quelle «souveraineté» dans la souveraineté alimentaire? Le cas népalais1. *Rédaction et Administration*. **28**: 87.
- Sharma, R. (2022). A bibliometric study of "Bhartiya Krishi Anushandhan Patrika" from 2016 to 2021. *Bhartiya Krishi Anusandhan Patrika*. **37(4)**: 311-315. doi: 10.18805/BKAP506.
- TOBBA Tassabih, G.H. (2023). Enquête alimentaire sur la consommation des compléments alimentaire chez les algériens à la willaya de Tébessa [PhD Thesis, Université Echahid Chikh Larbi Tébessi-Tébessa]. <http://dspace.univ-tebessa.dz:8080/jspui/handle/123456789/10670>.
- Truman, E., Lane, D. and Elliott, C. (2017). Defining food literacy: A scoping review. *Appetite*. **116**: 365-371.
- Wadhera, D. and Capaldi, E. D. (2012). Categorization of foods as "snack" and "meal" by college students. *Appetite*. **58(3)**: 882-888.
- West, K.P. and Darnton-Hill, I. (2008). Vitamin A Deficiency. In R.D. Semba, M.W. Bloem and P. Piot (Eds.), *Nutrition and Health in Developing Countries*. Pp: 377-433. Humana Press. https://doi.org/10.1007/978-1-59745-464-3_13.