



Effect of Conjoint Application of Inorganic and Organic Nutrient Sources on Yield and Economics of French Bean (*Phaseolus vulgaris* L.)

Kuldeep S. Thakur, Nitin Yadav, Sandeep Kansal¹

10.18805/LR-4979

ABSTRACT

Background: French bean is an important legume vegetable crop. Due to its low ability to fix atmospheric nitrogen, excess amount of nitrogenous fertilizers are required by this crop as compared to other legumes. Overwhelming application of inorganic inputs has severely impacted environment and soil. So, there is a need to reduce inorganic inputs by adding organic nutrient sources.

Methods: A three-year field experiment was laid comprising of eight treatment combinations and was replicated thrice during *Kharif* season of 2018, 2019 and 2020. Seeds of French bean cv. Green Wonder were sown at a spacing of 45 × 15 cm in each plot of size 3.0 m × 1.8 m. Recommended dose of fertilizers (NPK 50:100:50 kg ha⁻¹) was followed as per different treatment details.

Result: Application of 75 per cent NPK through inorganic and 25 per cent N through vermicompost resulted in minimum days to first pod harvest (61.89 days), maximum pod width (7.67 mm), number of pods per plant (30.96), plant height (51.29 cm), average pod weight (5.70 g), marketable pod yield (130.53 q ha⁻¹) and highest benefit cost ratio (2.18) along with maximum net returns (Rs. 313061.43 ha⁻¹).

Key words: Economics, Growth, INM, Inorganic, Organic, Vermicompost.

INTRODUCTION

French bean (*Phaseolus vulgaris* L.) belongs to family Fabaceae with chromosome number 2n = 14 is one of the most important vegetable crops and is widely cultivated legume globally. It is a short durational highly relished vegetable crop of North India. It has very high nutritional value containing 20.69 to 25.81 per cent crude protein, 1.72 per cent fats, 72.42 per cent carbohydrates and 5.83 mg of iron (Singh *et al.*, 2014). French bean fetches premium price in market as compared to other vegetables and is a popular vegetable grown under irrigated conditions almost throughout the year. It is of great importance due to its high production potential as well as high nutritive value. French bean is a tender warm season vegetable which cannot tolerate frost, high temperature and rainfall. Its seeds do not germinate below 15°C and a most favorable soil temperature for its seed germination ranges from 18-24°C. A mean air temperature of 20-25°C is optimum for its growth and high pod yield. It is grown for tender green pods for fresh consumption as well as for dry seeds which are used as pulse. The dried beans are rich in protein and closely compare with meat. In most of the tropical Asia, it is a major vegetable crop where indigenous pulses are preferred. In India, it is mostly grown for tender green pods, while in the USA, it is grown for processing in large quantities (Meena *et al.*, 2017). The crop can be grown all over the world with diverse environmental conditions ranging from tropical to temperate (Kumar *et al.*, 2021). Insufficient, late appearance and ineffectiveness of nodules are subjected to low nitrogen fixation ability of this crop. Due to this, higher dose of nitrogenous fertilizer is applied in French bean as compared to other legumes. Today's intensive agriculture is based on

Department of Vegetable Science, Dr. Y.S. Parmar University of Horticulture and Forestry, Solan-173 230, Himachal Pradesh, India.

¹Department of Plant Pathology, Dr. Y.S. Parmar University of Horticulture and Forestry, Solan-173 230, Himachal Pradesh, India.

Corresponding Author: Nitin Yadav, Department of Vegetable Science, Dr. Y.S. Parmar University of Horticulture and Forestry, Solan-173 230, Himachal Pradesh, India.

Email: nitinydv1900@gmail.com

How to cite this article: Thakur, K.S., Yadav, N. and Kansal, S. (2022). Effect of Conjoint Application of Inorganic and Organic Nutrient Sources on Yield and Economics of French Bean (*Phaseolus vulgaris* L.). Legume Research. DOI: 10.18805/LR-4979.

Submitted: 06-06-2022 **Accepted:** 21-07-2022 **Online:** 16-08-2022

use of chemical fertilizers, but price of these fertilizers has gone up considerably, which in turn increased the production cost and also decreased the fertility status of the soil markedly. Application of inorganic inputs has increased the productivity of produce, but has severely impacted environment and soil health (Yadav *et al.*, 2022). Integration of various sources of nutrients is more suitable because this reduces the application of chemical fertilizers and cost of cultivation, besides being an environment friendly approach (Ram and Mir, 2006). There is an ample scope for increasing productivity through application of chemical fertilizers in conjunction with organic nutrient sources that reduces the load of chemical fertilizers and maintains soil sustainability (Saikia *et al.*, 2018). So, present investigation was designed to evaluate the effect of conjoint application of inorganic and organic nutrient sources on yield and economics of French bean.

MATERIALS AND METHODS

This investigation was carried out during the *Kharif* season of 2018, 2019 and 2020 at Vegetable Research Farm, Department of Vegetable Science, Dr. Y S Parmar University of Horticulture and Forestry, Nauni, Solan (HP) for three consecutive years to evaluate the effect of integrated nutrients in French bean. The experiment comprised of eight treatments in Randomized Block Design (RBD) and was replicated thrice. Treatment details are presented in Table 1. The experimental site is located at an altitude of 1,270 meters above mean sea level lying between latitude of 30°5' North and longitude of 77°11' East. The area receives an annual rainfall of about 1100 mm and most of which is received during the monsoon period (mid June - mid September).

Seeds of French bean cv. Green Wonder were sown at a spacing of 45 cm × 15 cm in each plot of size 3.0 m × 1.8 m. Recommended dose of fertilizers *i.e.* NPK 50:100:50 kg ha⁻¹ (Anonymous, 2020) was adopted and applied as per treatment details presented in Table 1. Organic manures *i.e.* FYM and vermicompost were applied to different treatments on N-equivalent basis. Before sowing of seeds, soil was low in organic carbon (1.01%), normal in available nitrogen (322.16 kg ha⁻¹) and available potassium (314.19 ka ha⁻¹),

high in available phosphorus (52.45 kg ha⁻¹), normal pH (6.93) and electrical conductivity (0.43 dS m⁻¹). Soil properties before sowing were estimated by the methods given by Walkey and Black, 1934 (organic carbon); Subbiah and Asija, 1956 (nitrogen); Merwin and Peech, 1951 (potassium); Olsen *et al.*, 1954 (phosphorus); Jackson, 1967 (pH and EC). Data of three consecutive years was pooled and analysed by using OPSTAT developed by Sheoran *et al.*, 1998.

RESULTS AND DISCUSSION

On the basis of three years pooled data as depicted in Table 2, results revealed that majority of the growth, yield and yield components of French bean were significantly influenced by integrated nutrient management practices. All the growth, yield and yield contributing characters were superior over control.

Significantly minimum days to first pod harvest (61.89 days), maximum pod width (7.67 mm), number of pods per plant (30.96), plant height (51.29 cm), average pod weight (5.70 g) and marketable pod yield (130.53 q ha⁻¹) was recorded with the application of 75 per cent NPK through inorganic and 25 per cent N through Vermicompost (T₃) whereas, maximum pod length (15.89 cm) was recorded with the application of 50 per cent NPK through inorganic and 50 per cent N through vermicompost (T₅). The results indicate the superiority of inorganic manures in conjunction with organic manure over sole application of inorganic manure. Furthermore, vermicompost performed better than FYM in conjunction with inorganic manures.

Application of NPK through inorganic fertilizer in conjunction with organic manure resulted in enhanced photosynthetic activity, cell division, elongation and differentiation that ultimate led to increase in growth attributes. Enhanced plant growth attributes may be due to high composition of nitrogen in inorganic fertilizers along with vermicompost that reduced the leaching loss of nutrients which supplemented to the plant's vegetative phase. Moreover, vermicompost stimulates the microbial activity of soil, maintains soil temperature, increases soil porosity and

Table 1: Details of treatment.

Treatment	Details
T ₁	100% NPK through inorganic
T ₂	75% NPK through inorganic + 25 % N through FYM
T ₃	75% NPK through inorganic + 25 % N through vermicompost
T ₄	50% NPK through inorganic + 50 % N through FYM
T ₅	50% NPK through inorganic + 50 % N through vermicompost
T ₆	25% NPK through inorganic + 75 % N through FYM
T ₇	25% NPK through inorganic + 75 % N through vermicompost
T ₈	Control (no fertilizer)

Table 2: Growth, yield and yield contributing characters as influenced by integrated nutrient management in French bean during 2018, 2019 and 2020 (pooled data).

Treatments	Days to first pod harvest	Pod length (cm)	Pod width (mm)	Number of pods per plant	Plant height (cm)	Average pod weight (g)	Marketable pod yield (q ha ⁻¹)
T ₁	62.22	15.31	7.36	27.19	48.42	5.46	110.62
T ₂	64.22	14.86	7.39	28.47	47.44	5.33	117.53
T ₃	61.89	15.75	7.67	30.96	51.29	5.70	130.53
T ₄	64.11	15.05	7.34	27.16	48.71	5.01	108.15
T ₅	62.56	15.89	7.74	27.89	48.96	5.56	117.53
T ₆	66.22	14.68	7.04	28.07	41.79	5.20	104.03
T ₇	64.00	14.88	7.43	26.14	48.44	5.31	112.76
T ₈	69.22	13.89	7.12	23.99	40.66	4.90	94.65
CD (5%)	3.81	1.02	0.38	2.6	3.12	NS	6.99
CV (%)	5.35	5.82	4.93	8.36	7.76	8.82	7.53

infiltration of water, improves nutrient content and increases growth, yield and quality of the plant (Arora *et al.*, 2011). The increased growth with substitution of N by organic manures might be due to the fact that organic manures release nutrients slowly (Sharma *et al.*, 2015), resulting in enhanced nutrient use efficiency (Verma *et al.*, 2015; Mohanty *et al.*, 2017) and availability of micro nutrients (Nawalgatti *et al.*, 2009; Shubhashree *et al.*, 2011). Higher yield of treatment comprising of inorganic fertilizers in combination with organic manures may be due to the increased availability and uptake of macro and micro nutrients by the plant, resulting in higher rate of physiological (Barcchiya and Kushwah, 2017) and anabolic processes (Sen *et al.*, 2006; Ramana *et al.*, 2010). Sharma *et al.*, 2018 reported increased uptake of N, P and K by French bean significantly through the combined application of vermicompost and N over the use of farmyard manure and nitrogen application. The results are in consonance with Kumar *et al.*, 2009.

Economics of different treatments has been presented in Table 3. Lowest cost of cultivation (Rs. 122935 ha⁻¹) was recorded under control. It was due to no cost incurred for organic manure and their transportation, spreading and application. Maximum cost of cultivation (Rs. 169895 ha⁻¹) was recorded with the application of 25 per cent NPK through inorganic + 75 per cent N through Vermicompost (T₇). Maximum gross income (Rs. 456872.43 ha⁻¹) was recorded with the application of 75 per cent NPK through inorganic + 25 per cent N through Vermicompost (T₃) whereas, minimum (Rs. 331275.72 ha⁻¹) was recorded under control. Highest net income (Rs. 313061.43 ha⁻¹) and benefit cost ratio (2.18) was recorded with the application of 75 per cent NPK through inorganic and 25 per cent N through Vermicompost (T₃) whereas, minimum values (Rs. 208340.72 ha⁻¹ and 1.32) of these were recorded in control and with the application of 25% NPK through inorganic + 75% N through Vermicompost (T₇) respectively. This defines the practical utility of this treatment for the benefit of growers of French bean.

Our findings of various growth and yield components reveals that fertilizer requirement of French bean when met through 75 per cent NPK through inorganic and 25 per cent

N through vermicompost resulted in enhanced yield along with highest benefit cost ratio. It may be due to the potential of organic nutrient sources to influence both the growth as well as yield components as compared to the sole application of (100% NPK) through inorganic source.

CONCLUSION

Vermicompost performed better over FYM in the conjoint application of inorganic and organic nutrient sources. It can be concluded that the application of 75 per cent NPK through inorganic and 25 per cent N through vermicompost resulted in enhanced growth, yield and yield components of French bean along with highest net returns and benefit cost ratio.

ACKNOWLEDGEMENT

The authors are highly thankful to Indian Council of Agricultural Research for providing grants to carry out this study.

Conflict of interest: None.

REFERENCES

- Anonymous. (2020). Package of Practices of Vegetable Crops. Directorate of Extension Education, Dr. Y S Parmar University of Horticulture and Forestry, Solan, Himachal Pradesh, 43p.
- Arora, V.K., Singh, C.B., Sidhu, A.S. and Thind, S.S. (2011). Irrigation, tillage and mulching effects on soybean yield and water productivity in relation to soil texture. *Agricultural Water Management*. 98(4): 563-568.
- Barcchiya, J. and Kushwah, S.S. (2017). Influence of integrated nutrient management on growth, yield parameters and yield in French bean (*Phaseolus vulgaris* L.). *Legume Research*. 40: 920-923.
- Jackson, M.L. (1967). *Soil Chemical Analysis*. Prentice Hall of India Private Limited. New Delhi. pp. 111-126.
- Kumar, R.P., Singh, O.N., Singh, Y., Dwivedi, S. and Singh, J.P. (2009). Impact of integrated nutrient management on growth, yield and economics of French bean (*Phaseolus vulgaris* L.). *Indian Journal of Agricultural Sciences*. 79: 122-128.
- Kumar, U., Prasad, K., Prasad, S.S. and Sinha, B.M. (2021). Integrated nutrient management studies for higher productivity of French bean (*Phaseolus vulgaris* L.) in calcareous soil of Bihar, India. *Legume Research*. LR-4399: 1-6.
- Meena, J., Dhillon, T.S., Meena, A. and Singh, K.K. (2017). Studies on performance of French bean (*Phaseolus vulgaris* L.) genotypes for yield and quality traits under protected conditions. *Plant Archives*. 17(1): 615-619.
- Merwin, H.D. and Peech, M. (1951). Exchange ability of soil potassium in sand, silt and clay fraction as influenced by the nature and complementary exchangeable cations. *Soil Science Society of America Journal*. 15(C): 125-128.
- Mohanty, S., Sahu, G.S., Dash, S.K., Pradhan, S.R., Mangaraj S. and Nahak, S. (2017). Integrated nutrient management for seed production in French bean (*Phaseolus vulgaris* L.). *International Journal of Current Microbiology and Applied Science*. 6: 3295-3303.

Table 3: Economics of integrated nutrient management in French bean.

Treatments	Cost of cultivation (Rs ha ⁻¹)	Gross income (Rs ha ⁻¹)	Net income (Rs ha ⁻¹)	B:C Ratio
T ₁	130770	387160.49	256390.49	1.96
T ₂	136311	411358.02	275047.02	2.02
T ₃	143811	456872.43	313061.43	2.18
T ₄	141863	378518.52	236655.52	1.67
T ₅	156893	411358.02	254465.02	1.62
T ₆	147395	364115.23	216720.23	1.47
T ₇	169895	394650.21	224755.21	1.32
T ₈	122935	331275.72	208340.72	1.69

- Nawalgatti, C.M., Ashwini, G.M., Doddamani, M.B., Chetti, M.B. and Hiremath, S.M. (2009). Influence of organics, nutrients and plant growth regulators on growth, yield and yield components in French bean. *International Journal of Plant Sciences*. 4: 367-372.
- Olsen, S.R., Cole, C.V., Watenable, D.S. and Dean, L.A. (1954). Estimation of available phosphorus in soils by extraction with sodium bicarbonate. US Department of Agriculture Circular, Washington. 939p.
- Ram, T. and Mir, M.S. (2006). Effect of integrated nutrient management on yield and yield-attributing characters of wheat (*Triticum aestivum*). *Indian Journal of Agronomy*. 51(3): 189-192.
- Ramana, V., Ramakrishna, M., Purushotham, K. and Reddy, K.B. (2010). Effect of biofertilizers on growth, yield attributes and yield of French bean (*Phaseolus vulgaris* L.). *Legume Research*. 33: 178-183.
- Saikia, J., Saikia, L., Phookan, D.B. and Nath, D.J. (2018). Effect of biofertilizer consortium on yield, quality and soil health of French bean (*Phaseolus vulgaris* L.). *Legume Research*. 41(5): 755-758.
- Sen, S., Mondal, C.K., Mandal, A.R. and Paria, N.C. (2006). Effect of Rhizobium culture and different levels of nitrogen on growth, yield and nodulation of French bean (*Phaseolus vulgaris* L.). *The Horticulture Journal*. 19: 268-272.
- Sharma, A., Sharma, R.P., Katoch, V. and Sharma, G.D. (2018). Influence of vermicompost and split applied nitrogen on growth, yield, nutrient uptake and soil fertility in pole type Frenchbean (*Phaseolus vulgaris* L.) in an acid alfisol. *Legume Research*. 41: 126-131.
- Sharma, S.K., Prajapati, S. and Raghuwanshi, O. (2015). Effect of organic manures and inorganic fertilizers on yield and economics of cowpea production (*Vigna unguiculata* L.). *Indian Research Journal of Genetics and Biotechnology*. 7(1): 152-155.
- Sheoran, O.P., Tonk, D.S., Kaushik, L.S., Hasija, R.C. and Pannu, R.S. (1998). Statistical Software Package for Agricultural Research Workers. Recent Advances in information theory, Statistics and Computer Applications by D.S. Hooda and R.C. Hasija Department of Mathematics Statistics, CCS HAU, Hisar. 139-143p.
- Shubhashree, K.S., Alagundagi, S.C., Hiremath, S.M., Chittapur, B.M., Hebsur, N.S. and Patil, B.C. (2011). Effect of nitrogen, phosphorus and potassium levels on growth, yield and economics of Rajmash (*Phaseolus vulgaris*). *Karnataka Journal of Agricultural Sciences*. 24: 283-285.
- Singh, V.K., Singh, G.R. and Dubey, S.K. (2014). Effect of agronomic practices on growth, dry matter and yield of Rajmash (*Phaseolus vulgaris* L.). *African Journal of Agricultural Research*. 9(51): 3711-3719.
- Subbiah, B.V. and Asija, G.L. (1956). A rapid procedure for the estimation of the available nitrogen in soils. *Current Science*. 25: 259-260.
- Verma, H.P., Chovatia, P.K., Sharma, S. and Shivan, A.C. (2015). Growth and yield of cowpea as influenced by nitrogen and phosphorus levels on medium black soils of Gujarat. *Annals of Agricultural Research New series*. 36: 400-404.
- Walkley, A. and Black, T.A. (1934). An estimation of soil organic matter and proposed modification of the chromic acid titration method. *Soil Science*. 37: 29-38.
- Yadav, N., Thakur, K.S. and Mehta, D.K. (2022). A practice to enhance soil physico-chemical properties and viable microbial count as effected by organic nutrients sources in garden pea under mid hill zone of Himachal Pradesh. *Indian Journal of Agricultural Research*. A-5979.