



# Enhancing Morpho-physiological Characters and Productivity of *Kharif* Groundnut Through Application of Plant Growth Retardants under East and South Eastern Coastal Zone of Odisha

S. Lenka<sup>1</sup>, G.S. Kumar<sup>1</sup>, S.K. Swain<sup>2</sup>, R.R. Dash<sup>3</sup>, K.C. Pradhan<sup>3</sup>, A.K. Sethy<sup>4</sup>

10.18805/LR-5362

## ABSTRACT

**Background:** Groundnut grown in *kharif* season incurs more vegetative growth with higher plant height which causes the upper aerial pegs not to reach the ground. So, less numbers of sound mature pods resulting in lower yield compared to other seasons. The aim of the following investigation was to examine the effect of different growth regulators on plant height including other morphological and physiological parameters of *kharif* groundnut.

**Methods:** The experiment was conducted during *kharif* seasons of 2021 and 2022, taking three growth regulators with different concentrations in main plot i.e. Cycocel (CCC) @ 500 and 1000 ppm, Brassinosteroides (BR) @ 10 and 15 ppm, Maleic hydrazide (MH) @ 100 and 200 ppm and scheduling of applications in sub plots i.e. 30 DAS, 50 DAS and 30 and 50 DAS with an additional control for observation with cv. Dharani (TCGS-1043) variety.

**Result:** The application of cycocel @ 1000 ppm recorded lowest plant height with maximum no. of primary branches and leaves / plant compared to other growth regulators and control. All physiological parameters studied in the experiment were maximum under foliar spray with CCC @ 1000 ppm. The scheduled application of growth regulators at 30 and 50 DAS decreased plant height, increased branches and leaves/plant significantly at different growth stages and also all physiological parameters compared to single 30 DAS and 50 DAS.

**Key words:** Cycocel, Groundnut, Plant growth regulators (PGRs).

## INTRODUCTION

Groundnut (*Arachis hypogaea* L. 2n=4x=40) referring to the formation of pods in the soil. It ranks as the 13<sup>th</sup> most significant food crop and 4<sup>th</sup> most important oilseed globally. It is also known as peanut, earthnut, monkey nut and energy capsule. The groundnut seed contains 47-53% oil and 26% protein and 11.5% starch. The haulm and oil cakes are used as valuable organic manures and feeding material for livestock containing 7.3% N; 1.5% P<sub>2</sub>O<sub>5</sub> and 1.3% K<sub>2</sub>O. In India, it holds the title of "King of Vegetable Oil Seeds" and plays a crucial role in the national edible oil economy covering an area of 49.1 lakh ha in 2022-23 (Reddy, 2022). In India, it is grown in all the three seasons such as *kharif*, *rabi* and *summer*, but major portion of groundnut i.e. around 85% of it is cultivated in *kharif* season covering an area of 45.53 lakh ha with production of 83.69 lakh tonnes (*Kharif* 2022-23, agricoop.nic). Odisha covers around 68000 ha under *kharif* groundnut having a production and productivity 101 th MT and 1465 kg/ha, respectively, 1.23 lakh ha with productivity of 1982 kg/ha in *rabi* season (Odisha Agriculture Statistics, 2018-19). The productivity of *kharif* groundnut is low in comparison to other two season due to more vegetative growth with higher plant height because of rainfall, which causes the upper aerial pegs not to reach the ground. This results in production of less numbers of sound mature pods and in turn less yields. Having indeterminate growth habit type,

<sup>1</sup>Department of Agronomy, College of Agriculture, Orissa University of Agriculture and Technology, Bhubaneswar-751 003, Orissa, India.

<sup>2</sup>Co-ordination Cell College, of Agriculture, Orissa University of Agriculture and Technology, Bhubaneswar-751 003, Orissa, India.

<sup>3</sup>OIC AICRP on Weed management, College of Agriculture, Orissa University of Agriculture and Technology, Bhubaneswar-751 003, Orissa, India.

<sup>4</sup>AICRP on Groundnut, College of Agriculture, Orissa University of Agriculture and Technology, Bhubaneswar-751 003, Orissa, India.

**Corresponding Author:** S. Lenka, Department of Agronomy, College of Agriculture, Orissa University of Agriculture and Technology, Bhubaneswar-751 003, Orissa, India.

Email: subhrasini@gmail.com

Orcids: 0000-0002-9207-4503, 0009-0003-6970-4532, 0000-0009-8458-0885, 0009-0004-5284-8043, 0009-0001-2449-8240.

**How to cite this article:** Lenka, S., Kumar, G.S., Swain, S.K., Dash, R.R., Pradhan, K.C. and Sethy, A.K. (2024). Enhancing Morpho-Physiological Characters and Productivity of *Kharif* Groundnut Through Application of Plant Growth Retardants under East and South Eastern Coastal Zone of Odisha. Legume Research. 1-7. doi: 10.18805/LR-5362.

**Submitted:** 04-06-2024 **Accepted:** 19-11-2024 **Online:** 30-01-2025

the vegetative and reproductive stage goes on simultaneously till harvest of the crop. This can be altered by the application of specific growth regulators to regulate the plant metabolism and to modify apical dominance. Plant

growth regulators are the organic compounds other than nutrients which in small amount promotes or inhibit or otherwise modify any physiological response in plants (Tukey *et al.* 1954). Cycocel (chlormequat), a plant growth regulator that inhibits plant growth, increases leaf density, retards stem elongation, extends shelf life and produces crops with shorter internodes, stronger stems, greener and thicker leaves. Brassinosteroids, steroid-based hormones in plants, regulate cell elongation, division, gravitropism, stress resistance and differentiation. Maleic hydrazide is a plant growth regulator used to reduce growth, increasing dry matter, chlorophyll content and promoting dormancy in seeds. With the above backgrounds in view, a field investigation was executed to evaluate the impact of several plant growth regulators at different concentration and scheduling on morphological and physiological parameters of *kharif* groundnut.

## MATERIALS AND METHODS

The present investigation was conducted at research field of AICRP on groundnut, Horticulture Research Station (HRS), Odisha University of Agriculture and Technology, Bhubaneswar in the *kharif* season, 2021 and 2022. The soil of the experimental plot was sandy loam in texture. The experimental site received 1135.2 and 1042.9 mm rainfall in 2021 and 2022, respectively during crop growth period. The average maximum and minimum temperature were (32.5 and 24.6°C) and (33.1 and 25.6°C) in 2021 and 2023, respectively. The nutrient status of soil was medium in nitrogen (301 N kg/ha), high in phosphorus (34.3 P<sub>2</sub>O<sub>5</sub> kg/ha) and medium in potassium (230 K<sub>2</sub>O kg/ha) and slightly acidic in nature having pH 5.6 and EC 0.42 dS/m. The experiment was laid out in split plot design with 18 treatment combinations. It consisted of three growth regulators at different concentrations in main plot viz. M<sub>1</sub>: Cycocel (CCC) @ 500 ppm, M<sub>2</sub>: Cycocel @ 1000 ppm, M<sub>3</sub>: Brassinosteroides (BR) @ 10 ppm, M<sub>4</sub>: Brassinosteroides @ 15 ppm, M<sub>5</sub>: Maleic hydrazide (MH) @ 100 ppm, M<sub>6</sub>: Maleic hydrazide @ 200 ppm and scheduling of applications in subplot i.e. S<sub>1</sub>: 30 DAS, S<sub>2</sub>: 50 DAS and S<sub>3</sub>: 30 and 50 DAS with an additional control observation. Solutions of PGRs were prepared and sprayed on the foliage of plants at 30 and 50 DAS with the help of knapsack sprayer as per treatment while in untreated control distilled water was sprayed at 30 and 50DAS. Groundnut cv. Dharani (TCGS-1043) variety was fertilized with uniform dose of (20-40-40 kg ha<sup>-1</sup>) of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O, respectively and was grown with standard agronomic practices. For morphological parameters, observations were taken from ten plants, which were randomly selected from each net plot of each treatment from all the replications, then uprooted carefully on 45, 65 and 85 DAS and at harvest. The plant height (cm) was measured from ground level to the growing tip of freshly opened leaf emerging from the main shoot. The number of primary branches and number of functional leaves were recorded from the same observational plants. Physiological

parameters were measured from 5 plants at several intervals starting from 45 DAS till harvest. The dry matter production and leaf area recorded at different stages of the crop were used to calculate the growth analysis indices as per the standard formulae (Radford, 1967). Data collected for various studies were subjected to the analysis of variance (ANOVA) appropriate to the design as given by Gomez and Gomez (1984). The Critical differences were worked out for the effects which are significant at 5% level of probability.

## RESULTS AND DISCUSSION

### Effect of plant growth regulators (PGRs) and time of application on morphological parameters

#### Plant height (cm)

The application of different plant growth regulators significantly reduced the plant height except BR compared to control. Application of CCC @ 1000 ppm significantly reduced plant height from 45 DAS till harvest compared to all other PGRs at different concentration and control. However, the spraying of BR @ 15 ppm increased the plant height, followed by BR @ 10 ppm. The time of administration of PGRs at 30 and 50 DAS significantly reduced the length of the plant from 45 DAS (36.8 cm) till harvest (66.8 cm) over other two time of application (Table 1). The application of CCC @ 1000 ppm at 30 and 50 DAS significantly reduced plant height from 45 DAS (30.1 cm) to the harvest (59.3 cm) in comparison to other growth regulators with time of applications. This might be due to the application of cycocel which decreased the apical dominance with less intermodal elongation by inhibition of gibberellin synthesis and promotes in case of brassinosteroides. These findings were similar to those of, Moulana *et al.* (2020); Sadeghi *et al.* (2014); Bhadane *et al.* (2020); Faldu *et al.* (2018); Sengupta *et al.* (2015); Lenka *et al.* (2023) and Karuppuswamy *et al.* (2021).

#### Number of main branches/plant

The numbers of branches/plant were significantly higher in the foliar application of the CCC @ 1000 ppm (5.8) which was at par with the BR @ 10 ppm (5.6). The lower numbers of branches/plant were recorded with the application of MH @ 200 ppm (4.3). The scheduled application of PGR at 30 and 50 DAS increased the number of branches/plant (5.4) significantly at different intervals of growth stages compared to 30 DAS and 50 DAS (Table 1). The heightened presence of branches, particularly in higher hormone concentrations during the peak vegetative stage, can be attributed to the influence of Cycocel on physiological processes like cell division, ion absorption, enzyme activities, photosynthetic efficiency and source-sink regulation. Similar increases in branch numbers were reported by Kaur *et al.* (2015); Faldu *et al.* (2018); Netwal *et al.* (2018); Mohanabharathi *et al.* (2019); Kurmi *et al.* (2020); Patil *et al.* (2008); Lakshmi *et al.* (2015) and Lenka *et al.* (2023).

**Number of leaves/plant**

Among different PGRs, application of the CCC @ 1000 ppm significantly increased the number of leaves /plant consecutively from 45 DAS (42.6), to the till the time of harvest (72.9), being maximum at 85 DAS (107.8). The number of leaves/plant increased significantly from 45DAS (45.0) till 85DAS (107.9), then decreased at harvest (74.1) as a result of spraying growth regulators at the time of 30 and 50 DAS (Table 1). The application of CCC @ 1000 ppm at 30 and 50 DAS significantly increased the number of leaves/plant at all the growth stages till 85 DAS (79.7) (Table 3). The leaves/plant is crucial as it serve as channel for supplying photosynthates to the pod development site, making a higher leaf count conducive to higher yield of groundnut. These results were supported with findings of, Prakash *et al.* (2006); Sengupta *et al.* (2015); Netwal *et al.* (2018); Sagar *et al.* (2020) and Siri *et al.* (2020).

**Physiological parameters****Leaf area index (LAI)**

The foliar application of the cycocel @ 1000 ppm significantly increased the LAI compared to other growth retardants from 45 DAS to 75 DAS. The LAI significantly increased with the application of PGRs at 30 and 50 DAS at all successive growth stages to 75 DAS (Fig 1) and decreased thereafter due to senescence and ageing of leaves. This may be due to good number of leaves with LAI, which was resulted higher production of photosynthates and then more partitioning of dry matter transported from leaf to pods and ultimately gave higher pod yield. Similar results were also cited by Upadhyay and Ranjan (2015); Prakash *et al.* (2006); Fatima *et al.* (2015); Sengupta *et al.* (2015) and Sagar *et al.* (2020).

**Crop growth rate ( $\text{g}^2\text{day}^{-1}$ )**

A perusal of data on crop growth rate showed that it increased up to 65 days and thereafter followed a decreasing trend. At 30-45DAS, CGR significantly increased in the BR @ 15 ppm ( $9.8 \text{ gm}^2 \text{ day}^{-1}$ ) but at par with the BR @ 10 ppm ( $9.1 \text{ gm}^2 \text{ day}^{-1}$ ). But from 45-65 DAS to 85 DAS-harvest, there was significant increase in crop growth rate with CCC @ 1000 ppm compared to other treatments. The spraying schedule of 30 and 50 DAS significantly increased CGR from 30-45 days ( $9.2 \text{ gm}^2 \text{ day}^{-1}$ ) till 85 DAS-harvest ( $5.2 \text{ gm}^2 \text{ day}^{-1}$ ) compared to 30 DAS and 50 DAS (Table 2). CGR is an indicator of productive efficiency of a crop and depends on genetic variables such as leaf area index, photosynthetic rate and leaf orientation and used as an indicative of light absorption. The crop growth increased during the peak period at 65 days and thereafter followed a decreasing trend due to transport of photosynthates towards pod during pod development stage. These results aligned with finding of Jadhav and Bhamburdekar (2014) and Meena *et al.* (2023).

**Table 1:** Effect of different levels of plant growth regulators and time of application on morphological parameters in *kharif* groundnut at successive stages of growth.

| Treatments                                 | Plant height (cm) |        |        | Main Branches/plant |        |        | Leaves/plant |         |         |
|--------------------------------------------|-------------------|--------|--------|---------------------|--------|--------|--------------|---------|---------|
|                                            | 45 DAS            | 65 DAS | 85 DAS | Harvest             | 45 DAS | 65 DAS | 85 DAS       | Harvest | Harvest |
| <b>Main plots -Plant growth regulators</b> |                   |        |        |                     |        |        |              |         |         |
| M <sub>1</sub> :CCC @500 ppm               | 33.7              | 43.7   | 56.1   | 65.7                | 4.4    | 4.4    | 4.5          | 4.7     | 69.6    |
| M <sub>2</sub> :CCC @1000 ppm              | 31.4              | 41.4   | 52.7   | 62.2                | 4.4    | 5.1    | 5.4          | 5.8     | 79.2    |
| M <sub>3</sub> :BR @ 10 ppm                | 40.9              | 54.5   | 65.2   | 72.8                | 4.0    | 4.9    | 5.3          | 5.6     | 75.6    |
| M <sub>4</sub> :BR@ 15 ppm                 | 43.8              | 57.6   | 68.2   | 75.6                | 4.2    | 4.6    | 4.8          | 5.1     | 70.7    |
| M <sub>5</sub> :MH@100 ppm                 | 36.3              | 48.2   | 59.7   | 68.4                | 4.0    | 4.3    | 4.4          | 4.7     | 65.1    |
| M <sub>6</sub> :MH @200 ppm                | 36.0              | 45.5   | 56.8   | 65.2                | 4.0    | 4.3    | 4.6          | 4.9     | 63.2    |
| SEM±                                       | 0.36              | 0.73   | 0.71   | 0.71                | 0.05   | 0.04   | 0.04         | 0.05    | 0.32    |
| CD (P=0.5)                                 | 1.12              | 2.30   | 2.25   | 2.23                | 0.14   | 0.13   | 0.12         | 0.15    | 0.99    |
| <b>Sub plots- Time of application</b>      |                   |        |        |                     |        |        |              |         |         |
| S <sub>1</sub> :30 DAS                     | 36.9              | 49.2   | 60.7   | 69.5                | 4.3    | 4.5    | 4.8          | 5.1     | 68.84   |
| S <sub>2</sub> : 50 DAS                    | 37.4              | 48.9   | 60.3   | 68.7                | 3.8    | 4.5    | 4.8          | 5.1     | 71.44   |
| S <sub>3</sub> : 30 and 50 DAS             | 36.8              | 47.4   | 58.3   | 66.8                | 4.5    | 4.9    | 5.2          | 5.5     | 74.14   |
| SEM±                                       | 0.23              | 0.49   | 0.48   | 0.49                | 0.04   | 0.03   | 0.05         | 0.05    | 0.34    |
| CD (P=0.05)                                | 0.62              | 1.31   | 1.29   | 1.34                | 0.10   | 0.09   | 0.12         | 0.14    | 1.00    |
| Control                                    | 40.0              | 51.5   | 62.8   | 71.3                | 3.8    | 4.1    | 4.4          | 4.6     | 63.7    |


**Relative growth rate ( $\text{g g}^{-1} \text{day}^{-1}$ ) (RGR)**

The application of BR @ 15 ppm significantly noted higher RGR at 30-45 DAS ( $0.051 \text{ g g}^{-1} \text{ day}^{-1}$ ), being at par with the CCC @ 1000 ppm. From 45-65 DAS till 85 DAS-harvest, CCC @ 1000ppm registered significantly higher RGR over all other treatments along with control. The foliar spraying of PGRs at 30 and 50 DAS significantly increased RGR at all the successive growth stages over other scheduling of applications. However, there was no significant difference between scheduling of growth regulators application at time of harvest (Fig 2). The increase in RGR (increase in dry matter of crop per unit of existing dry matter over time) through foliar application of growth regulator over control could be attributed to enhanced photosynthetic efficiency achieved through increased leaf thickness, chlorophyll content and improved translocation of carbon.

Foliar application of growth retardant in groundnut resulted peak relative growth rate prior to pod initiation, followed by decrease during pod filling period over control. Similar results were also reported by Bharud and Pawar (2005) and Meena *et al.* (2023).

**Net Assimilation Rate ( $\text{mg cm}^{-2} \text{day}^{-1}$ ) (NAR) and Absolute Growth Rate ( $\text{g day}^{-1}$ ) (AGR)**

Foliar application of CCC @ 1000 ppm significantly enhanced the NAR and AGR from 30-45 DAS till 45-65 DAS over all other PGRs and then followed a declining trend till harvest. Similar trend was also observed with spraying schedule of 30 and 50 DAS (Fig 3). Net assimilation rate is a measure of photosynthesis of leaves in crop community due to optimum LAI. Increase in NAR and AGR at 65 DAS *i.e.* at pod development stage might be due to increased efficiency of leaves for photosynthesis



**Fig 1:** Effect of different levels of plant growth regulators and time of application on leaf area index in *kharif* groundnut at successive stages of the growth.

**Table 2:** Effect of different levels of plant growth regulators and time of application on pod yield (kg/ha), haulm yield (kg/ha) and shelling percentage (%) in *kharif* groundnut.

| Treatments                        | Pod yield(kg/ha) | Haulm yield(kg/ha) | Shelling% |
|-----------------------------------|------------------|--------------------|-----------|
| <b>Plant growth regulator (M)</b> |                  |                    |           |
| M <sub>1</sub> : CCC@ 500 ppm     | 1551.9           | 3544.5             | 71.8      |
| M <sub>2</sub> : CCC@ 1000 ppm    | 1847.1           | 3677.8             | 72.9      |
| M <sub>3</sub> : BR @ 10 ppm      | 1771.5           | 4348.7             | 72.4      |
| M <sub>4</sub> : BR @15 ppm       | 1681.4           | 4652.3             | 71.8      |
| M <sub>5</sub> : MH @100 ppm      | 1552.4           | 3516.9             | 71.6      |
| M <sub>6</sub> : MH @ 200 ppm     | 1747.9           | 3550.0             | 72.1      |
| SEm±                              | 32.4             | 56.8               | 0.22      |
| CD(P=0.05)                        | 102.1            | 179.0              | 0.68      |
| <b>Time of application(S)</b>     |                  |                    |           |
| S <sub>1</sub> : 30 DAS           | 1560.5           | 3947.7             | 71.3      |
| S <sub>2</sub> : 50 DAS           | 1718.7           | 4053.6             | 72.5      |
| S <sub>3</sub> : 30 and 50 DAS    | 1796.8           | 4193.8             | 72.6      |
| SEm±                              | 21.9             | 58.0               | 0.12      |
| CD(P=0.05)                        | 59.3             | 156.8              | 0.32      |
| Control                           | 1432             | 4757.2             | 70.8      |


resulting in higher production of photosynthates as a response of photosynthetic apparatus to increase the demand for assimilates by growing pod. Similar findings on increase in NAR by PGRs were also reported by Meshram *et al.* (2018) and Saheen *et al.* (2019).

#### Leaf area ratio ( $\text{cm}^2 \text{g}^{-1}$ )

The treatment imposed with foliar application CCC @ 1000 ppm incurred significantly maximum LAR at 30-45 DAS and decreased in subsequent intervals till harvest under all growth regulators. The spraying of PGRs at 30 and 50 DAS significantly increased LAR at 30-45 DAS, followed by decreasing trend till harvest (Table 2). This trend of result implies that cellular level of developmental process were mostly completed before flowering. Management of nutrition and hormone during this growth phase can add to growth and productivity of *kharif* groundnut by effectively regulating metabolic processes. The present findings were in close conformity of Sagar *et al.* (2020).

#### Yield and yield component


Compared to the control, the application growth regulators significantly enhanced the pod yield and yield attributes in *kharif* groundnut. The application of the CCC @ 1000 ppm significantly increased the pod yield (1847.1 kg/ha) and shelling% (72.9%) and haulm yield was higher under control (4757.2 kg/ha). The foliar spray growth retardants at the time of 30 and 50 DAS significantly increased the pod yield and shelling% to 1796.8 kg/ha and 2.6%, respectively. Groundnut pod yield is a function of dry matter accumulation and yield attributes. In groundnut, the application of the plant growth retardants significantly increased the yield and yield components compared to the control. The application of plant growth regulators increased the total biomass and then might have resulted in increased transport of assimilates from source to sink and their ultimate conversion into final reserved food (Vardhini and Rao, 1999). These results were supported with the findings




**Fig 2:** Effect of different levels of plant growth regulators and time of application on relative growth rate ( $\text{g g}^{-1} \text{day}^{-1}$ ) in *kharif* groundnut at successive stages of the growth.

**Table 3:** Effect of different levels of plant growth regulators and time of application on crop growth rate and leaf area ratio in *kharif* groundnut at successive stages of growth.

| Treatments                               | Crop growth rate ( $\text{g}^2 \text{day}^{-1}$ ) |        |        |         | Leaf area ratio ( $\text{cm}^2 \text{g}^{-1}$ ) |        |        |         |
|------------------------------------------|---------------------------------------------------|--------|--------|---------|-------------------------------------------------|--------|--------|---------|
|                                          | 45 DAS                                            | 65 DAS | 85 DAS | Harvest | 45 DAS                                          | 65 DAS | 85 DAS | Harvest |
| <b>Mainplots-Plant growth regulators</b> |                                                   |        |        |         |                                                 |        |        |         |
| M <sub>1</sub> : CCC @500 ppm            | 6.7                                               | 16.3   | 12.6   | 4.0     | 39.4                                            | 29.5   | 26.4   | 24.5    |
| M <sub>2</sub> : CCC @1000 ppm           | 7.6                                               | 26.4   | 15.2   | 5.6     | 47.4                                            | 28.3   | 25.0   | 24.3    |
| M <sub>3</sub> : BR @ 10 ppm             | 9.1                                               | 17.8   | 14.3   | 4.6     | 41.6                                            | 28.1   | 26.0   | 24.5    |
| M <sub>4</sub> : BR@ 15 ppm              | 9.8                                               | 22.6   | 12.9   | 4.5     | 39.7                                            | 29.8   | 25.7   | 24.2    |
| M <sub>5</sub> : MH@100 ppm              | 5.7                                               | 16.5   | 12.6   | 3.8     | 41.2                                            | 31.8   | 26.9   | 25.7    |
| M <sub>6</sub> : MH @200 ppm             | 6.9                                               | 17.0   | 14.2   | 4.4     | 40.6                                            | 28.8   | 25.8   | 23.8    |
| <b>SEm(±)</b>                            | 0.23                                              | 0.24   | 0.32   | 0.18    | 0.147                                           | 0.113  | 0.10   | 0.097   |
| <b>CD (P=0.5)</b>                        | 0.69                                              | 0.73   | 0.96   | 0.56    | 0.431                                           | 0.337  | 0.312  | 0.285   |
| <b>Sub plots- Time of application</b>    |                                                   |        |        |         |                                                 |        |        |         |
| S <sub>1</sub> : 30 DAS                  | 8.5                                               | 16.3   | 12.5   | 4.2     | 43.0                                            | 28.1   | 25.5   | 23.6    |
| S <sub>2</sub> : 50 DAS                  | 4.8                                               | 20.5   | 13.6   | 4.7     | 38.1                                            | 29.3   | 26.1   | 24.5    |
| S <sub>3</sub> : 30 and 50 DAS           | 9.2                                               | 21.3   | 14.8   | 5.2     | 44.5                                            | 31.4   | 27.5   | 26.3    |
| <b>SEm±</b>                              | 0.09                                              | 0.08   | 0.13   | 0.07    | 0.690                                           | 0.425  | 0.405  | 0.415   |
| <b>CD (P=0.05)</b>                       | 0.26                                              | 0.23   | 0.38   | 0.20    | 1.987                                           | 1.215  | 2.502  | 1.155   |
| <b>Control</b>                           | 9.7                                               | 18.0   | 13.9   | 4.8     | 39.4                                            | 29.5   | 26.4   | 24.5    |



**Fig 3:** Effect of different levels of plant growth regulators and time of application on net assimilation rate ( $\text{cm}^2 \text{day}^{-1}$ ) in *kharif* groundnut at successive stages of the growth.



**Fig 4:** Effect of different levels of plant growth regulators and time of application on absolute growth rate ( $\text{g day}^{-1}$ ) in *kharif* groundnut at successive stages of the growth.

of Janarjuna *et al.* (2001); Sadeghi *et al.* (2014); Khalilzadeh *et al.* (2016); Patel *et al.* (2019) and Kurmi *et al.* (2020).

## CONCLUSION

The application of plant growth regulators has a significant effect on the morphological and physiological parameters of *kharif* groundnut. Foliar application of CCC @ 1000 ppm at 30 and 50 DAS reduced the plant height and enhanced primary branches, leaves per plant and several physiological parameters such as LAI, CGR and NAR over other growth regulators, adding to groundnut growth and productivity in *kharif* season.

## ACKNOWLEDGEMENT

The present study was supported by ICAR-Directorate of Groundnut Research, Junagadh, Gujarat and Directorate of Research, Odisha University of Agriculture and Technology, Bhubaneswar.

## Disclaimers

The views and conclusions expressed in this article are solely those of the authors and do not necessarily represent the views of their affiliated institutions. The authors are

responsible for the accuracy and completeness of the information provided, but do not accept any liability for any direct or indirect losses resulting from the use of this content.

## Conflict of interest

The authors declare that there is no conflict of interest regarding the publication of this article. No funding or sponsorship influenced the design of the study, data collection, analysis, decision to publish, or preparation of the manuscript.

## REFERENCES

- Bhadane, R.S., Prajapati, K.R., Ombase, K.C. and Patel, D.B. (2020). Effect of seed priming and foliar spraying of pgrs on morpho-physiology, growth and yield in greengram (*Vigna radiata* L.). *Legume Research*. 45: 435-444. doi: 10.18805/LR-4434.
- Bharud, R.W. And Pawar, M.R. (2005). physiological basis of yield variation in groundnut varieties under summer conditions. *Journal of Maharashtra Agricultural University*. 30(1): 100-102.
- Faldu, T.A., Kataria, G.K., Singh, C.K. And Paghadal, C.B. (2018). Influence of plant growth regulators on morphological and physiological parameters of groundnut (*Arachis hypogaea* L.) cv.gjg-9. *Journal of Pharmacognosy and Phytochemistry*. 7: 2341-2344.

- Fatima, K.G., Nuaimi, A.L., Laith, M.J. And Shamma, A.L. (2015). Effect of chemical mutagens on some morphological traits of *Vicia faba* L. cv. aqadulce. Iraqi Journal of Science. 56: 2506-2512.
- Gomez, K.A. and Gomez, A.A. (1984). Statistical Procedures for Agriculture Research.
- Jadhav, S.H. and Bhamburdekar, S.B. (2014). Effect of salicylic acid on germination performance in groundnut. Inter-National Journal of Applied Biology and Pharmaceutical Technology. 2: 224-227.
- Janarjuna, S., Shivraj, B., Nanjudappa, G. and Sridhara, S. (2001). Uptake and recovery of n,p,k in groundnut as influenced by application of Ca, Mo and plant growth regulators alone and in combination. Indian Agriculturist. 45(1 and 2): 61-64.
- Karuppusamy, G., Jeyakumar, P., Chandrasekhar, C.N., Jayamani, P. and Gopal, N.O.(2021). Effect of plant growth regulating compounds and nutrients on morphophysiological characters in pigeonpea (*Cajanus cajan*). Journal of Pharmacognosy Phytochemistry. 9: 1794-1797.
- Kaur, J., Ram, H. and Gill, B.S. (2015). Agronomic performance and economic analysis of soybean (*Glycine max* L.) in relation to growth regulating substances in punjab, India. Legume Research. 38: 603-608. doi: 10.18805/lr.v38i5.5936.
- Khalilzadeh, R., Sharifi, S.R. and Jalilian, J. (2016). Antioxidant status and physiological responses of wheat (*Triticum aestivum* L.) to cycocel application and biofertilizers under water limitation condition. Journal of Plant Interactions. 11(1): 130-137.
- Kurmi, A.K., Naruka, I.S., Kushwah, S.S. and Shaktawa, R.P.S. (2020). Effect of plant growth regulators on growth, yield and quality of coriander (*Coriandrum sativum* L.). Environment and Ecology. 38: 710-714.
- Lakshmi, J., Rame, G., Parashivamurthy, Narayanaswamy, S. and Shivanandam, V.N. (2015). Influence of pre-flowering pinching and maleic hydrazide spray on plant growth, seed yield and quality attributes in fenugreek. Legume Research. 38(3): 353-357. doi: 10.5958/0976-0571.2015.00097.1.
- Lenka, S., Swain, S.K., Pradhan, K.C. and Dhal, A. (2023). Effect of different levels and time of application of paclobutrazol on morphology, yield and yield attributing characters and economics of groundnut (*Arachis hypogaea* L.). Legume Research. 46(4): 428-431. doi: 10.18805/LR-4391.
- Meena, B.P., Kataria, G.K., Singh, C.K., Godha, U.N. And Pandya, Y. (2023). Impact of plant growth modulators on morpho-physiological characteristics of *kharif* groundnut variety GJG-31. The Pharma Innovation Journal. 12: 1419-1422.
- Meshram, S.D.R.D., Deotale, K.H., Chute, G.N., Jadhav and G.A. Padghan. (2018). Morpho-physiological and yield responses of soybean foliar sprays of chitosan and IBA. Journal of Soils and Crops. 28: 121-127.
- Mohanabharathi, M., Sritharan, N., Senthil, A. and Ravikesavan, R. (2019). Physiological studies for yield enhancement in finger millet under drought condition. Journal of Pharmacognosy and Phytochemistry. 8: 3308-3312.
- Moulana, S., Prasad, V.M. and Bahadur, V. (2020). Effect of different levels of cycocel (ccc) on two different cultivars of okra under prayagraj agroclimatic conditions (*Abelmoschus esculantus* L.). International Journal of Chemical Studies. 8: 133-136.
- Netwal, M., Choudhary, M.R., Rajkumar, J., Devi, S. and Suman, C. (2018). Exogenous application of brassinoid and salicylic acid enhances on growth, yield and nutritional quality of indian bean (*Lablab purpureus* L. var. typicus). Journal of Pharmacognosy and Phytochemistry. 7: 2093-2096.
- Odisha Agriculture Statistics. (2018-19). Directorate of Agriculture and Food Production, Government of Odisha.
- Patel, R., Thakral, S.K. and Dudi, S. (2019). Wheat yield maximization through nutrient management and use of growth retardants. Journal of Pharmacognosy and Phytochemistry. 8(6): 1046-1047.
- Patil, C.N., Mohorkar, V.K., Dod, V.N., Peshattiwar, D.D., Kayande, N.V. and Gomase, D.G. (2008). Effect of seed treatment with gibberellic acid and maleic hydrazide on growth, seed yield and quality of okra. Asian Journal of Horticulture. 3: 74-78.
- Prakash, M., Saravanan, K., Sunil Kumar, B., Sabesan, T., Gokulakrishnan, J., Suganthi, S. and Ganesan, J. (2006). Response of brassinosteroids on growth and pod yield of groundnut. Crop Research. s1: 419-423.
- Radford, P.J. (1967). Growth analysis formulae- Their use and abuse. Crop Science. 7: 171-175.
- Reddy, G.R. (2022). Crop outlook report of andhra pradesh, january to december. centre for agriculture and policy research (carp), ANGRAU Andhra Pradesh. <https://angrau.ac.in/downloads/amic/outlookreports/2022>.
- Sadeghi, S.M., Vakili, D. and Hashemi, S.A. (2014). Investigation of cycocel and mycorrhiza effecting yield and yield components of peanut planting (*Arachis hypogaea* L.) in different planting distances. Advanced Biomedical Research. 5: 136-139.
- Sagar, M.M., Dhanoji, M.K., Meena, Suma, T.C. and Yadahalli, G.S. (2020). Physiological approaches to improve productivity of safflower (*Carthamus tinctorius* L.) through foliar nutrition. journal of pharmacognosy and phytochemistry, 9: 1257-1259.
- Sengupta, K. and Tamang, D. (2015). Response of green gram to foliar application of nutrients and brassinolide. Journal Crop and Weed. 11: 43-45.
- Shaheen, A.M., Ragab, M.E., Rizk, F.A., Mahmoud, H., Soliman, M.M. and Omar, N.M. (2019). Effect of some active stimulants on plant growth, tubers yield and nutritional values of potato plants grown in newly reclaimed soil. Journal of Animal and Plant Science. 29: 215-275.
- Siri, C., Patil, R.P., Amaregouda, A., Suma, T.C. and Kenganal, M. (2020). Influence of different agrochemicals on yield attributes and yield of chilli (*Capsicum annum* L.) grown under hk region of Karnataka. International Journal of Chemical Studies. 8: 2629-2631.
- Tukey, H.B., Went, F.W., Muir, R.M. and Overbreek, J.V. (1954). Nomenclature of chemical plant regulators. American Society of Plant Physiology. 29: 307-308.
- Upadhyay, R.G., Ranjan, R. (2015). Effect of growth hormones on morphological parameters, yield and quality of soybean (*Glycine max* L.). Academia Journal of Agricultural Research. 33: 1899-1904.
- Vardhini, B.V. And Rao, S.S .R. (1999). Effect of brassinosteroids on nodulation and nitrogenase activity in groundnut (*Arachis hypogaea* L.). Plant Growth Regulation. 28: 165-167.