



# Adaptive Response of Chickpea Varieties with Diverse Breeding Pathways Across Six Contrasting Semi-arid Environments using AMMI and GGE Biplot Analysis

Fatiha Zine-Zikara<sup>1,2</sup>, Fatma Zohra Bouras<sup>2</sup>, Wahiba Tiliouine<sup>1</sup>,  
Samia Hamdi-Daoud<sup>1</sup>, Amar Haireche<sup>3</sup>, Oussama Baghdadli<sup>3</sup>,  
Rabah Rahim<sup>3</sup>, Mohamed Réda Bououlcis<sup>3</sup>, Meriem Laouar<sup>2</sup>

10.18805/LRF-901

## ABSTRACT

**Background:** This study aimed to evaluate the variety  $\times$  environment interaction for high grain yields and stability of chickpea varieties with different breeding pathways, across six contrasting semi-arid environments.

**Methods:** During one cropping season, trials were conducted at the farmers' fields. Ten genotypes were evaluated in a completely randomized block design, with three replicates in six semi-arid contrasting environments. Analysis of variance, AMMI and GGE biplot models were used.

**Result:** Genetic variability of grain yield was strongly influenced by the environment that explained almost 26% of the total variance, whereas the  $G \times E$  interaction explained 16%, highlighting the importance of genotype-specific adaptation. Two discriminating environments, E5 and E4 were selected. The introduced variety G8-IV showed excellent stability and high yield ( $1.32 \text{ t-ha}^{-1}$ ), demonstrating its plasticity. Similarly, participatory variety G4-PS showed high stability despite its selection in a specific terroir, demonstrating a wider-than-expected capacity for adaptation. Conversely, some local varieties, such as G1-LV, showed narrower adaptation limited to environments such as E3. These elements allow breeding to be oriented towards the greater resilience of farming systems in the face of climatic variability.

**Key words:** Additive main effects and multiplicative interaction (AMMI), Chickpea, Genotype plus genotype-by-environment interaction biplot (GGE), Semi-arid, Varieties pathway, Yield stability.

## INTRODUCTION

Chickpea (*Cicer arietinum* L.) is a globally important food legume and a major source of protein in human diets (Baite *et al.*, 2017). It contributes substantially to soil fertility through biological nitrogen fixation, making it an ideal rotational crop in sustainable cereal-based systems (Saraf *et al.*, 1998).

Chickpea is cultivated under rainfed systems in semi-arid Mediterranean environments (Bouri *et al.*, 2019). It requires relatively low water (300-400 mm), but its production is increasingly constrained by irregular rainfall, terminal drought during flowering and pod filling (Street *et al.*, 2016) and high temperatures. In fact, terminal drought can potentially reduce yields by 58-95% compared to irrigated conditions (Farooq *et al.*, 2017).

The semi-arid Algerian environments are broadly representative of Mediterranean rainfed systems (Djellouli, 2025), where water scarcity is increasingly critical from eastern to western regions (Alrteimei *et al.*, 2022).

Developing varieties with drought tolerance and adaptation to variable rainfall is crucial to sustain productivity and food security under climate variability (Arriagada *et al.*, 2022). Chickpea breeding pathways include: (i) pure (bred) varieties, genetically homogeneous; (ii) participatory selection varieties, co-developed with farmers to enhance local adaptation and adoption; and (iii) heterogeneous local varieties, maintained by farmers and characterized by high

<sup>1</sup>INRAA, National Institute of Agronomic Research of Algeria, Biotechnology and Plant Breeding Division, Algiers, Algeria.

<sup>2</sup>ENSA, National Higher School of Agronomy-El Harrach, Laboratory of Integrated Improvement of Plant Productions (C2711100), Algiers, Algeria.

<sup>3</sup>OAIC, Algerian Inter professional Cereals Office, Algeria.

**Corresponding Author:** Fatiha Zine-Zikara, Meriem Laouar; INRAA, National Institute of Agronomic Research of Algeria, Biotechnology and Plant Breeding Division, Algiers, Algeria; ENSA, National Higher School of Agronomy-El Harrach, Laboratory of Integrated Improvement of Plant Productions (C2711100), Algiers, Algeria.

Email: fatiha.zine.zikara@gmail.com; meriem.laouar@edu.ensa.dz

**How to cite this article:** Zine-Zikara, F., Bouras, F.Z., Tiliouine, W., Hamdi-Daoud, S., Haireche, A., Baghdadli, O., Rahim, R., Bououlcis, M.R. and Laouar, M. (2025). Adaptive Response of Chickpea Varieties with Diverse Breeding Pathways Across Six Contrasting Semi-arid Environments using AMMI and GGE Biplot Analysis. *Legume Research*. 1-11. doi: 10.18805/LRF-901.

**Submitted:** 08-09-2025 **Accepted:** 19-11-2025 **Online:** 05-12-2025

intra-population diversity and resilience under marginal rainfed conditions (Raggi *et al.*, 2019).

Expanding the genetic base enhances chickpea resilience to drought and heat stress while improving productivity (Kharrrat *et al.*, 2023). Nevertheless, some self-

pollinating homogeneous varieties retain notable tolerance to abiotic stresses (Panigrahi *et al.*, 2024). In the context of increasing climatic variability and a limited genetic pool (Varshney *et al.*, 2024), it is therefore important to determine not only the best-adapted genotypes but also the most appropriate type of classical selection to adopt to maximize productivity for long-term resilience.

Understanding genotype  $\times$  environment is critical for guiding varietal selection toward yield stability and resilience under variable semi-arid conditions. While multi-year trials are ideal, evaluating varieties across contrasting environments within a single year can provide robust insights into stability and adaptation, as demonstrated in recent studies (Khan *et al.*, 2024).

To capture these interactions, this study employs Additive Main Effects and Multiplicative Interaction (AMMI) and Genotype plus Genotype  $\times$  Environment (GGE) biplot analyses, integrating robust statistical modeling with visual interpretation (Laxuman *et al.*, 2025; Shekhawat *et al.*, 2024; Annicchiarico, 2020; Zobel *et al.*, 1988).

The specific objectives of this study are to: (i) evaluate the performance of chickpea varieties differing by breeding pathway across six contrasting Algerian environments; (ii) quantify G $\times$ E interactions for grain yield and stability; and (iii) identify varieties most stable and adapted for rainfed semi-arid conditions, provide guidance for sustainable varietal deployment.

## MATERIALS AND METHODS

### Genetic material

Ten varieties of chickpea were obtained through different selection methods; their pedigree, origins and supplying institutes are listed in Table 1. They correspond to local

varieties (LV), participatory selected varieties (PS) and introduced varieties (IV) selected by ICARDA.

### Experimental sites and environmental variability

Six sites, favorable for chickpea cultivation, with contrasting conditions (climatic and edaphic) were selected for the study and are illustrated in Fig 1 (S1 to S6).

Multiple Factor Analysis was carried out using climatic data (Algerian National Office of Meteorology) for twenty-three years (2000 to 2022) and soil analysis (2021-2022). This enabled us to confirm the environmental variability of the sites along the axes 1-2-3.

For each site, we assigned a code to the corresponding environment by identifying its most distinctive characteristics (Table 2).

### Trial setup

The experiments were conducted during the 2021-2022 season on a farmer's field. The climatic and edaphic characteristics of each experiment by site are presented in Table 2.

Climatic data for 2021-2022 (Fig 2) indicate higher or stable rainfall from March to May (early spring) compared to the 23-year average, but reduced precipitation from December to February (winter) and after May. This atypical winter deficit contrasts with the usual spring water stress in these semi-arid regions and aligns with recent reports of declining winter rainfall, shorter wet seasons and greater variability in northern Algeria (Bessaoud and Montaigne, 2009; Meddi *et al.*, 2010; Djellouli and Laborde, 2022). Mean temperatures remained close to the historical average.

### Experimental design and recorded traits

The experiment was laid out as a completely randomized block design with three replicates. Elementary plots




Fig 1: Algerian production chickpea areas and location of study sites.

**Table 1:** Breeding Pathway, codes, names, pedigrees, origins, source and target environments of ten chickpea varieties.

| Breeding pathway                    | Code   | Variety             | Pedigree                                                   | Origin        | Source | Selection purpose                                                   |
|-------------------------------------|--------|---------------------|------------------------------------------------------------|---------------|--------|---------------------------------------------------------------------|
| Local variety (LV)                  | G1-LV  | Tadjena             | Not available                                              | Farmer/Chlef  | OAIC   | Large grain                                                         |
|                                     | G2-LV  | Rabat               |                                                            | Farmer/Skikda |        |                                                                     |
|                                     | G3-LV  | Bouazza             |                                                            | Farmer/SBA    |        |                                                                     |
|                                     | G4-PS  | F10-5*              |                                                            | Not available |        |                                                                     |
| Participatory selected variety (PS) | G5-PS  | F10-49*             |                                                            |               |        | Height plant, medium grain size, anthracnose tolerance              |
|                                     | G6-PS  | F4-82*              |                                                            |               |        | Large grain size, drought resistance                                |
| Introduced variety (IV)             | G7-IV  | ILC 32-79           | Line                                                       | ICARDA        |        | Large grain, tolerant to anthracnose                                |
|                                     |        |                     |                                                            |               |        | Height plant, upright habit, productive, very resistant to disease. |
|                                     |        |                     |                                                            |               |        | Height plant, disease tolerance, stable yield.                      |
|                                     | G8-IV  | Ghab4 (FLIP 93-93c) |                                                            |               |        |                                                                     |
|                                     |        |                     | X89TH258/<br>(flip85-122CX<br>flip82- 150C)<br>/flip86-77C |               |        |                                                                     |
|                                     | G9-IV  | Flip 90-13C         | X87TH317/<br>(Pl.Se.Be.81-40X<br>ILC195)Xflip84-46C        |               |        | Anthracnose tolerance                                               |
|                                     |        |                     | X2001TH16/<br>(Flip97- 25CXs98588)<br>XS99481              |               |        |                                                                     |
|                                     | G10-IV | Flip 05-153C        |                                                            |               |        | Large grain                                                         |

\*: Selected by Dr. Labdi M. and Mrs. Hamdi S., at National Institute of Agronomic Research of Algeria, INRAA, OAIC: Algerian Inter professional Cereals Office, ICARDA: International center for agricultural research in the dry areas.

measured 6 m × 1.8 m and consisted of four rows sown at 20 seeds m<sup>-1</sup> (150 kg ha<sup>-1</sup>) with a thousand-seed weight of 340 g. The crop cycle was between December and June for all sites except site 1 (sowing in February).

Six agronomic traits were selected based on their relevance to the yield, plant architecture and adaptation to environmental constraints (Singh *et al.*, 2020): grain yield, plant height, number of primary branches, stem length to the first pod, 100-seed weight and the harvest index.

### Statistical analysis

Analysis of variance (ANOVA) was conducted to assess yield differences among varieties within each environment, aiming to test the variety effect and to classify the ten cultivars individually or according to their breeding pathways. In addition, a factorial ANOVA was applied to several agronomic traits measured in the six environments, considering the main effects of the selection pathway, environment and their interaction. Analyses were carried out using RStudio software (version 2023.12.0+369) Multiple comparisons of the means were performed using Tukey's test.

Statistical analyses of the Additive Main Effects and Multiplicative Interaction (AMMI) Zobel *et al.* (1988) and genotypic main effect + genotype-by-environment interaction (GGE) biplot models (Ponsiva *et al.*, 2024) were performed using the "metan" package in RStudio software version 12.0. AMMI models were applied to study the G×E interaction, adaptation and stability of varieties across the six environments.

According to Gauch *et al.* (2008), the display of nominal variety yields (AMMI-1) clearly describes the winning varieties and adaptive responses.

## RESULTS AND DISCUSSION

### Grain yield performance across environments

ANOVA for grain yield revealed significant differences (1.01 t·ha<sup>-1</sup> to 2.18 t·ha<sup>-1</sup>) among varieties across most environments, except for E1 (Table 3).

Participatory selection varieties showed the most consistent yields across environments, performing best under moderately productive conditions. Local varieties were well adapted to favorable environments, particularly under adequate spring rainfall and exhibited moderate stability. In contrast, introduced varieties displayed the highest variability, occasionally achieving high yields but often underperforming, indicating greater sensitivity to environmental conditions.

### Effects of breeding pathways, environment and their interaction on morpho-agronomic traits

The environment had a highly significant effect on all traits, except Number of primary branches (Table 4), highlighting the major role of local conditions in shaping phenotypic expression. The pathway × environment interaction was especially pronounced for 100-grain weight.

**Table 2:** Climatic and soil variations of the six study sites.

| Site/Code                                | S1/E1                                       | S2/E2        | S3/E3        | S4/E4                                  | S5/E5            | S6/E6                                  |
|------------------------------------------|---------------------------------------------|--------------|--------------|----------------------------------------|------------------|----------------------------------------|
| Average rainfall (mm)_Year               | 495.79                                      | 499.08       | 464.76       | 426.6                                  | 508.8            | 574.44                                 |
| Average rainfall (mm)_cycle culture      | 299.10                                      | 380.26       | 225.07       | 294.13                                 | 292.21           | 259.37                                 |
| MinT (°C)/month                          | 5.41/January                                | 7.66/January | 6.54/January | 8.68/January                           | 6.96/January     | 0.51/January                           |
| MaxT (°C)/month                          | 34.39/July                                  | 33.76/August | 35.52/August | 31.43/August                           | 37.10/July       | 33.90/July                             |
| Soil texture                             | Silty                                       | Silty clay   | Silty clay   | Silty                                  | Clayey silt      | Silty clay                             |
| Soil-EC (ds/m)                           | 0.14                                        | 0.13         | 0.16         | 0.38                                   | 0.11             | 0.24                                   |
| Soil-TAL (%)                             | 10.07                                       | 2.65         | 4.5          | 6.64                                   | 5.36             | 3.15                                   |
| Soil-OM (%)                              | 2.79                                        | 1.29         | 1.32         | 0.44                                   | 1.84             | 3.27                                   |
| Soil-P <sub>2</sub> O <sub>5</sub> (ppm) | 4.87                                        | 4.26         | 4.56         | 6.39                                   | 3.64             | 11.25                                  |
| Soil-K <sub>2</sub> O (mg/kg)            | 0.27                                        | 0.24         | 0.25         | 0.24                                   | 0.63             | 3.09                                   |
| Characteristics                          | Dry, with calcareous and nutrient-rich soil | Intermediate | Intermediate | Hot, with low rainfall and saline soil | Moderately rainy | Cold and rainy, with fertile/rich soil |

EC: Electrical conductivity, TAL: Total active limestone, OM: Organic matter.




Fig 2: Monthly temperatures and precipitation over the 2021-2022 growing season and inter annual monthly mean values recorded over the period 2000-2022 at the six environments (E1 to E6).

The results indicate that the selection method influences the genetic response of the lines depending on the environment, emphasizing the importance of evaluating varieties under diverse conditions to fully capture their potential.

### AMMI and GGE analyses and multivariate exploration of yield stability and genotype × environment interaction

Analysis of variance (Table 5) revealed significant effects of genotype and environment, as well as a highly significant

**Table 3:** Analysis of variance (ANOVA) for grain yield (t·ha<sup>-1</sup>) of ten varieties corresponding to three selection types across six environments, with hierarchical mean comparison using the LSD test.

| Nature of breeding pathway             | Varieties | Environment          |                     |                     |                      |                      |
|----------------------------------------|-----------|----------------------|---------------------|---------------------|----------------------|----------------------|
|                                        |           | E2                   | E3                  | E4                  | E5                   | E6                   |
| Local varieties (LV)                   | G1        | 0.92 <sup>de</sup>   | 0.59 <sup>d</sup>   | 1.83 <sup>c</sup>   | 1.64 <sup>a</sup>    | 0.74 <sup>d</sup>    |
|                                        | G2        | 0.74 <sup>de</sup>   | 1.01 <sup>bc</sup>  | 2.31 <sup>bc</sup>  | 1.43 <sup>ab</sup>   | 1 <sup>bcd</sup>     |
|                                        | G3        | 1.36 <sup>abc</sup>  | 1.13 <sup>ab</sup>  | 1.86 <sup>c</sup>   | 0.90 <sup>bcd</sup>  | 1.33 <sup>ab</sup>   |
|                                        | Mean      | 1.01 <sup>b</sup>    | 0.91 <sup>b</sup>   | 2.00                | 1.33 <sup>a</sup>    | 1.02 <sup>b</sup>    |
| Participative selection varieties (PS) | G4        | 1.19 <sup>abcd</sup> | 0.87 <sup>bcd</sup> | 2.15 <sup>bc</sup>  | 1.20 <sup>abc</sup>  | 1.42 <sup>a</sup>    |
|                                        | G5        | 1.27 <sup>abc</sup>  | 1.47 <sup>a</sup>   | 0.89 <sup>d</sup>   | 1.20 <sup>abc</sup>  | 1.16 <sup>abc</sup>  |
|                                        | G6        | 1.50 <sup>ab</sup>   | 1.08 <sup>abc</sup> | 3.15 <sup>a</sup>   | 0.84 <sup>cd</sup>   | 1.24 <sup>ab</sup>   |
|                                        | Mean      | 1.32 <sup>a</sup>    | 1.14 <sup>a</sup>   | 2.07                | 1.08 <sup>ab</sup>   | 1.27 <sup>a</sup>    |
| Introduced varieties (IV)              | G7        | 1.01 <sup>bcd</sup>  | 0.73 <sup>cd</sup>  | 1.93 <sup>c</sup>   | 1.08 <sup>abcd</sup> | 1.05 <sup>abcd</sup> |
|                                        | G8        | 1.68 <sup>a</sup>    | 0.98 <sup>bcd</sup> | 2.16 <sup>bc</sup>  | 0.94 <sup>bcd</sup>  | 0.97 <sup>bcd</sup>  |
|                                        | G9        | 0.66 <sup>e</sup>    | 1.28 <sup>ab</sup>  | 2.44 <sup>abc</sup> | 0.77 <sup>cd</sup>   | 0.83 <sup>cd</sup>   |
|                                        | G10       | 0.73 <sup>de</sup>   | 0.96 <sup>bcd</sup> | 2.97 <sup>ab</sup>  | 0.57 <sup>d</sup>    | 1.00 <sup>abcd</sup> |
|                                        | Mean      | 1.02 <sup>b</sup>    | 0.99 <sup>ab</sup>  | 2.38                | 0.85 <sup>b</sup>    | 0.96 <sup>b</sup>    |
| LSD (P<0.05)_type of selection         |           | ***                  | *                   | NS                  | *                    | *                    |
| LSD (P<0.05)_varieties                 |           | ***                  | **                  | ***                 | **                   | **                   |
| Total mean                             |           | 1.11                 | 1.01                | 2.18                | 1.08                 | 1.08                 |

\*P<0.05, \*\*P<0.01 and \*\*\* P<0.001, NS: Not significant; LSD: Least significant difference.

**Table 4:** Analysis of variance morpho-agronomic traits of three breeding pathways (ten chickpea varieties) across six environments.

| Source of variation    | Degrees of freedom                       | Sum of squares | Mean square | F-test |
|------------------------|------------------------------------------|----------------|-------------|--------|
| <b>Line</b>            | <b>Plant height (cm)</b>                 |                |             |        |
| Breeding pathways (BP) | 2                                        | 243.31         | 121.66      | ***    |
| Environment (E)        | 5                                        | 2134.33        | 426.87      | ***    |
| BP × E                 | 10                                       | 498.43         | 49.84       | ***    |
| <b>Trait</b>           | <b>Number of primary branches</b>        |                |             |        |
| Breeding pathways (BP) | 2                                        | 1.45           | 0.72        | *      |
| Environment (E)        | 5                                        | 15.32          | 3           | ***    |
| BP × E                 | 10                                       | 2.97           | 0.29        | NS     |
| <b>Line</b>            | <b>Stem length to the first pod (cm)</b> |                |             |        |
| Breeding pathways (BP) | 2                                        | 239.82         | 119.91      | ***    |
| Environment (E)        | 5                                        | 1831.77        | 366.35      | ***    |
| BP × E                 | 10                                       | 167.8          | 16.78       | **     |
| <b>Trait</b>           | <b>100-seed weight (g)</b>               |                |             |        |
| Breeding pathways (BP) | 2                                        | 5686.5         | 2843.24     | ***    |
| Environment (E)        | 5                                        | 2457           | 491.4       | ***    |
| BP × E                 | 10                                       | 2055           | 205.5       | ***    |
| <b>Line</b>            | <b>Grain yield (t·ha<sup>-1</sup>)</b>   |                |             |        |
| Breeding pathways (BP) | 2                                        | 0.27           | 0.13        | NS     |
| Environment (E)        | 5                                        | 26.08          | 5.21        | ***    |
| BP × E                 | 10                                       | 3.24           | 0.32        | *      |
| <b>Trait</b>           | <b>Harvest index</b>                     |                |             |        |
| Breeding pathways (BP) | 2                                        | 0.00893        | 0.004466    | NS     |
| Environment (E)        | 5                                        | 0.48           | 0.096822    | ***    |
| BP × E                 | 10                                       | 0.4            | 0.040322    | ***    |

\*P<0.05, \*\*P<0.01, \*\*\*P<0.001 significance.

G×E interaction. The first principal component (PC1) explained 60.3% of the interaction variance, reflecting a clear environmental gradient and well-structured response.

The AMMI analysis identified G6-PS, G8-IV and G2-LV as the most productive varieties. G8-IV, G4-PS and G7-LV exhibited the highest stability values (Table 6). Considering both high yield and low instability, the introduced variety G8-IV emerged as the best-performing and most broadly adapted genotype, followed by G4-PS and G2-LV.

Regarding environments, E4 was clearly separated along the negative side of PC1, reflecting specific conditions highlighting the performance of certain varieties (Fig 3). E2, E5 and E6 were positioned on the positive side of PC1, representing moderately differentiated environments favorable to other genotypic profiles. E1 and E3, near the center of PC1, generated fewer G×E interactions and represented average environmental conditions.

The nominal stability plot (Fig 4) provided complementary insight: varieties with steep slopes (G6-PS, G10-IV, G5-PS) exhibited strong specific interactions, whereas G8-IV, G2-LV and G4-PS displayed more regular responses indicative of stability.

Fig 4 also highlighted significant environmental differences. E4 generated pronounced variation among varieties, reflecting strong specific interactions, while E1 and E3 elicited more homogeneous responses, identifying them as reference environments. E2, E5 and E6 showed

intermediate structuring, differentiating certain varieties while maintaining overall consistency.

Finally, the GGE biplot (Fig 5) illustrated environmental structure. E5, with a long vector, was the most discriminating environment.

The evaluation of performance and stability of chickpea genotypes derived from different selection methods under contrasting environments is essential for identifying varieties with high resilience potential. AMMI and GGE biplot analyses proved effective in characterizing genotype × environment (G×E) interactions, facilitating the selection of varieties that combine high productivity and stability, key traits for adaptation to semi-arid conditions with increasingly irregular and limited rainfall. Environmental factors accounted for the largest portion of variation (over 25% of total variation), followed by G×E interactions and genotypic differences, a pattern reported in recent studies (Istanbuli *et al.*, 2025; Eskezia *et al.*, 2025). This confirms the predominant influence of the environment on chickpea grain yield, with GEI playing a secondary but significant role. These findings align with Din *et al.* (2024), who emphasized the contribution of both environmental and G×E effects to the stability and adaptation of Desichickpea genotypes.

The contrasting performance of both homogeneous and heterogeneous varieties across environments suggests that stability is not solely determined by genetic background. Local varieties, often heterogeneous and co-

**Table 5:** Analysis of variance of main effect and genotype × environment interactions for chickpea grain yield (t·ha<sup>-1</sup>).

| Source of variation | Degrees of freedom (df) | Sum of squares (SS) | Mean squares (MS) | F value (F) | Probability (>F) | Proportion (%) |
|---------------------|-------------------------|---------------------|-------------------|-------------|------------------|----------------|
| ENV                 | 5                       | 92.52               | 18.50             | 63.77       | ***              | -              |
| REP(ENV)            | 12                      | 3.48                | 0.29              | 0.75        | NS               | -              |
| GEN                 | 9                       | 7.51                | 0.83              | 2.18        | *                | -              |
| GEN:ENV             | 45                      | 57.18               | 1.27              | 3.32        | ***              | -              |
| PC1                 | 13                      | 11.48               | 0.88              | 2.31        | **               | 60.3           |
| Residuals           | 468                     | 178.87              | 0.38              | -           | -                | -              |
| Total               | 584                     | 358.63              | 0.61              | -           | -                | -              |

\* $P < 0.05$ , \*\* $P < 0.01$  and \*\*\* $P < 0.001$ , NS: Not significant; LSD: Least significant difference; ENV: Environment; GEN: Genotype (variety).

**Table 6:** Mean chickpea grain yield and AMMI analysis of the ten varieties tested across six environments.

| Varieties | Type of selection | Mean grain yield (t·ha <sup>-1</sup> ) | ASV  | rASV | YSI | rYSI | GSI |
|-----------|-------------------|----------------------------------------|------|------|-----|------|-----|
| G8        | IV                | 1.32                                   | 0.30 | 2    | 4   | 2    | 4   |
| G4        | PS                | 1.29                                   | 0.27 | 1    | 5   | 4    | 5   |
| G2        | LV                | 1.32                                   | 0.58 | 4    | 7   | 3    | 7   |
| G6        | PS                | 1.49                                   | 1.72 | 8    | 9   | 1    | 9   |
| G3        | LV                | 1.29                                   | 0.73 | 5    | 10  | 5    | 10  |
| G7        | IV                | 1.08                                   | 0.37 | 3    | 13  | 10   | 13  |
| G1        | LV                | 1.15                                   | 0.99 | 7    | 14  | 7    | 14  |
| G9        | IV                | 1.14                                   | 0.88 | 6    | 14  | 8    | 14  |
| G10       | IV                | 1.26                                   | 2.13 | 9    | 15  | 6    | 15  |
| G5        | PS                | 1.12                                   | 2.79 | 10   | 19  | 9    | 19  |


ASV: AMMI stability value; rASV: Rank of each genotype according to AMMI stability value; YSI: Yield stability index; rYSI: Rank of each genotype according to yield stability index; GSI: Genotype stability index.

adapted, continue to evolve under local conditions, whereas introduced varieties, developed through intensive breeding in foreign environments, display distinct morphological and agronomic traits.


The analysis across six experimental sites confirmed significant genetic variability and underscored the importance of G×E interactions in chickpea agronomic performance (Jagan *et al.*, 2023; Ligarreto-Moreno and Pimentel-Ladino, 2022). Environments E1, E4 and E5 were the most discriminating, but only E4 and E5 displayed high productive potential, supporting the relevance of current chickpea zoning and multi-location breeding strategies (Pérez *et al.*, 2005). This highlights the need to consider agro-climatic specificity rather than generalizing

performance across apparently similar zones (Srivastava *et al.*, 2024).


Contrary to common assumptions, varieties introduced through international programs such as ICARDA are not inherently less plastic than local varieties. For instance, G8 (FLIP 93-93C, Ghab 4), selected in the WANA region, exhibited both high yield and stability, confirming its adaptation to Algeria's diverse semi-arid conditions. This observation agrees with Meddi *et al.* (2010); Annicchiarico (2017) and Dhuria and Babbar (2019), demonstrating that varieties developed for Mediterranean arid zones can combine high yield potential with broad adaptation. Similarly, the locally participatory-selected variety G4 displayed substantial plasticity across environments,



**Fig 3:** AMMI1 biplot for 10 chickpea varieties across six environments for grain yield.



**Fig 4:** Nominal yield plot based on AMMI1 scores and mean grain yield of chickpea varieties.



**Fig 5:** PCA Biplot of grain yield performance of ten chickpea varieties across six environments.

challenging the traditional dichotomy between plastic local varieties and specialized improved varieties.

Selecting genotypes such as G8-IV, G4-PS and G2-LV, which perform stably across most environments, provides a strong foundation for breeding broadly adapted varieties. These genotypes represent valuable resources for addressing climate change challenges, including drought, salinity and increasing precipitation variability (Bessah *et al.*, 2023; Djellouli and Laborde, 2022; Khoshro and Maleki, 2025).

Joint AMMI and GGE analyses revealed that no single site encompassed the full range of constraints and opportunities in semi-arid zones. Environments E5 and E4 proved strategic for selection, offering a balanced compromise between experimental efficiency and representativeness by capturing both general and specific adaptations of chickpea varieties (Yan and Rajcan, 2002). Other sites showing redundancy or low discriminatory power could be excluded in an optimized experimental design (Gauch and Zobel, 1997).

## CONCLUSION

Promising candidates, G8-IV, G4-PS and G2-LV, appear to be for direct use by farmers or breeding stocks in selection programmes. Their contrasting profiles, derived from different breeding methods, underscore the importance of diversifying the genetic sources for chickpea breeding. Multi-environment analysis showed that the E5 and E4 environments are particularly informative for breeding. These elements reinforce the relevance of flexible agro-ecological zoning for chickpeas in Algeria, enabling breeding to be oriented towards greater resilience of farming systems in the face of climatic variability.

## ACKNOWLEDGEMENT

This study was carried out in collaboration between Algeria's National Institute of Agronomic Research (INRAA) and the Algerian Interprofessional Office of Cereals and Dry Vegetables (OAIC). We sincerely thank Mrs. Dalila Boukecha (INRAA) for her technical assistance and guidance and the farmers who provided their land for the field trials.

## Disclaimers

The views and conclusions expressed in this article are solely those of the authors and do not necessarily represent the views of their affiliated institutions. The authors are responsible for the accuracy and completeness of the information provided, but do not accept any liability for any direct or indirect losses resulting from the use of this content.

## Informed consent

All the experimental procedures and handling techniques were approved by the National Higher School of Agronomy (ENSA), Laboratory of Integrated Improvement of Plant Production.

## Conflict of interest

The authors declare no conflicts of interest regarding the publication of this article. No funding or sponsorship influenced the conduct of the study, data collection, analysis, decision to publish, or manuscript preparation.

## REFERENCES

- Alrteimei, H.A., Ash'aari, Z.H., Muharram, F.M. (2022). Last decade assessment of the impacts of regional climate change on crop yield variations in the mediterranean region. *Agriculture*. **12**: 1787. doi: org/10.3390/agriculture12111787.

- Annicchiarico, P. (2017). Feed legumes for truly sustainable crop-animal systems. *Italian Journal of Agronomy*. **12**(2): 151-160. doi: 10.4081/ija.2017.880.
- Annicchiarico, P. (2020). Advances in the Definition of Adaptation Strategies and Yield Stability Targets in Plant Breeding. In: Quantitative Genetics, Genomics and Plant Breeding. [Kang, M.S., Basford, K.E. (Eds)], CABI. (2<sup>nd</sup> ed, pp 194-209). doi: 10.1079/9781789240214.0194.
- Arriagada, O., Cacciuttolo, F., Cabeza, R.A., Carrasco, B., Schwember, A.R. (2022). A comprehensive review on chickpea (*Cicer arietinum* L.) breeding for abiotic stress tolerance and climate change resilience. *Int. J. Mol. Sci.* **23**: 6794. doi: org/10.3390/ijms23126794.
- Baite, M.S., Dubey, S.C., Upadhyay, B.K. (2017). Genetic diversity of *Ascochyta rabiei* causing blight of chickpea in India. *Research Journal of Biotechnology*. **12**(8): 29-37.
- Bessah, E., Daryanto, S., Tsegaye, T. (2023). Climate-smart breeding for dry lands: Pathways for adaptation. *Frontiers in Plant Science*. **14**: 1178190.
- Bessaoud, O. and Montaigne, E. (2009). Algerian agriculture put to the test of sustainable development. *CIHEAM Options Méditerranéennes*. **64**: 51-91.
- Bouri, A., Mahdad, M.Y., Sidhoum, M., Gaouar, S.B.S. (2019). Effect of the genotype  $\times$  environment interaction on the main components of yield in chickpea (*Cicer arietinum* L.) in tlemcen, Algeria. *Gen. Biodv. J.* **3**(1): 63-69.
- Dhuria, N., Babbar, A. (2019). Genotype  $\times$  Environment (G $\times$ E) interaction and stability analysis of diverse elite *kabuli* chickpea lines for phenological and yield related traits under normal and heat stress environment. *Legume Research*. **44**(1): 31-35. doi: 10.18805/LR-4064.
- Din, A., Gul, R., Khan, H., Garcia-Abadillo Velasco, J., Persa, R., Isidro y Sánchez, J., Jarquin, D. (2024). Assessing the genotype-by-environment g $\times$ e interaction in desi chickpea via the bayesian additive main effects and multiplicative interaction model. *Agriculture*. **14**: 215. doi: org/10.3390/agriculture14020215.
- Djellouli, F. (2025). Assessing agricultural and hydrological drought trends in Algeria's semi-arid basins. *Pure and Applied Geophysics, Springer*. doi: org/10.1007/s00024-025-03773-4.
- Djellouli, Y. and Laborde, S. (2022). Assessment of climate variability and trends in the Algerian semi-arid region: Implications for water management. *Journal of Arid Environments*. **196**: 104676.
- Eskezia, A., Nakachew, K., Tadesse, M., Kassa, M. (2025). Genotype-by-environment interaction and yield stability of *Kabuli* chickpea (*Cicer arietinum* L.) in Northern Ethiopia. *Legume Science*. **7**(2): 1-15. doi: org/10.1002/leg3.70038.
- Farooq, M., Gogoi, N., Barthakur, S., Baroowa, B., Bharadwaj, N., Alghamdi, S.S., Siddique, K.H. (2017). Drought stress in grain legumes during reproduction and grain filling. *J. Agron. Crop Sci.* **203**: 81-102. doi: abs/10.1111/jac.12169.
- Gauch, H.G. and Zobel, R.W. (1997). Identifying mega-environments and targeting genotypes. *Crop Science*. **37**(2): 311-326.
- Gauch, H.G., Piepho, H.P., Annicchiarico, P. (2008). Statistical analysis of yield trials by AMMI and GGE: Further considerations. *Crop Science*. **48**: 866-889.
- Istanbuli, T., Alsamman, A.M., Al-Shamaa, K., Abu Assar, A., Adlan, M., Kumar, T., Tawkaz, S., Hamwieh, A. (2025). Selection of high nitrogen fixation chickpea genotypes under drought stress conditions using multi-environment analysis. *Front. Plant Sci.* **16**: 1490080. doi: 10.3389/fpls.2025.1490080.
- Jagan, M.R.P., Sandhyakishore, N., Srinivasan, S., Sandeep, S., Praveen, G., Neelima, G., Kumar, G.A. (2023). AMMI and GGE stability analysis of drought tolerant chickpea (*Cicer arietinum* L.) genotypes for target environments. *Legume Research*. **46**(9): 1105-1116. doi: 10.18805/LR-5156.
- Khan, A.M., Ali, Z., Mohsin, A.U., Soomro, A.A., Hussain, K., Rafiq, M., Hussain, N. et al. (2024). Genotype  $\times$  environment interactions analysis for chickpea grain yield and related traits by a mixed model approach. *Discovery Agriculture*. **2**: 90. doi: org/10.1007/s44279-024-00110-6.
- Kharrrat, M., Ben Salah, H., Maatougui, M.H. (2023). Study on the food and fodder legume sector in North Africa. *Tunis*. 72 pages. doi: 10.4060/cc0460fr.
- Khoshro, H.H., Maleki, H.H. (2025). Clarifying interactions between genotype and environment and management in chickpea by focusing on plant and soil attributes. *Sci. Rep.* **15**: 11401. doi: org/10.1038/s41598-025-95807-z.
- Laxuman, Avinalappa, H., Sidramappa, Kuchanur, P.H., Kumar, K.S., Kumar, G.A. (2025). Seed yield stability assessment of chickpea genotypes through AMMI and biplot analysis. *Legume Research*. **48**(2): 218-223. doi: 10.18805/LR-4911.
- Ligarreto-Moreno, G., Pimentel-Ladino, C. (2022). Grain yield and genotype  $\times$  environment interaction in bean cultivars with different growth habits. *Plant Production Science*. **25**(2): 232-241.
- Meddi, M., Meddi, H., Hubert, P. (2010). Impact of climate change on rainfall and runoff in the northwest of Algeria. *Hydrological Sciences Journal*. **55**(2): 316-332.
- Pandey, P.R., Gaur, P.M., Sajja, S.B. (2019). Pulses value chain development for achieving food and nutrition security in South Asia: Current status and future prospects. *SAARC Agriculture Centre*. p 181.
- Panigrahi, S., Kumar, U., Swami, S., Singh, Y., Balyan, P., Singh, K.P., Dhankher, O.P., Varshney, R.K., Roorkiwal, M., Amiri, K.M., Mir R.R. (2024). Meta QTL analysis for dissecting abiotic stress tolerance in chickpea. *BMC Genomics*. **25**(1): 439. doi: 10.1186/s12864-024-10336-9. PMID: 38698307; PMCID: PMC11067088.
- Pérez, P., Morcuende, R., Martín del Molino, I., Martínez-Carrasco, R. (2005). Reaction of chickpea lines to drought in different environmental conditions. *Agronomie*. **25**(6): 447-455.
- Ponsiva, S.T., Senthilkumar, N., Vanitha, J., Anandan, A., Satheeskumar, K., Mahendran, R., Thirugnanakumar, S. (2024). Detection of superior rice genotypes and yield stability using AMMI and MTSI models. *Electronic Journal of Plant Breeding*. doi: 10.37992/2024.1502.057.
- Raggi, L., Tiranti, B., Negri, V. (2019). Italian landraces of chickpea (*Cicer arietinum* L.): An evaluation of their agronomic performances and genetic diversity. *Agronomy*. **9**(8): 438. doi: 10.3390/agronomy9080438.

- Saraf, C.S., Rupella, O.P., Yadav, R.L., Shivkumar, B.G., Bhattarai, S. (1998). Biological nitrogen fixation and residual effects of winter grain legumes in rice and wheat cropping system of indo-gangetic plain. *Oxford and IHB Publishing*. pp 14-30.
- Shekhawat, H.V.S., Meena, V.K., Choudhary, K., Sharma, B.P., Sharma, J.K., Meena, S.C. (2024). Assessment of genetic stability in chickpea varieties through GGE and AMMI analyses. *Legume Research*. doi: 10.18805/LR-5246.
- Singh, D., Yadav, A., Kaur, G. (2020). Genotype  $\times$  environment interaction and stability analysis in chickpea (*Cicer arietinum* L.) for yield and its component traits. *Legume Research*. **43(1)**: 30-36.
- Srivastava, A.K., Mondal, B., Jha, U.C., Singh, A., Biradar, R.S., Praween, N., Singh, N., Dixit, G.P. and Kumar, Y. (2024). Selecting stable chickpea genotypes under rainfed cultivation using GGE biplot analysis. *Legume Research*. **47(12)**: 2141-2146. doi: 10.18805/LR-4832.
- Street, K., Bari, A., Mackay, M., Amri, A. (2016). How the Focused Identification of Germplasm Strategy (FIGS) is used to Mine Plant Genetic Resources Collections for Adaptive Traits. In: Enhancing Crop Genepool Use: Capturing Wild Relative and Landrace Diversity for Crop Improvement, [Maxted, N., Dulloo, M.E., Ford-Lloyd, B.V. (Eds)], CABI. (pp 54-63).
- Varshney, R.K., Roorkiwal, M., Amiri, K.M., Mir, R.R. (2024). Meta QTL analysis for dissecting abiotic stress tolerance in chickpea. *BMC Genomics*. **25(1)**: 439. doi: 10.1186/s12864-024-10336-9. PMID: 38698307; PMCID: PMC11067088.
- Yan, W. and Rajcan, I. (2002). Biplot analysis of test sites and trait relations of soybean in Ontario. *Crop Science*. **42(1)**: 11-20.
- Zobel, R., Wright, M.J., Gauch, H.G. (1988). Statistical analysis of yield trial. *Agronomy Journal*. **80**: 388-393.