



Cytotoxicity and Apoptosis Induction of *Lactobacillus acidophilus* Cell-free Supernatant against Human Colon Carcinoma Cell Line

Mustafa A. Hadid¹, Ahmed Flayyih Hasan², Mohammad F. Al-Halbosiy²,
Noorhan Sabih Al-Maliki⁴, Abdulwahid B. Al-Shaibani^{3,4},
Sameer Abed Mohammed⁵, Hany M. El-Wahsh⁶

10.18805/ag.RF-383

ABSTRACT

Background: *Lactobacillus acidophilus* is one of the promising probiotics that attracts the attention of the scientific community to research its therapeutic characteristics that can provide an option to fight colon cancer.

Methods: The cytotoxic effect of cell-free supernatant of *Lactobacillus acidophilus* (LACFS) was evaluated by using the MTT 3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide) test to measure cell viability and the high content screening assay (HCS) to detect apoptotic cell death.

Result: LACFS was found to possess the highest anticancer activity on Caco-2 at 50 µg/ml. NCM425 cells were more tolerant toward LACFS than the Caco-2 when the Half-maximal inhibitory concentration (IC₅₀) was 161.8 µg/ml on NCM425 compared to 30.63 µg/ml on Caco-2. When 6.25, 12.5, 25 and 50 µg/ml concentrations of LACFS were applied, a substantial drop in mitochondrial membrane potential occurred. A significant rise in cytochrome C release and total nuclear intensity was recorded at 25 and 50 µg/ml concentrations and the viable cell count was decreased at 12.5, 25 and 50 µg/ml, while membrane permeability was dropped at 50 µg/ml. The expression of caspase 3 was observed at 50 µg/ml only while caspase 9 expression was seen at 25 and 50 µg/ml which further indicates that the harmful effect on the Caco2 cells causes them to die through apoptosis.

Key words: Apoptosis, Colon cancer, Cytotoxicity, *Lactobacillus acidophilus*, Probiotics.

INTRODUCTION

The digestive system of humans is a receptacle for a diverse and dynamic microbial population that mostly contains bacteria that greatly impact the body in homeostasis and illness (Liśkiewicz *et al.*, 2018). The intestinal flora grows and flourishes within the first three years of human childhood (Gonzalez *et al.*, 2011). Probiotic bacteria include live microorganisms' dietary supplements that are classified as healthy ingredients. They improve a host animal's health by preserving gut microbial flora (Er *et al.*, 2015). Probiotics could help stop colorectal cancer through several strategies such as modifications in the microbiota of the intestine, inactivation of cancer-causing chemicals, improving the host's immunological response and anti-proliferative activities through apoptotic and cellular differentiation control, combat against harmful and putrefactive bacteria and undigested food fermentation (Uccello *et al.*, 2012).

The link between the *Lactobacillus*-enriched diet and the lower risk of colon cancer has been demonstrated (Goldin *et al.*, 1980). Among the most prevalent probiotics are the *Lactobacillus* species which possess health-improving characteristics that are firmly related to the reduction of allergy reactions, in addition to anti-tumor and anti-inflammatory properties (Ding *et al.*, 2018 ; Kahouli *et al.*, 2017 ; El-Deeb *et al.*, 2018). They induce programmed cell death and prevent colonic tumorigenesis (Escamilla

¹Department of Applied Pathological Analysis, College of Sciences, Al-Nahrain University, 10011 Baghdad, Iraq.

²Biotechnology Research Center, Al-Nahrain University, 10011 Baghdad, Iraq.

³Department of Biology, Al-Farabi University College, 10011 Baghdad, Iraq.

⁴College of Biotechnology, Al-Nahrain University, 10011 Baghdad, Iraq.

⁵Collage of Pharmacy, University of Anbar, Anbar, Iraq.

⁶Department of Marine Biology, Faculty of Marine Sciences, King Abdulaziz University, Saudi Arabia.

Corresponding Author: Ahmed Flayyih Hasan, Biotechnology Research Center, Al-Nahrain University, 10011 Baghdad, Iraq.
Email: ahmed_flayyih@nahrainuniv.edu.iq

How to cite this article: Hadid, M.A., Hasan, A.F., Al-Halbosiy, M.F., Al-Maliki, N.S., Al-Shaibani, A.B., Mohammed, S.A. and El-Wahsh, H.M. (2025). Cytotoxicity and Apoptosis Induction of *Lactobacillus acidophilus* Cell-free Supernatant against Human Colon Carcinoma Cell Line. *Agricultural Reviews*. 1-8. doi: 10.18805/ag.RF-383.

Submitted: 03-04-2025 **Accepted:** 31-05-2025 **Online:** 24-06-2025

et al., 2012). *Lactobacillus acidophilus* inhibited the growth of both colonic and gastric cancer cells (Lee *et al.*, 2004).

Cancer incidence is expected to increase from 11.5 billion in 2007 to 16.0 billion by 2030, driven by an ageing and rising global population (Hasan *et al.*, 2024). As ranking third, colorectal cancer (CRC) is considered to be among

the most frequent malignancies affecting both the colon and the rectum (Siegel *et al.*, 2019). Colorectal cancer is often managed using surgery, radiation, chemotherapy, or immunotherapy (Sadreddini *et al.*, 2019). In a study designed to examine how modified circulatory cells can be used as targeted carriers for anticancer drug delivery, traditional therapies might well be associated with unfavorable adverse effects, including chemoresistance, toxicities and cancer recurrence (Singh *et al.*, 2020).

MATERIALS AND METHODS

Lactobacillus acidophilus cell-free supernatant (LACFS)

A *Lactobacillus acidophilus* isolate was obtained from the Department of Biology/College of Science/ Baghdad University. It was activated twice by incubating its stock solution anaerobically for 24 and 48 hours at 37°C while subculturing it in MRS broth (Oxoid). One hundred and twenty ml of MRS broth containing 1×10^8 CFU/ml of a 24-hour-grown *L. acidophilus* culture was incubated anaerobically for 24 hours at 37°C. The broth was then centrifuged at 6000 rpm for 15 min and filtered through 0.22 µm pore-size filter papers (Microlab) before being lyophilized and stored at -20°C until it was needed (El-Mokhtar *et al.*, 2020).

MTT assay

Five concentrations (50, 25, 12, 5, 6.25 and 3.1 µg/ml) of the LACFS of *L. acidophilus* were prepared using an MTT kit (Intron Ltd) after 0.5 ml of distilled water was added to the lyophilized product. These concentrations were then tested on human colon adenocarcinoma Caco-2 (RRID: CVCL_0025) and NCM425 (RRID: CVCL_D876) cell cultures (Sigma-Aldrich). Cells were cultivated on RPMI1640 for each concentration at 37°C for 24 hours. They were then extracted using an EDTA/trypsin solution and resuspended in a medium that included 10-20% bovine serum albumin before being plated on a 96-well microtiter plate (All steps were done in triplicate.) The MTT assay was used to examine the anticancer potential activity and the results were read at 517 nm after 24 hours (Vinken *et al.*, 2015). The cell viability was measured by the formula below:

Percentage of cell viability =

$$\frac{\text{OD of sample} - \text{OD of blank}}{\text{OD of control} - \text{OD of blank}} \times 100\%$$

Cell death parameters

The high content screening (thermo fisher) approach was used to measure nuclear intensity, viable cell count, membrane permeability, cytochrome C release and mitochondrial membrane potential on Caco-2 in order to estimate cell death parameters at doses of 6.25, 12.5, 25 and 50 µg/ml. This method determines morphological traits and molecular reactivity after chemical treatment using fluorescent markers (Hoechst 33342, SYTOX Green/Red, mito tracker red CMXRos and Alexa Fluor 488/594-conjugated antibodies) (O'Brien *et al.*, 2017).

Caspase assay

The Caspase-Glo® 3/7 and Caspase-Glo® 9 Assay Kits (Promega) were used to measure Caspase-3 and Caspase-9 activities as stated by the manufacturer's instructions. Dimethyl Sulfoxide (DMSO) (Thermo Fisher, USA) and Tamoxifen (Dabur Pharma) were used as positive and negative controls, respectively (Hadid *et al.*, 2022).

Statistical analysis

ANOVA and Dunnett's multiple comparisons tests were used to analyze the data. Statistical significance was established at a p-value of less than 0.05. Each and every statistical analysis was conducted using GraphPad Prism version 8.4.3.

RESULTS AND DISCUSSION

Cytotoxicity of *Lactobacillus acidophilus* cell-free supernatant

The cytotoxic effect of *L. acidophilus* was evaluated on the colorectal cancer cell line (Caco-2) by using an MTT assay. This experiment aimed to investigate the viability of cells with various dosages of *L. acidophilus* LACFS on Caco-2 and it was shown in form of IC₅₀, which is the concentration required to reduce 50% of cell viability. A lesser IC₅₀ value indicates that the *L. acidophilus* supernatant is more effective in inhibiting cancer cell growth and vice-versa. Results indicated that *L. acidophilus* cell-free supernatant (LACFS) possessed cytotoxicity activity on colonic cancer with IC₅₀ = 30.63 µg/ml (Fig 1). Earlier studies had demonstrated that *L. acidophilus* has cytotoxic activity against Caco-2 (Hasan *et al.*, 2021 ; Shokryazdan *et al.*, 2017). When tested at different incubation times, cytotoxicity had been shown on HepG2, Chang and MDA-MB-231 cell cultures (20). On the other hand, cytotoxicity on NCM425 cells was drastically low compared to those of cancer cell line with IC₅₀ higher than on cancer cells by nearly 5 times; and it was insignificant with untreated (control) group when tested at all concentrations (Fig 1) (except at 25 and 50 µg/ml). This result agrees with earlier research (Nami *et al.*, 2014), which described that *L. acidophilus* CFS has a negligible effect on normal cell lines when tested for cytotoxic activity.

The MTT assay results showed that cytotoxicity of LACFS as expected, was dose-dependent, with a maximum cytotoxic impact reported at a concentration of 50 µg/ml on both Caco-2 and NCM425. As shown in (Fig 1), the cytotoxicity of LACFS on Caco-2 was significantly higher than on NCM425 at all tested concentrations as well as in untreated (control) cell lines. These results were close to those found by (Dehghani *et al.*, 2021; Alsadee *et al.*, 2024), which revealed that *L. rhamnosus* supernatant suppressed the development of HT-29 cancer cells in a dose- and time-dependent manner. These outcomes indicate a remarkable finding that *L. acidophilus* CFS has safer activity on normal colonic cells than on cancer cells when cell viability recorded 73.5% and 26.5% at 50 µg/ml respectively. Similar results were reported by (21) when *L. acidophilus* CFS appeared to be safer on the normal cell line.

Cell death caused by *Lactobacillus acidophilus* cell-free supernatant (LACFS)

The High Content Screening (HCS) assay was performed on the colonic cell line to further understand the cell-health indicators. HCS gives several criteria for determining the cytotoxicity of a substance at the single-cell level (Abraham *et al.*, 2008). The tested indicators were mitochondrial membrane potential (MMP) changes, cytochrome C discharge *via* mitochondria, the permeability of the cell membrane, viable cell count and morphology of nucleus intensity.

The MMP reduction causes cytochrome C to leak from mitochondria to cytosol so MMP along with cytochrome C

were tested by HCS assay and the results showed that there was an effect of LACFS on mitochondrial membrane potential (MMP) at concentrations of 25 and 50 µg/ml which had significant differences from that of untreated (control) group (Fig 2). At 25 and 50 µg/ml, cytochrome C levels rose considerably (Fig 3). In a study, by (Madempudi *et al.*, 2017; Al-Mashhadani *et al.*, 2024). MMP decreased significantly evidenced with rising in cytochrome C levels on colonic cancer cells after treatment with supernatant of *Bacillus coagulans*. The reduction of mitochondrial membrane potential (MMP) and the induction of apoptosis may be ascribed to the downregulation of the anti-apoptotic gene Bcl-2 and an elevation in Bax and caspase-9 which are




Fig 1: Cytotoxic effect of *Lactobacillus acidophilus* cell-free supernatant (LACFS) on Caco-2 and NCM425 cell lines.




Fig 2: Activity of *Lactobacillus acidophilus* cell-free supernatant (LACFS) on mitochondrial membrane permeability of Caco-2 cell line.

regarded as apoptotic signals that activate the endogenous route of apoptosis (Al-Saily *et al.*, 2019). Such processes induce alterations in the membrane permeability as a result of the expansion of the transition pore for mitochondrial permeability, which allows minute molecules and ions such as Ca^{2+} ions to pass through leading the respiratory chain

to decouple and release cytochrome C into the cytoplasm. Eventually, cytochrome C release triggers caspase-9, which leads to the creation of caspase-3 and the initiation of cysteine proteases, which are primarily responsible for the disintegration and digesting of the cell from within (Cha *et al.*, 2010; Tafani *et al.*, 2002). On cell membrane permeability (Fig 4) findings revealed that LACFS has activity only at high concentration (50 $\mu\text{g}/\text{mL}$) which was significant ($P\text{-value} \leq 0.05$) in comparison to the untreated (control) group. A previous study by reported a comparable impact of *L. acidophilus* on the permeability of the cells when experienced against the Caco-2 cell line. Modifications in cellular membrane permeability have been linked to apoptotic or toxic responses, in addition, the collapse of cell membrane integrity seems to be a typical morphological hallmark of severe cytotoxic effects (Hassan *et al.*, 2018).

Results (Fig 5) showed that *L. acidophilus* cell-free supernatant (LACFS) has a highly significant effect on the viable count of Caco-2 at concentrations of 25 and 50 $\mu\text{g}/\text{mL}$, compared to the untreated (control) group. The distinction in such survivability findings as contrasted to MTT experiments would be that in HSC the findings convey the impact of previous exposure to the drug for the assessment of morphology and cellular defects in cultures after only a few hours of exposure. However, in MTT, the tested cells were exposed to the agent for a prolonged period. Apoptosis is marked by chromatin condensation and DNA fragmentation, followed by a decrease in nuclear intensity as fragmentation advances (Khan *et al.*, 2010).

In this study breakdown of DNA was also seen by a substantial increase in overall nuclear intensity at 25 and 50 $\mu\text{g}/\text{mL}$ as shown in (Fig 6), while no effect was noticed at other treatments in comparison to the control (untreated)




Fig 3: Activity of *Lactobacillus acidophilus* cell-free supernatant (LACFS) on cytochrome c releasing of Caco-2 Cell Line.




Fig 4: Activity of *Lactobacillus acidophilus* cell-free supernatant (LACFS) on cell membrane permeability of Caco-2 Cell Line.

group. The nuclear intensity outcomes of (Kamil *et al.*, 2021). also showed an increase when dextran extracted from probiotics *Leuconostoc* was applied to the MCF7 cell line. As shown in (Fig 7), the impact of LACFS on caspase 3 was only significant at 50 $\mu\text{g/ml}$ concentration when it was

extremely significant at 25 and 50 $\mu\text{g/ml}$ concentrations on caspase 9 compared to DMSO (negative control) (Fig 8). It is worth noting that prior probiotic anti-cancer investigations have yielded similar outcomes. High doses of probiotic supernatant, cell walls and pellets might have a potent




Fig 5: Activity of *Lactobacillus acidophilus* cell-free supernatant (LACFS) on Caco-2 viable count.




Fig 6: Activity of *Lactobacillus acidophilus* cell-free supernatant (LACFS) on the total nuclear intensity of Caco-2 Cell Line.

anti-tumor impact (Desrouillères *et al.*, 2016 ; Rasheed *et al.*, 2025). Probiotic *L. delbrueckii*, for instance, suppresses the development of colorectal carcinoma cells SW620 and triggers death *via* a caspase 3-dependent internal

mechanism (Wan *et al.*, 2014). When *L. acidophilus* CICC 6074 concentrations were raised in comparison to the untreated control group, the quantity of caspase-3 and caspase-9 rose dramatically (Guo *et al.*, 2020; Ghiath *et al.*, 2025).




Fig 7: Effect of *Lactobacillus acidophilus* cell-free supernatant (LACFS) on caspase-3 expression of Caco-2 Cell Line.




Fig 8: Effect of *Lactobacillus acidophilus* cell-free supernatant (LACFS) on caspase-9 expression of Caco-2 Cell Line.

CONCLUSION

It is noteworthy that *Lactobacillus acidophilus* cell-free supernatant (LACFS) was cytotoxic on human colon cancer cells with lesser cytotoxicity on normal cells which may propose an alternative promising anticancer therapy. Also, LACFS promotes apoptosis in colonic cells as an antitumor combating mechanism.

Conflict of interest

All authors declare that they have no conflict of interest.

REFERENCES

- Abraham, V.C., Towne, D.L., Waring, J.F., Warrior, U. (2008). Burns, D.J. Application of a high-content multiparameter cytotoxicity assay to prioritize compounds based on toxicity potential in humans. *J. Biomol Screen*. Sep. 13(6): 527-537.
- Al-Mashhadani, T.A., Nafea, M.H., Obayes, K.R., Hussein, M.S. and Hasan, A.F. (2024). Biochemical effects of silver nanoparticles prepared by chemical reduction method on male rat kidney functions and antioxidant defense systems. *Agricultural Science Digest*. 1-7. doi: 10.18805/ag.DF-657.
- Alsadee, A.S., Mohammed, A.S., Hasan, F.A. (2024). Hepatotoxicity of Oral Sub-chronic exposure to thiamethoxam and lambda cyhalothrin in rats. *Agricultural Reviews*. 45(4): 728-733. doi: 10.18805/ag.RF-350.
- Al-Saily, H., Al-Halbosi, M., Al-Hady, F. (2019). Cytotoxic and apoptotic effects of cyproterone acetate against cancer cells and normal cells. *Journal of biotechnology research center*. 2019. 13(1): 68-74.
- Cha, J., Kim, Y., Kim, J.Y. (2010). Essential oil and 1,8-cineole from *Artemisia lavandulaefolia* induces apoptosis in KB cells via mitochondrial stress and caspase activation. *Food Sci Biotechnol*. 2010. 19(1): 185-191.
- Dehghani, N., Tafvizi, F., Jafari, P. (2021). Cell cycle arrest and anti-cancer potential of probiotic *Lactobacillus rhamnosus* against HT-29 cancer cells. *BiolImpacts*. 2021. 11(4): 245-252.
- Desrouillères, K., Millette, M., Jamshidian, M., Maherani, B., Fortin, O., Lacroix, M. (2016). Cancer preventive effect of a specific probiotic fermented milk components and cell walls extracted from a biomass containing *L. acidophilus* CL1285, *L. casei* LBC80R and *L. rhamnosus* CLR2 on male F344 rats treated with 1,2-dimethylhydrazine. *J Funct Foods*. 2016 Oct 1. 26:373-384.
- Ding, C., Tang, W., Fan, X., Wu, G. (2018). Intestinal microbiota: A novel perspective in colorectal cancer biotherapeutics. *Onco Targets Ther*. 2018. 11: 74-79.
- El-Deeb, N.M., Yassin, A.M., Al-Madboly, L.A., El-Hawiet, A. (2018A). Novel purified *Lactobacillus acidophilus* 20079 exopolysaccharide, LA-EPS-20079, molecularly regulates both apoptotic and NF-KB inflammatory pathways in human colon cancer. *Microb Cell Fact*. 2018 Feb 21: 17(1).
- El-Mokhtar, M.A., Hassanein, K.M., Ahmed, A.S., Gad, G.F.M., Amin, M.M., Hassanein, O.F.E. (2020). Antagonistic activities of cell-free supernatants of lactobacilli against extended-spectrum β -Lactamase Producing *Klebsiella pneumoniae* and *Pseudomonas aeruginosa*. *Infect Drug Resist*. 2020.13: 543.
- Er, S., Koparal, A.T., Kivanç, M. (2015). Cytotoxic effects of various lactic acid bacteria on Caco-2 cells. *Turkish Journal of Biology*. 2015 Jan 27. 39(1): 23-30.
- Escamilla, J., Lane, M.A., Maitin, V. (2012). Cell-free supernatants from probiotic *Lactobacillus casei* and *Lactobacillus rhamnosus* GG decrease colon cancer cell invasion *in vitro*. *Nutr Cancer*. 64(6): 871-878.
- Ghiath, Y., Mtashar, B.A., AL-Zuhairy, N.A.H.S., Hussein, M.S. and Hasan, A.F. (2025). Interplaying correlation of some genetic and inflammatory factors among patients with polycythemia vera. *Asian Journal of Dairy and Food Research*. 1-6. doi: 10.18805/ajdfr.DRF-492.
- Goldin, B., Gorbach, S. (1980). Effect of *Lactobacillus acidophilus* dietary supplements on 1,2-dimethylhydrazine dihydrochloride-induced intestinal cancer in rats. *J. Natl Cancer Inst*. 64(2): 263-265.
- Gonzalez, A., Stombaugh, J., Lozupone, C., Turnbaugh, P.J., Gordon, J.I., Knight, R. (2011). The mind-body-microbial continuum. *Dialogues Clin Neurosci*. 13(1): 55.
- Guo, Y., Zhang, T., Gao, J., Jiang, X., Tao, M., Zeng, X. *et al*. (2020). *Lactobacillus acidophilus* CICC 6074 inhibits growth and induces apoptosis in colorectal cancer cells *in vitro* and in HT-29 cells induced-mouse model. *J. Funct Foods*. 1. 75: 104290.
- Hadid, M.A., Al-Halbosi, M.M.F., Al-Shaibani, A.B. (2022). Anticancer and Cytotoxic Effect of Verotoxin 1 on Colon Cancer Cell Line. *Microbiology and Biotechnology Letters*. 50(3): 387-394.
- Hasan, A.F., Hameed, H.M., Hadid, M.A., Tousson, E. (2024). Impact of chia (*Salvia hispanica*) seeds extract on ehrlich ascites model induced kidney toxicity in female mice. *Asian Journal of Dairy and Food Research*. 43(4): 750-756. doi: 10.18805/ajdfr.DRF-397.
- Hasan, R.N. (2021). Cytotoxicity potential of local isolates of *Lactobacillus acidophilus* extracts on colon cancer cell line. *Annals of the Romanian Society for Cell*. 25: 6750-6765.
- Hassan, F.A. (2018). Cytotoxic, Antioxidant and Caspase-9 activity of 2-(3,5-dibromo-4-methoxyphenyl)3(3-mercapto-5(pyridine-4-Yl)4H1,2,4-triazol-4-yl)-thiazolidine-4-one against various human tumor cell lines. *Journal of Al-Nahrain University of Science*. 21(3): 32-38.
- Kahouli, I., Malhotra, M., Westfall, S., Moulay, A., Alaoui-Jamali, Prakash, P. (2017). Design and validation of an orally administered active *L. fermentum*-*L. acidophilus* probiotic formulation using colorectal cancer ApcMin/+ mouse model. *Appl Microbiol Biotechnol*. 101(5): 1999-2019.
- Kamil, W., Shalash, W.K., Al-Ezzy, R.M., Ahmaed, A.S. (2021). Antitumor activity of dextran produced from localized leuconostoc mesenteroides isolates. *Biochem Cell Arch*. 21(1): 2371- 2374.
- Khan, N., Mukhtar, H. (2010). Cancer Chemoprevention. *Comprehensive Toxicology*, Second Edition. 14: 417-431.
- Lee, J.W., Shin, J.G., Kim, E.H., Kang, H.E., Yim, I.B., Kim, J.Y. *et al*. (2004). Immunomodulatory and antitumor effects *in vivo* by the cytoplasmic fraction of *Lactobacillus casei* and *Bifidobacterium longum*. *J. Vet Sci*. 5: 41-48.
- Liśkiewicz, P.J., Pelka-Wysiecka, J., Wroński, M., Bąba-Kubiś, A., Samochowiec, J. (2018). Intestinal flora and the pathophysiology of depression and anxiety disorders- current state of the art and future perspectives. *Psychiatry*. 15(2): 70-76.
- Madempudi, R.S., Kalle, A.M. (2017). Antiproliferative effects of *Bacillus coagulans* unique IS2 in colon cancer cells. *Nutr Cancer*. 69(7): 1062-1068.

- Nami, Y., Abdullah, N., Haghshenas, B., Radiah, D., Rosli, R., Khosroushahi, A.Y. (2014). Probiotic potential and biotherapeutic effects of newly isolated vaginal *Lactobacillus acidophilus* 36YL strain on cancer cells. *Anaerobe*. 28: 29-36.
- O'Brien, P.J., Edvardsson, A. (2017). Validation of a multiparametric, high-content-screening assay for predictive/investigative cytotoxicity: Evidence from technology transfer studies and literature review. *Chem Res Toxicol*. 30(3): 804-829.
- Rasheed, S.S., Fadhil, R.Z., AL-Chalabi, S.M.M., Khalaf, R.A., Al Mahdawi, F.A. and Hasan, A.F. (2025). Assessment of quercetin influence on multiple physiological, biochemical and immunological parameters in diabetic rats. *Asian Journal of Dairy and Food Research*. 1-6. doi: 10.18805/ajdfr.DF-703.
- Sadreddini, S., Baradaran, B., Aghebati-Maleki, A., Sadreddini, S., Shanehbandi, D., Fotouhi, A. *et al.* (2019). Immune checkpoint blockade opens a new way to cancer immunotherapy. *J. Cell Physiol*. 234(6): 8541-8549.
- Shokryazdan, P., Jahromi, M.F., Bashokouh, F., Idrus, Z., Liang, J.B. (2017). Antiproliferation effects and antioxidant activity of two new *Lactobacillus* strains. *Brazilian Journal of Food Technology*. 21. doi: 10.1590/1981-6723.6416.
- Siegel, R., Miller, K., Jemal, A., Cancer, S. (2019). Cancer statistics. *Cancer Journal for Clinicians*. 69(1): 7-34.
- Singh, B., Mitragotri, S. (2020). Harnessing cells to deliver nanoparticle drugs to treat cancer. *Biotechnol Adv*. 42: 107339.
- Tafari, M., Karpnich, N., Hurster, K.A., Pastorino, J.G., Schneider, T., Russo, M.A., *et al.* (2002). Cytochrome c release upon Fas receptor activation depends on translocation of full-length bid and the induction of the mitochondrial permeability transition. *Journal of Biological Chemistry*. 277(12): 10073-10082.
- Uccello, M., Malaguarnera, G., Basile, F., Dagata, V., Malaguarnera, M., Bertino, G. *et al.* (2012). Potential role of probiotics on colorectal cancer prevention. *BMC Surg*. (SUPPL. 1).
- Vinken, M., Rogiers, V. (2015). Protocols in *in vitro* hepatocyte research. *Methods in Molecular Biology*. New York: Springer New York; 1-390 p.
- Wan, Y., Xin, Y., Zhang, C., Wu, D., Ding, D., Tang, L., *et al.* (2014). Fermentation supernatants of *Lactobacillus delbrueckii* inhibit growth of human colon cancer cells and induce apoptosis through a caspase 3-dependent pathway. *Oncol Lett*. 7(5): 1738-1742.