



Unravelling the Role of Phytohormones against Heat Stress in Maize

V. Manonmani¹, V. Revathy¹, S. Ambika², R. Paramasivam³, S. Kavitha¹,
M. Nivethitha⁴, M. Umanath⁵

10.18805/IJARE.A-6237

ABSTRACT

Background: Drought stress is a significant challenge for maize production, causing significant harm to crop growth and yield. In this experiment two set of pot culture experiment was conducted with three replication to assess the effect of phytohormone foliar spray on maize COH(M) 8.

Methods: This experiment study was conducted at the Seed Science and Technology, TNAU in the summer season during 2022 and 2023, respectively. One set of potted plants were kept under ambient temperature ($34\pm 2^{\circ}\text{C}$). Another set of potted plants were raised at elevated temperature of 42°C in the Open Top Chamber. The plants were sprayed with sodium nitroprusside (50 μM and 75 μM), brassinolides (0.2 ppm and 0.5 ppm) and salicylic acid (50 ppm and 75 ppm) at 40 and 47 days after sowing along with control.

Result: Foliar spraying with sodium nitroprusside 50 μM and salicylic acid 75 ppm recorded higher plant height, early flowering, pollen viability, chlorophyll content and enzyme activities (catalase, peroxidase and superoxide dismutase) compared to the control plants. As a result, these findings suggest that use of sodium nitroprusside 50 μM enhances plant defence mechanisms against heat stress in maize.

Key words: Abiotic stress, Enzymes, Heat stress, Maize, Phytohormones.

INTRODUCTION

Heat stress is a major abiotic factor that reduces productivity and quality, leading to significant economic losses (Sharma and Manjeet, 2020). Heat stress has a significant negative impact on plant growth, development and yield (Youldash *et al.*, 2020). When temperatures surpass a particular level, which is typically approximately 30°C for the majority of crops (Sattar *et al.*, 2020) plants undergo changes such as reduced plant height, root growth, leaf size, uptake of water, nutrients (Ray *et al.*, 2012), photosynthesis activity, change of source-sink relationships (Rennenberg *et al.*, 2006), respiration, reduce chlorophyll content, and disrupt the balance of carbon assimilation that can affect their productivity (Tiwari *et al.*, 2019). Heat stress has a significant impact on physiological processes such as seed germination and vigor, leaf enlargement, root growth, photosynthesis, reproductive development, rate of filling and, grain filling duration ultimately reducing yield and grain quality. Different proteins, membranes, RNA species, and cytoskeleton structures are affected differently by heat stress. It also modifies the efficiency of enzymatic reactions within the cell, which causes a barrier to major physiological processes and metabolic imbalance (Suzuki *et al.*, 2011). Heat stress also causes an oxidative burst, which results in overproduction of ROS that cause membrane lipid peroxidation, DNA damage, protein oxidation, and cell apoptosis enzyme inactivation and macromolecule damage (Iqbal *et al.*, 2022; Choudhary *et al.*, 2020) as well as disruption of cell elongation and differentiation (Bazzaz *et al.*, 2020). Rubisco becomes less active or inactive at high temperatures, which leads to a reduction in the fixing of carbon dioxide.

¹Department of Seed Science and Technology, Tamil Nadu Agricultural University, Coimbatore-641 003, Tamil Nadu, India.

²Department of Seed Science and Technology, SRM College of Agricultural Sciences, Chennai-603 203, Tamil Nadu, India.

³School of Social Sciences and Languages, Vellore Institute of Technology, Chennai-603 203, Tamil Nadu, India.

⁴Don Bosco Agriculture College, Arakkonam, Chennai-631 151, Tamil Nadu, India.

⁵Madras Institute of Development Studies, Chennai-600 020, Tamil Nadu, India.

Corresponding Author: S. Ambika, Department of Seed Science and Technology, SRM College of Agricultural Sciences, Chennai-603 203, Tamil Nadu, India. Email: ambikasingaram@gmail.com.

How to cite this article: Manonmani, V., Revathy, V., Ambika, S., Paramasivam, R., Kavitha, S., Nivethitha, M. and Umanath, M. (2025). Unravelling the Role of Phytohormones against Heat Stress in Maize. *Indian Journal of Agricultural Research*. **59(9)**: 1360-1367. doi: 10.18805/IJARE.A-6237.

Submitted: 04-04-2024 **Accepted:** 03-10-2024 **Online:** 12-11-2024

In maize during flowering and grain filling stage, temperature above 40°C will affect the productivity (Shiferaw *et al.*, 2011). The degree and duration of exposure also increases the effect on plant. Average optimum temperature is $30-34^{\circ}\text{C}$. Increase in temperature by 2°C reduces yield by 13%. Temperature above 35°C causes increased pollen sterility and therefore reduces the yield (Suwa *et al.*, 2010). High temperature negatively affects the reproductive stage by reducing the flower number, size and deformation of floral organs which results in reduction of the seed yield (Lobell *et al.*, 2011).

External application of various phytohormones like salicylic acid, brassinolide and sodium nitroprusside plays an important role in alleviating the heat stress by acting as messenger molecules and also plays a serious role in growth and development of plants. The hormonal cross talk coordinates the defence responses in plants to heat stress.

Foliar application of salicylic acid reduced the heat stress effects by membrane damage reduction through reduced leakage of electrolytes, malondialdehyde content and hydrogen peroxide content (Hameed and Ali, 2016). Salicylic acid (SA) is an endogenous plant growth regulator plays a vital role in plant growth, ion uptake, transport, interaction with other organisms and in the responses to environmental stress (Kuchlan and Kuchlan, 2023). Foliar application of brassinolides protects the plants through amplification responses to heat stress and saving the expression of plant growth promoters. Foliar application of sodium nitroprusside reduced heat stress by decreasing ion leakage and increasing the activity of anti-oxidant enzymes (Deng and Song, 2012) and also alleviate membrane lipid peroxidation damage under drought (Zhao *et al.*, 2024). The effectiveness of exogenously applied phytohormones at optimum concentration for protecting the crop against high temperature needs to be studied.

MATERIALS AND METHODS

This experiment study was conducted at the Seed Science and Technology, TNAU in the summer season during 2022 and 2023, respectively. The crop was raised in the pots in two sets. One set of potted plants were kept under ambient temperature ($34\pm 2^\circ\text{C}$) for the total duration upto the flowering stage (Fig 1a and 1b). Another set of potted plants were raised at the ambient temperature until the completion of flowering stage. Then the plants were exposed to the elevated temperature of 42°C in the Open Top Chamber for 10 days during the flowering stage. Nutrients and plant protection management were carried out as per the recommendation of TNAU crop production guide (NPK @ $175:90:90\text{ kg ha}^{-1}$). The crop was foliar sprayed with phytohormones using knapsack sprayer at two growth stages. The growth parameters (plant height) and physiological parameters (chlorophyll content (mg/g), relative water content (%), proline content (mg/g), catalase ($\mu\text{g of H}_2\text{O}_2/\text{g/minute}$), peroxidase activity (g tissue/min),

superoxide dismutase ($\text{U mg}^{-1}\text{ protein min}^{-1}$) were observed and the leaf samples were collected before spray and 3 days after second foliar spray and analysed for the enzyme activity. The mean value was calculated. The significance of data from several studies was determined using the “F” test, as described by (Panse and Sukhatme, 1957). Treatment details are given in Table 1.

RESULTS AND DISCUSSION

Heat stress is a critical concern for food crop yield in the changing climatic scenario and it is a great challenge for everyone in achieving food security (Chaturvedi *et al.*, 2017). Maize plants are vulnerable to heat stress (temperatures above 30°C) (Fig 2a and 2b).



Fig 1 (a): View of pot culture experiment at the ambient condition ($34\pm 2^\circ\text{C}$).



Fig 1 (b): View of pot culture experiment in the open top chamber (42°C).

Table 1: Treatment details.

Foliar spray with phytohormones	Concentration	Temperature stress	Stages of foliar spray
Control	-		-
Salicylic acid	50 ppm 75 ppm	Ambient condition ($34\pm 2^\circ\text{C}$)	40 and 47 days after sowing
Brassinolides	0.2 ppm 0.5 ppm		
Sodium nitroprusside	50 μM 75 μM	Open top chamber (42°C)	

Significant increase in the plant height was noticed due to foliar application of phytohormones. Among the foliar spraying treatments, maize plants sprayed with salicylic acid 75 ppm recorded the highest plant height of 185.5 cm and 188.6 cm which was on par with sodium nitroprusside 50 μ M (180.7 cm and 185.6 cm at 60 and 90 DAS respectively) whereas the lowest plant height was observed for the control which recorded 160.5 cm 164.3 cm at 60 and 90 DAS respectively (Table 2). Salicylic acid stimulated cell division in apical meristem in plants (Shakirova *et al.*, 2003). Foliar application of salicylic acid increased the plant growth in *Cicer arietinum* (Hayat *et al.*, 2012) and *Rosmarinus officinalis* (El-Esawi *et al.*, 2017). Salicylic acid controls the functions of plant in a concentration dependent manner where the lower concentrations of the salicylic acid induced the functions of the plant while higher concentrations inhibited the plant functions. Sodium nitroprusside significantly increased growth and its related parameters compared to the control at all stages of the crop growth (Hajhashemi *et al.*, 2021).

Flowering was delayed in the open top chamber (42°C) when compared to the ambient condition (34°C). Plants foliar sprayed with sodium nitroprusside 50 μ M and salicylic acid 75 ppm (52 days) flowered 4 days earlier when compared to control (56 days) at the ambient condition (34°C) (Table 2). Plants foliar sprayed with sodium nitroprusside 50 μ M and salicylic acid 75 ppm (54 days) flowered early by 4 days when compared to control (58 days) at the open top chamber (42°C). Days to 50% flowering was 55 days in plants foliar sprayed with sodium nitroprusside 50 μ M which was on par with salicylic acid 75 ppm foliar spraying whereas control seeds have taken 58 days to attain 50% flowering at the ambient condition (34°C). The treatments such as sodium nitroprusside 50 μ M and salicylic acid 75 ppm taken less days for 50% flowering compared to control at the open top chamber (42°C). Salicylic acid application enhanced the flowering (Aghdam *et al.*, 2016). Nitric oxide (NO), which SNP (sodium nitroprusside) is a donor of, has been demonstrated to be involved in the way plants react to environmental stressors, such as heat stress (Qiao and Fan, 2008). Application of SNP to plants can mitigate certain adverse consequences of thermal stress, such as diminishing oxidative stress, preserve the integrity of the plant cell membrane, enhancing photosynthesis and elevating antioxidant activity and enzymes involved in stress signalling (Yu *et al.*, 2013; Siddiqui *et al.*, 2017a; Kharbech *et al.*, 2020).

The maximum pollen viability was observed in the plants foliar sprayed with sodium nitroprusside 50 μ M (88%) whereas the minimum was observed in control plants (83%) at the ambient condition (34°C) (Table 2). However, the pollen viability was reduced at the elevated temperature (42°C) in the open top chamber. Sodium nitroprusside 50 μ M (80%) applied plants recorded higher pollen viability when compared to control (68%). Pollen abortion caused by different stresses is always a result of their abnormal

development of tapetum (Trost, 2014), which directly contacts the male gametophyte and plays a serious role in the development and maturation of microspores. Pollen viability, stigma receptivity and ovule viability was reduced in plants exposed to heat stress but increased with the exogenous application of sodium nitroprusside as a consequence of improvement in leaf and anther function to significantly increase physiological function and yield in heat stressed lentil (Esim and Atici, 2014; Sita *et al.*, 2021).

The plants foliar sprayed with salicylic acid 75 ppm and sodium nitroprusside 50 μ M registered maximum proline content (360 mg/g) when compared to control 320 mg/g at the ambient condition. Similar trend was observed for the plants kept at the elevated temperatures in the open top chamber. However, the proline content was increased when compared to the plants kept at the ambient condition.

The plants foliar sprayed with sodium nitroprusside 50 μ M registered maximum proline content (420) followed by salicylic acid 75 ppm (410). The minimum proline content was observed in control (340) (Fig 3). Aftab *et al.* (2010) reported that salicylic acid has been involved in regulation of essential plant biological processes like nitrogen and




Fig 2 (a): Effect of heat stress on maize plant.




Fig 2 (b): Influence of phytohormones spray on maize plant under heat stress.

proline metabolism, photosynthesis, glycine betaine production, multiple antioxidant defense mechanism and water relations in plants under stress situations. Tufail *et al.* (2013) suggested that action of salicylic acid under heat stress increased the production of proline which improved the osmotic potential allowing the plant for increased water uptake which results in positive effect of on stomatal complex and photosynthetic process which leads to improved efficiency of photosystem II and activity of Rubisco increases that collectively resulted in improved photosynthesis. Salicylic acid modifies both the osmolytes and metabolites production and also the nutrient status of the plants and plays defensive role under the abiotic stress situations (Ghani *et al.*, 2015).

The plants foliar sprayed with sodium nitroprusside 50 μ M (0.947 mg/g) at 40 and 47 DAS recorded the maximum chlorophyll content over control in the open top chamber (Table 3). This was supported by Mittler (2002) that application of sodium nitroprusside led to higher levels of chlorophylls a and b due to less degradation of chlorophyll under heat stress. The application of sodium nitroprusside significantly improved chlorophyll contents and transpiration rate, which could be attributed to the inhibition of reactive oxygen species production or to maintaining photosynthesis stability under stress conditions (Tian *et al.*, 2015; Muthulakshmi *et al.*, 2017). Exogenous application of sodium nitroprusside treatment improved relative water content, leaf chlorophyll concentration, and electrolyte content in heat-stressed *Zingiber officinale* (Li *et al.*, 2013). The restoration of net photosynthetic rate (Haldimann and Feller, 2004) and recovery from chlorophyll bleaching were detected during sodium nitroprusside treatment by delayed reduction of photosynthetic pigment in *Quercus pubescens*, *Triticum aestivum*, *Vicia faba*, *Solanum Lycopersicon* and

Oryza sativa (Yang *et al.*, 2011). Heat stress is accompanied by oxidative stress as noticed by the burst of hydrogen peroxide in cells after high temperature exposure thus reducing yield (Sharma *et al.*, 2020).

The present study proved that the plants foliar sprayed with salicylic acid 50 and 75 ppm recorded the maximum Relative Water Content (18%) (Table 3) at the ambient condition. Plants foliar sprayed with sodium nitroprusside 50 μ M and salicylic acid 75 ppm (14%) registered increased relative water content in the open top chamber (42°C) when compared to control (11%). Similar findings were observed by Li *et al.* (2013) that exogenous application of sodium nitroprusside retrieved relative water content and chlorophyll concentration of leaf in heat-stressed *Zingiber officinale*.

The maximum catalase activity was observed in the leaves foliar sprayed with sodium nitroprusside 50 μ M (127.3 μ g of H₂O₂/g/minute) and minimum was observed in the control (121.0 μ g of H₂O₂/g/minute) at the ambient condition (Table 4). In the open top chamber (42°C) the maximum catalase activity was observed with sodium nitroprusside 50 μ M (132.1 μ g of H₂O₂/g/minute) and minimum was observed in the control (123.2 μ g of H₂O₂/g/minute). The peroxidase and superoxide dismutase activity was maximum for sodium nitroprusside 50 μ M applied plants (145.4 g tissue/min, 1.47, respectively) whereas the minimum peroxidase activity was observed in the control plants (127.3 g tissue/min, 1.12, respectively) at the ambient condition (34°C). The plants foliar sprayed with sodium nitroprusside 50 μ M registered maximum superoxide dismutase activity (1.60) followed by salicylic acid 75 ppm (1.40) and the minimum superoxide dismutase activity was observed in control (1.10) at open top chamber (Fig 4). By encouraging the expression of

Table 2: Effect of foliar spray with phytohormones on plant height (cm), days to first flowering and 50% flowering in maize COH (M) 8 under temperature stress.

Foliar spray treatments	Plant height		Ambient condition (34°C)		Open top chamber (42°C)		Ambient condition (34°C)	Open top chamber (42°C)
	60 DAS	90 DAS	Days to first flowering	Days to 50% flowering	Days to first flowering	Days to 50% flowering	Pollen viability (%)	Pollen viability (%)
Control	160.5	164.3	56	58	58	61	83 (65.65)	68 (55.55)
Salicylic acid 50 ppm	167.4	169.2	54	56	55	57	86 (68.03)	72 (58.05)
75 ppm	185.5	188.6	52	55	54	56	87 (68.87)	75 (60.00)
Brassinolides 0.2 ppm	169.8	174.2	55	57	57	60	84 (66.42)	70 (56.79)
0.5 ppm	166.4	168.6	54	57	57	59	85 (67.22)	73 (58.69)
Sodium 50 μ M	180.7	185.6	52	55	54	56	88 (69.73)	80 (64.43)
nitroprusside 75 μ M	172.5	175.6	53	56	55	57	80 (63.44)	74 (59.34)
Mean	171.828	175.16	53.857	56.428	55.714	58.00	84.714	73.143
SEd	1.566	2.622	0.743	0.617	0.859	0.817	1.036	0.767
CD (P=0.05)	3.291**	5.452**	1.545	1.284	1.788	1.699	2.154	1.595

HSPs in plants, sodium nitroprusside can help them further resist heat stress (Garg *et al.*, 2012). Application of sodium nitroprusside (0.5 mM) increased enzymatic antioxidants and non-enzymatic antioxidants such as peroxidase, superoxide dismutase and catalase enzymes under heat stress (43°C) and reduced electrolyte leakage and lipid

peroxidation in wheat (Karpets *et al.*, 2011; Hasanuzzaman *et al.*, 2012). Mung bean leaves treated with sodium nitroprusside recorded lower membrane leakage, H₂O₂ production and lipid peroxidation (48%) compared to the control (Yang *et al.*, 2006). The same treatment increased the carbonic anhydrase, nitrate reductase and RuBisCo





Fig 3: Effect of foliar spray with phytohormones on proline (mg/g DW) in maize COH (M) 8 under temperature stress.

Table 3: Effect of foliar spray with phytohormones on chlorophyll content (mg/g) in maize COH (M) 8 under temperature stress.

Foliar spray treatments		Ambient condition (34°C)			Open top chamber (42°C)			Ambient condition (34°C)	Open top Chamber (42°C)
		Chlorophyll 'a'	Chlorophyll 'b'	Total chlorophyll	Chlorophyll 'a'	Chlorophyll 'b'	Total chlorophyll	Relative water content	
Control		0.356	0.312	0.738	0.335	0.539	0.869	16	11
								(23.57)	(19.37)
Salicylic acid	50 ppm	0.375	0.441	0.743	0.302	0.601	0.896	18	13
								(25.10)	(21.13)
	75 ppm	0.359	0.514	0.868	0.341	0.595	0.930	18	14
								(25.10)	(21.97)
Brassinolides	0.2 ppm	0.336	0.626	0.955	0.338	0.512	0.845	17	13
								(24.33)	(21.13)
	0.5 ppm	0.350	0.628	0.971	0.350	0.521	0.866	17	13
								(24.35)	(21.13)
Sodium nitroprusside	50 µM	0.353	0.628	0.974	0.351	0.604	0.947	17	14
								(24.35)	(21.97)
	75 µM	0.367	0.636	0.995	0.347	0.597	0.938	17	13
								(24.35)	(21.13)
Mean		0.357	0.541	0.892	0.338	0.567	0.899	17.143	13.00
SEd		0.005	0.009	0.012	0.004	0.007	0.010	0.334	0.163
CD (P=0.05)		0.009	0.019	0.024	0.007	0.016	0.021	0.694	0.339

Table 4: Effect of foliar spray with phytohormones on catalase (μg of $\text{H}_2\text{O}_2/\text{g}/\text{minute}$) and peroxidase (g tissue/min) activity in maize COH (M) 8 under temperature stress.

Foliar spray treatments		Ambient condition		Open top chamber		Ambient condition		Open top chamber	
		(34°C)		(42°C)		(34°C)		(42°C)	
		Catalase				Peroxidase			
		Before spray	After spray	Before spray	After spray	Before spray	After spray	Before spray	After spray
Control		120.3	121.0	115.6	123.2	126.2	127.4	125.4	127.3
Salicylic acid	50 ppm	119.5	123.9	114.4	126.6	128.9	129.3	127.3	135.1
	75 ppm	118.4	125.8	116.5	130.5	127.2	132.2	130.2	142.5
Brassinolides	0.2 ppm	119.3	122.9	115.4	125.9	126.3	127.4	126.4	138.2
	0.5 ppm	120.4	124.4	116.8	127.2	128.0	128.5	128.5	133.2
Sodium nitroprusside	50 µM	121.6	127.3	115.9	132.1	125.1	130.4	129.4	145.4
	75 µM	122.0	124.7	116.9	130.2	125.4	129.6	130.6	140.3
Mean		120.214	124.286	115.928	127.957	126.728	129.257	128.257	137.428
Ed		1.431	1.591	1.831	1.610	1.512	1.702	1.824	1.274
CD (P=0.05)		NS	3.308	NS	3.348	NS	NS	NS	2.650

**Fig 4:** Effect of foliar spray with phytohormones on superoxide dismutase activity ($\text{U mg}^{-1} \text{ protein min}^{-1}$) in maize COH (M) 8 under temperature stress.

seedlings showed increased transcription of enzymes such as small heat shock protein 26, sucrose phosphate synthase and delta-1-pyrroline-5-carboxylate synthase (Uchida *et al.*, 2002). Sodium nitroprusside increased the H_2S accumulation and activities of L-cysteine disulfhydryase, resulting in higher survival rate of maize seedlings under heat stress (Li *et al.*, 2013). Exogenous application of salicylic acid reduces the negative effects of abiotic stress in crops and increased plant resistance to heat stress (Hu *et al.*, 2010; Khan *et al.*, 2013). Treating *Triticum aestivum* with 0.5 mM salicylic acid can reduce heat impact by limiting ethylene formation (Khan *et al.*, 2013). Salicylic acid can regulate plant mechanisms in as well as osmolytes like glycine betaine and proline in tomato (Siddiqui *et al.*,

2017b). Under heat stress, paddy both optimum and stress conditions *via* cross-talk signalling with other phytohormones (Horvath *et al.*, 2007).

CONCLUSION

This study clearly demonstrates that heat stress affects maize plants growth, and biochemical parameters. It can be overcome by spraying with sodium nitroprusside 50 μM and salicylic acid 75 ppm. It is a simple and easy method to follow by farmers to mitigate the heat stress. We suggest that the farmers of maize could choose nitroprusside 50 μM and salicylic acid 75 ppm for large-scale commercial applications to increase growth, yield and antioxidant activity of maize plant under heat stress.

Conflict of interest

All authors declare that they have no conflicts of interest.

REFERENCES

- Aftab, T., Masroor, M., Khan, A., Idrees, M. and Naeem, M. (2010). Salicylic acid acts as potent enhancer of growth, photosynthesis and artemisinin production in *Artemisia annua* L. *Journal of crop science and biotechnology*. **13(3)**: 183-188.
- Aghdam, M.S., Jannatizadeh, A., Sheikh-Assadi, M. and Malekzadeh, P. (2016). Alleviation of postharvest chilling injury in anthurium cut flowers by salicylic acid treatment. *Scientia Horticulturae*. **202**: 70-76.
- Bazzaz, M.M., Hossain, A., Farooq, M., Alharby, H., Bamagoos, A. and Nuruzzaman, M. (2020). Phenology, growth and yield are strongly influenced by heat stress in late sown mustard (*Brassica spp*) varieties. *Pakistan Journal of Botany*. **52**: 1189-95.
- Chaturvedi, A.K., Bahuguna, R.N., Pal, M., Shah, D., Maurya, S. and Jagadish, K.S. (2017). Elevated CO₂ and heat stress interactions affect grain yield, quality and mineral nutrient composition in rice under field conditions. *Field Crops Research*. **206**: 149-157.
- Choudhary, A., Kumar, A. and Kaur, N. (2020). ROS and oxidative burst: Roots in plant development. *Plant Divers*. **42**: 33-43.
- Deng, Z. and Song, S. (2012). Sodium nitroprusside, ferricyanide, nitrite and nitrate decrease the thermo-dormancy of lettuce seed germination in a nitric oxide-dependent manner in light. *South African Journal of Botany*. **78**: 139-146.
- El-Esawi, M.A., Elansary, H.O., El-Shanhorey, N.A., Abdel-Hamid, A.M., Ali, H.M. and Elshikh, M.S. (2017). Salicylic acid-regulated antioxidant mechanisms and gene expression enhance rosemary performance under saline conditions. *Frontiers in Physiology*. **8**: 716.
- Esim, N. and Atici, O. (2014). Nitric oxide improves chilling tolerance of maize by affecting apoplastic antioxidative enzymes in leaves. *Plant Growth Regulation*. **72(1)**: 29-38.
- Garg, D., Sareen, S., Dalal, S., Tiwari, R. and Singh, R. (2012). Heat shock protein based SNP marker for terminal heat stress in wheat (*Triticum aestivum* L.). *Australian Journal of Crop Science*. **6**: 1516-21.
- Ghani, A., Khan, I., Ahmed, I. and Mustafa, I. (2015). Abd-Ur-Rehman. Amelioration of lead toxicity in *pisum sativum* (L.) by foliar application of salicylic acid. *Journal of Environmental and Analytical Toxicology*. **5(4)**: 292.
- Hajhashemi, S., Skalicky, M., Brestic, M. and Pavla, V. (2021). Effect of sodium nitroprusside on physiological and anatomical features of salt-stressed *Raphanus sativus*. *Plant Physiology and Biochemistry*. **169**: 160-170.
- Haldimann, P. and Feller, U. (2004). Inhibition of photosynthesis by high temperature in oak (*Quercus pubescens* L.) leaves grown under natural conditions closely correlates with a reversible heat dependent reduction of the activation state of ribulose 1, 5 biphosphate carboxylase/oxygenase. *Plant, Cell and Environment*. **27(9)**: 1169-1183.
- Hameed, S. and Ali, M.K. (2016). Exogenous application of salicylic acid: Inducing thermotolerance in cotton (*Gossypium hirsutum* L.) seedlings. *International Journal of Agricultural and Food Research*. **5(4)**: 9-18.
- Hasanuzzaman, M., Nahar, K., Alam, M.M. and Fujita, M. (2012). Exogenous nitric oxide alleviates high temperature induced oxidative stress in wheat (*Triticum aestivum* L.) seedlings by modulating the antioxidant defense and glyoxalase system. *Australian Journal of Crop Science*. **6(8)**: 1314-1323.
- Hayat, S., Alyemeni, M.N. and Hasan, S.A. (2012). Foliar spray of brassinosteroid enhances yield and quality of *Solanum lycopersicum* under cadmium stress. *Saudi Journal of Biological Science*. **19(3)**: 325-335.
- Horvath, E., Szalai, G. and Janda, T. (2007). Induction of abiotic stress tolerance by salicylic acid signaling. *Journal of Plant Growth Regulation*. **26(3)**: 290-300.
- Hu, X., Li, Y., Li, C., Yang, H., Wang, W. and Lu, M. (2010). Characterization of small heat shock proteins associated with maize tolerance to combined drought and heat stress. *Journal of Plant Growth Regulation*. **29(4)**: 455-464.
- Iqbal, N., Sehar, Z., Fatma, M., Umar, S., Sofo, A. and Khan, N.A. (2022). Nitric oxide and abscisic acid mediate heat stress tolerance through regulation of osmolytes and antioxidants to protect photosynthesis and growth in wheat plants. *Antioxidants*. **11**: 372.
- Karpets, Y.V., Kolupaev, Y.E. and Yastreb, T. (2011). Effect of sodium nitroprusside on heat resistance of wheat coleoptiles: Dependence on the formation and scavenging of reactive oxygen species. *Russian Journal of Plant Physiology*. **58(6)**: 1027-1033.
- Khan, M.I.R., Iqbal, N., Masood, A., Per, T.S. and Khan, N.A. (2013). Salicylic acid alleviates adverse effects of heat stress on photosynthesis through changes in proline production and ethylene formation. *Plant Signaling and Behaviour*. **8(11)**: e26374.
- Kharbech, O., Sakouhi, L., Massoud, M.B., Mur, L.A.J., Corpas, F.J., Djebali, W. (2020). Nitric oxide and hydrogen sulfide protect plasma membrane integrity and mitigate chromium-induced methylglyoxal toxicity in maize seedlings. *Plant Physiology and Biochemistry*. **157**: 244-255.
- Kuchlan, P. and Kuchlan, M.K. (2023). Effect of salicylic acid on plant physiological and yield traits of soybean. *Legume Research*. **46(1)**: 56-61. doi: 10.18805/LR-4527.
- Li, Z.G., Yang, S.Z., Long, W.B., Yang, G.X. and Shen, Z.Z. (2013). Hydrogen sulphide may be a novel downstream signal molecule in nitric oxide induced heat tolerance of maize (*Zea mays* L.) seedlings. *Plant, Cell and Environment*. **36(8)**: 1564-1572.
- Lobell, D.B., Schlenker, W. and Costa-Roberts, J. (2011). Climate trends and global crop production since 1980. *Science*. **333(6042)**: 616-620.
- Mittler, R. (2002). Oxidative stress, antioxidants and stress tolerance. *Trends in Plant Science*. **7(9)**: 405-410.
- Muthulakshmi, S., Santhi, M. and Lingakumar, K. (2017). Effect of sodium nitroprusside (SNP) on physiological and biochemical responses of *Sorghum vulgare* press under salt stress. *International Journal of Applied Research*. **3**: 33-37.

- Panase, V. and Sukhatme, P. (1957). Genetics of quantitative characters in relation to plant breeding. *Indian Journal of Genetics and Plant Breeding*. **17(2)**: 318-328.
- Qiao, W. and Fan, L.M. (2008). Nitric oxide signalling in plant responses to abiotic stresses. *Journal of Integrative Plant Biology*. **50**: 1238-1246.
- Ray, P.D., Huang, B.W. and Tsuji, Y. (2012). Reactive oxygen species (ROS) homeostasis and redox regulation in cellular signaling. *Cellular Signalling*. **24(5)**: 981-990.
- Rennenberg, H., Loreto, F., Polle, A., Brilli, F., Fares, S., Beniwal, R. and Gessler, A. (2006). Physiological responses of forest trees to heat and drought. *Plant Biology*. **8(05)**: 556-571.
- Sattar, A., Sher, A., Ijaz, M., Ullah, M.S., Ahmad, N. and Umar, U.U. (2020). Individual and combined effect of terminal drought and heat stress on allometric growth, grain yield and quality of bread wheat. *Pakistan Journal of Botany*. **52**: 405-412.
- Shakirova, F.M., Sakhabutdinova, A.R., Bezrukova, M.V., Fatkhutdinova, R.A. and Fatkhutdinova, D.R. (2003). Changes in the hormonal status of wheat seedlings induced by salicylic acid and salinity. *Plant Science*. **164(3)**: 317-322.
- Sharma, S. and Manjeet. (2020). Heat stress effects in fruit crops: A Review. *Agricultural Reviews*. **41(1)**: 73-78. doi: 10.18805/ag.R-1951.
- Sharma, A., Kumar, V., Shahzad, B., Ramakrishnan, M., Singh Sidhu, G.P., Bali, A.S., Handa, N., Kapoor, D., Yadav, P. and Khanna, K. (2020). Photosynthetic response of plants under different abiotic stresses: A review. *Journal of Plant Growth Regulation*. **39(2)**: 509-531.
- Shiferaw, B., Prasanna, B.M., Hellin, J. and Banziger, M. (2011). Crops that feed the world 6. Past successes and future challenges to the role played by maize in global food security. *Food Security*. **3(3)**: 307-327.
- Siddiqui, M.H., Alamri, S.A., Al-Khaishany M.Y.Y., Al-Qutami M.A., Ali H.M. and Khan, M.N. (2017a). Sodium nitroprusside and indole acetic acid improve the tolerance of tomato plants to heat stress by protecting against DNA damage. *Journal of Plant Interaction*. **12**: 177-186.
- Siddiqui, M., Alamri, S.A., Mutahhar, Y., Al-Khaishany, M., Al-Qutaytami, H. and Nasir Khan, M. (2017b). Nitric oxide and calcium induced physiobiochemical changes in tomato (*Solanum lycopersicum*) plant under heat stress. *Fresenius Environmental Bulletin*. **26(2a)**: 1663-1672.
- Sita, K., Sehgal, A., Bhardwaj, A., Bhandari, K., Kumar, S., Prasad, P.V., Jha, U., Siddique, K.H. and Nayyar, H. (2021). Nitric oxide secures reproductive efficiency in heat-stressed lentil (*Lens culinaris* Medik.) plants by enhancing the photosynthetic ability to improve yield traits. *Physiology and Molecular Biology of Plants*. **27(11)**: 2549-2566.
- Suwa, R., Hakata, H., Hara, H., El-Shemy, H.A., Adu-Gyamfi, J.J., Nguyen, N.T., Kanai, S., Lightfoot, D.A., Mohapatra, P.K. and Fujita, K. (2010). High temperature effects on photosynthate partitioning and sugar metabolism during ear expansion in maize (*Zea mays* L.) genotypes. *Plant Physiology and Biochemistry*. **48(2-3)**: 124-130.
- Suzuki, N., Miller, G., Morales, J., Shulaev, V., Torres, M.A. and Mittler, R. (2011). Respiratory burst oxidases: The engines of ROS signaling. *Current Opinion Plant Biology*. **14**: 691-699.
- Tian, X., He, M., Wang, Z., Zhang, J., Song, Y., He, Z. and Dong, Y. (2015). Application of nitric oxide and calcium nitrate enhances tolerance of wheat seedlings to salt stress. *Plant Growth Regulation*. **77(3)**: 343-356.
- Tiwari, Y.K. and Yadav, S.K. (2019). High temperature stress tolerance in maize (*Zea mays* L.): Physiological and molecular mechanisms. *Journal of Plant and Biology*. **62**: 93-102.
- Trost, G. (2014). Poly (A) Polymerase 1 (PAPS1) influences organ size and pathogen response in *Arabidopsis thaliana*. Universitat Potsdam.
- Tufail, A., Arfan, M., Gurmani, A.R., Khan, A. and Bano, A. (2013). Salicylic acid induced salinity tolerance in maize (*Zea mays*). *Pakistan Journal of Botany*. **45(S1)**: 75-82.
- Uchida, A., Jagendorf, A.T., Hibino, T., Takabe, T. and Takabe, T. (2002). Effects of hydrogen peroxide and nitric oxide on both salt and heat stress tolerance in rice. *Plant Science*. **163(3)**: 515-523.
- Yang, F., Jorgensen, A.D., Li, H., Søndergaard, I., Finnie, C., Svensson, B., Jiang, D., Wollenweber, B. and Jacobsen, S. (2011). Implications of high temperature events and water deficits on protein profiles in wheat (*Triticum aestivum* L. cv. Vinjett) grain. *Proteomics*. **11(9)**: 1684-1695.
- Yang, J.D., Yun J.Y., Zhang, T.H. and Zhao, H.L. (2006). Presoaking with nitric oxide donor SNP alleviates heat shock damages in mung bean leaf discs. *Botanical Studies*. **47**: 129-136.
- Youldash, K.M., Barutcular, C., El Sabagh, A., Toptas, I., Kayaalp, G.T. and Hossain, A. (2020). Evaluation of grain yield in fifty-eight spring bread wheat genotypes grown under heat stress. *Pakistan Journal of Botany*. **52**: 33-42.
- Yu, L., Gao, R., Shi, Q., Wang, X., Wei, M. and Yang, F. (2013). Exogenous application of sodium nitroprusside alleviated cadmium induced chlorosis, photosynthesis inhibition and oxidative stress in cucumber. *Pakistan Journal of Botany*. **45**: 813-819.
- Zhao, W., Wei, X.H. and Dong, S.K. (2024). Effects of irrigation with sodium nitroprusside on physiological characteristics of soybean under drought stress. *Legume Research*. **47(5)**: 817-822. doi: 10.18805/LRF-784.