



# Impact of Seasonal Weather Variations on Physical Attributes of Robusta Coffee Beans

E. Maneerat<sup>1</sup>, C. Saensano<sup>1</sup>, R. Chiarawipa<sup>1</sup>, V. Wongvarodom<sup>1</sup>  
P. Teerawattanapong<sup>2</sup>, M. Rueangkhanab<sup>3</sup>

10.18805/IJARE.AF-865

## ABSTRACT

**Background:** Robusta coffee is a type of cultivated agricultural commodity. It is grown to produce high-quality coffee. The weather conditions have a significant impact on the characteristics of green coffee beans, such as their aroma and taste. As a result, fluctuations in weather conditions can affect the quality of the final product.

**Methods:** This study assessed Robusta coffee beans' physical properties and defects in southern Thailand. The characteristics of green coffee beans and the weather variations of the southern peninsula were observed.

**Result:** It was found that significantly different proportions of normal, pea berry and defective coffee beans were found across geographical regions. In areas with high rainy days and total rainfall, full black disorder had a higher incidence (16.46%) of defective coffee beans. Additionally, the geometric mean diameter (GMD) and surface area of green coffee beans had a significant positive correlation with minimum temperature (Tmin) after the 6-to 9-month period of fruit setting. This study indicates that recent weather variations have had a significant impact on the characteristics and quality of Robusta green beans. The improvement of green coffee bean practices to achieve high-quality Robusta beans in response to pre-harvest weather conditions should be further investigated.

**Key words:** Coffee agroecology, Defect, Fine Robusta, Green bean, Weather variation.

## INTRODUCTION

The quality of coffee beans is influenced by various factors, including geographic location, cultivation conditions and bean characteristics (Worku *et al.*, 2018). Geographic location significantly impacts coffee bean quality, with temperature and sunlight affecting flavor and aroma (Tolessa *et al.*, 2017). Robusta coffee is a significant agricultural commodity in multiple regions across the globe and is extensively cultivated in tropical areas (Ababu and Getahun, 2021). In addition, Robusta coffee is a botanical entity that thrives exceptionally well in the southern regions of Thailand, constituting approximately 51% of the nation's entire coffee output (Office of Agricultural Economics, 2022). The quality of Robusta coffee is influenced by various environmental factors, including temperature, humidity and rainfall, thereby impacting the occurrence of defects in unripen coffee beans (Ahmed *et al.*, 2021). Moreover, the defect identified in the beans has a discernible adverse effect on the aroma and taste of the coffee (Specialty Coffee Association, 2018).

Farmers in the southern region of Thailand engage in the cultivation of Robusta coffee, which is highly favored in areas with mixed orchards and rubber plantations (Chiarawipa and Sirikantayakul, 2015). The growing conditions for coffee beans can range from open to shaded environments, with varying intensities (Chiarawipa *et al.*, 2021). Although Robusta coffee can grow in a wide range of climates, the coffee plants might produce smaller beans with lower-quality seeds when inadequate water is applied during cherry development (DaMatta *et al.*, 2018).

<sup>1</sup>Agricultural Innovation and Management Division, Faculty of Natural Resources, Prince of Songkla University, Songkhla 90110, Thailand.

<sup>2</sup>Yala Horticultural Research Center, Horticulture Research Institute, Department of Agriculture, Yala 95150, Thailand.

<sup>3</sup>Office of Agricultural Research and Development Region 8, Department of Agriculture, Songkhla 90110, Thailand.

**Corresponding Author:** R. Chiarawipa, Agricultural Innovation and Management Division, Faculty of Natural Resources, Prince of Songkla University, Songkhla 90110, Thailand.  
Email: rawee.c@psu.ac.th

**How to cite this article:** Maneerat, E., Saensano, C., Chiarawipa, R., Wongvarodom, V., Teerawattanapong, P. and Rueangkhanab, M. (2024). Impact of Seasonal Weather Variations on Physical Attributes of Robusta Coffee Beans. Indian Journal of Agricultural Research. 58(6): 1259-1265. doi: 10.18805/IJARE.AF-865.

**Submitted:** 29-02-2024 **Accepted:** 02-07-2024 **Online:** 08-08-2024

Furthermore, pre-harvest weather conditions significantly impact the physical characteristics of coffee beans (Bunya-atichart *et al.*, 2018). Both altitude and climatic variations have a direct impact on the quality of coffee beans produced (Cassamo *et al.*, 2022). The physical attributes of coffee beans exhibit variability based on the environmental factors in the planting area, including topography, climate, soil fertility (Wintgens, 2012; Hareesh and Jayarama, 2024) and pest population in the coffee plantation (Rimbing *et al.*, 2023).

To ensure the quality of the coffee beans, it is important to consider the suitability of the planting site's conditions

during the pre-harvest stage, as well as the management of the harvesting and post-harvest periods. The main aim of this particular study was to assess the physical characteristics and defect properties of Robusta coffee beans in the Southern Thailand region, while also evaluating the variations in growing seasonal weather.

## MATERIALS AND METHODS

The study was conducted at four major Robusta coffee plantations in four provinces, in Southern Thailand: Ranong-Kra Buri (RN-KR) (altitude 82 m), Chumphon-Tha Sae (CP-TS) (altitude 483 m), Surat Thani-Phanom (SR-PN) (altitude 143 m) and Krabi-Plai Phraya (KB-PY) (altitude 135 m). In these regions, coffee plantations have been established on a large scale. The arrangements varied according to the geographical locations, which are rainfed orchard systems. The dry process of Robusta coffee beans was used to retain the natural flavors. After the collection of the cherries, they were subjected to solar drying. To conduct the dry process processing, a set of coffee cherries was selected at random and positioned within a pristine and sunny area for 3-4 weeks until reaching a humidity level of 9-12%. When the coffee cherries were dry, they could then be put into a huller machine to remove the outer husk. The beans were also laid out in the sun for a few days to ensure that the moisture content was reduced. To evaluate the characteristics of beans, it was then followed by sorting the beans according to weight, color and physical properties.

The categorization of coffee beans comprises three distinct types: regular beans, pea berry and defect. For analysis, three separate samples, each weighing 350 g, were assessed at various locations using an analytical balance (model ES-1200HA, Zepper scales LTD, China). Subsequently, the proportions of each coffee bean type were recorded.

After that, the defective beans were classified according to their disorders as described by the Uganda Coffee Development Authority (UCDA) standards. It has also implemented a method for sorting and verifying the integrity of the beans to produce quality coffee beans (Coffee Quality Institute, 2018).

All coffee beans were classified according to the severity that affects coffee quality into 6 primary defects: Full black, full sour, cherry/pod, fungus damage, foreign matter and severe insect damage and 11 secondary defects: Partial black, partial sour, parchment, floater, immature/unripe, withered, shell, broken/chipped/cut, hull/husk, slight insect damage and chalky (Coffee Quality Institute, 2018) to compare the degree of abnormality of coffee beans at each location. Each defective bean of Robusta coffee was then weighed to calculate the respective percentages.

The physical properties of Robusta coffee beans were assessed at each location by randomly selecting 1 kg per location and conducting three replicates of 100 seeds. To determine the width (a), length (b) and thickness (c) of the coffee beans, a digital vernier caliper was used.

The equations used to calculate the physical characteristics were adapted from Mohsenin method (Mohsenin, 1971; Kattalee and Inprasit, 2012), specifically the following equations:

$$\text{Geometric mean diameter (GMD) (cm)} = (a*b*c)^{1/3} \quad \dots(1)$$

$$\text{Sphericity (\%)} = \frac{\text{GMD}}{b} \quad \dots(2)$$

$$\text{Volume (cm}^3\text{)} = \frac{[(\pi*a*c)*b^2]}{[6*(2b-(a*c)^{1/2})]} \quad \dots(3)$$

$$\text{Surface area (cm}^2\text{)} = \pi*(\text{GMD})^2 \quad \dots(4)$$

Bulk density ( $\rho$ ) (g/cm<sup>3</sup>); the volume of the cylinder (volume; v) is then filled with coffee beans and weighing the mass of coffee beans (mass; m)

$$\frac{m}{v} \quad \dots(5)$$

To analyze the color of Robusta coffee beans, 30 beans from 350 g were randomly sampled per area and assessed using a colorimeter (CR-400 Chroma meter, Minolta, Japan). The CIELAB system (L\*, a\* and b\* values) was recorded and compared to determine the average color values.

Monthly datasets of air temperature, including maximum temperature (Tmax) and minimum temperature (Tmin), total rainfall and relative humidity (RH) were observed during 2019 to 2020. The meteorological data were collected from agricultural meteorological stations located in four provinces. The relationship between monthly weather variables (at the 3<sup>rd</sup>, 6<sup>th</sup>, 9<sup>th</sup> and 12<sup>th</sup> months after fruit setting periods) and physical properties of green beans from those stations were analyzed using linear and non-linear regressions. Furthermore, the average altitude above sea level was recorded, encompassing the diverse locations of the coffee orchards.

To analyze the variance of the data, a completely randomized design was used and ANOVA (Analysis of Variance) was conducted. The mean was compared using the Duncan's new multiple range test (DMRT) method at a 95% confidence level, which was carried out using the R program. Regression analysis was utilized to assess the geographical and weather variables that exert the most significant influence on the physical properties and size of coffee beans ( $r^2 \geq 0.65$ ).

## RESULT AND DISCUSSION

### Defect classification of robusta coffee beans

In KB-PY, normal beans were the most prevalent, accounting for 74.71% of the samples, showing a statistically significant difference compared to coffee beans from RN-KR, where normal beans were least common at 58.25%. On the other hand, CP-TS had the highest percentage of pea berry beans at 18.62%, which was similar to RN-KR's 17.44%, but significantly different from SR-PN (16.15%) and

KB-PY (13.44%). Notably, RN-KR (24.31%) had the highest occurrence of total defects, which differed significantly from the other locations. The lowest total defect was found in KB-PY (11.85%) (Table 1).

In terms of major defects (category 1) and minor defects (category 2) observed in Robusta coffee beans, RN-KR exhibited the highest occurrence of primary defects at 16.82%, while KB-PY had the lowest percentage of secondary and minor defects at 7.80% and 4.60%, respectively. When considering the total of defects, RN-KR had the highest percentage of total defects at 24.63%, whereas KB-PY had the lowest percentage of total defects at 12.40%. The primary defect that stood out the most was full black in RN-KR, accounting for 16.46%. As for minor defects, slight insect damage was the most prominent

across all locations, ranging from 2.10% to 4.38%. However, no instances of parchment, floater, or shell disorders were observed in any of the locations (Table 2).

The full black defect in coffee beans is a type of category 2 defect according to the Specialty Coffee Association (SCA) standards (Specialty Coffee Association, 2018). This defect is characterized by blackened, darkened, or burnt beans that can occur as a result of over fermentation or over drying of the beans during processing. Additionally, this defect can be attributed to the occurrence of fungal infections or damage caused by insects on the beans. To identify full black beans, one can observe their dark and lustrous appearance, which can adversely affect the coffee's flavor profile by imparting a burnt or bitter taste (Gonzalez-Sanchez *et al.*, 2024). Meanwhile, the most common minor defects in green beans are slight insect damage, broken or chipped flesh and cuts that damage the seeds. These defects can occur during the post-harvest process, such as during harvesting, transportation, or storage. Farmers should sort the coffee beans to identify and remove these defects, as they can affect the quality, including its flavor and aroma.

Approximately 10% to 30% of coffee defects emerge throughout harvesting. Additionally, coffee production is subject to the influence of climate change, whereby variations in rainfall patterns and occurrences of extreme weather events directly affect the yield and quality of coffee (Legesse, 2022; Rimbing *et al.*, 2023). Similarly, this study

**Table 1:** Classification of percentage of normal bean, pea berry and defect.

| Coffee beans | Normal bean (%) | Pea berry (%) | Defect (%)  |
|--------------|-----------------|---------------|-------------|
| RN-KR        | 58.25±1.84c     | 17.44±1.17b   | 24.31±2.20a |
| CP-TS        | 65.56±4.05b     | 18.62±1.29a   | 15.82±3.53c |
| SR-PN        | 64.97±1.32b     | 16.15±1.33c   | 18.88±1.94b |
| KB-PY        | 74.71±2.00a     | 13.44±0.98d   | 11.85±2.05d |

Mean (±S.D.) in the same column with different letters are significantly different at  $P \leq 0.05$  by DMRT.

**Table 2:** Defects of robusta coffee beans in southern Thailand as UCDA standard.

| Defect category      | Green beans |             |             |            |
|----------------------|-------------|-------------|-------------|------------|
|                      | RN-KR       | CP-TS       | SR-PN       | KB-PY      |
| <b>Category 1:</b>   |             |             |             |            |
| Full black           | 16.46±2.05a | 9.23±1.80bc | 10.86±2.12b | 7.74±1.32c |
| Full sour            | 0.24±0.14   | 0.00        | 0.16±0.13   | 0.00       |
| Cherry/pod           | 0.13±0.07b  | 0.09±0.04b  | 0.51±0.19a  | 0.00       |
| Fungus damage        | 0.13±0.11   | 0.00        | 0.00        | 0.00       |
| Foreign matter       | 0.09±0.03   | 0.09±0.05   | 0.10±0.05   | 0.08±0.08  |
| Severe insect damage | 0.19b       | 0.30±0.14b  | 0.75±0.23a  | 0.11±0.05b |
| Total category 1 (%) | 16.82       | 9.62        | 12.19       | 7.80       |
| <b>Category 2:</b>   |             |             |             |            |
| Partial black        | 0.99±0.53b  | 1.08±0.57ab | 1.48±0.47a  | 0.98±0.41b |
| Partial sour         | 0.00        | 0.00        | 0.08±0.09   | 0.05       |
| Parchment            | 0.00        | 0.00        | 0.00        | 0.00       |
| Floater              | 0.00        | 0.00        | 0.00        | 0.00       |
| Immature/unripe      | 0.00        | 0.00        | 0.16±0.09b  | 0.55a      |
| Withered             | 0.50±0.04   | 0.00        | 0.20±0.08   | 0.36±0.39  |
| Shell                | 0.00        | 0.00        | 0.00        | 0.00       |
| Broken/chipped/cut   | 4.64±0.67a  | 3.25±0.39b  | 1.27±0.35c  | 0.42±0.16d |
| Hull/husk            | 0.08±0.03   | 0.33±0.09   | 0.10±0.05   | 0.02±0.02  |
| Slight Insect damage | 2.10±1.78d  | 3.63±0.59b  | 4.38±1.07a  | 2.80±0.05c |
| Chalky               | 0.00        | 0.08±0.60   | 0.00        | 0.00       |
| Total category 2 (%) | 7.81        | 7.94        | 7.26        | 4.60       |
| Total defect (%)     | 24.63       | 17.56       | 19.45       | 12.40      |

Means±S.D. in the same column with different letters are significantly different at  $P \leq 0.05$  by DMRT.

found that the impacts of elevated total rainfall levels resulted in a rise in the prevalence of full black beans in the RN-KR location. Moreover, the study revealed that rainfall variation was associated with discolored beans having a higher blue-green color (Table 3). A large number of the main defects found, in full black, were caused by fermentation in processing or microbial infection from dropped or overripe seeds. This is most prominent in the RN-KR samples. To avoid these defects, farmers should harvest the cherries at the appropriate maturity and follow the climatic conditions (Worku *et al.*, 2018). Although farmers cannot avoid harvesting the coffee cherries during periods of heavy rainfall, it is imperative that they subsequently desiccate the cherries in an area that is sheltered and facilitates air circulation.


This study has provided evidence to suggest that climatic conditions substantially influence the imperfection

of Robusta green coffee beans (Bunn *et al.*, 2015). It has been ascertained that an excess of rainfall is correlated with a heightened manifestation of defect attributes in the province of RN. To mitigate such losses, agriculturalists ought to implement measures to prevent the occurrence of

**Table 3:** Color shades of Robusta coffee beans using the CIELAB system ( $L^*$ ,  $a^*$ ,  $b^*$ ).

| Locations | Color parameters |            |             |
|-----------|------------------|------------|-------------|
|           | $L^*$            | $a^*$      | $b^*$       |
| RN-KR     | 53.78±5.68b      | 3.86±2.02d | 20.31±5.41b |
| CP-TS     | 54.75±5.63b      | 6.23±2.84b | 22.16±3.10a |
| SR-PN     | 50.88±6.05c      | 7.42±2.05a | 22.90±3.55a |
| KB-PY     | 58.92±5.18a      | 5.03±2.08c | 22.08±2.23a |

Means±S.D. in the same column with different letters are significantly different at  $P \leq 0.05$  by DMRT.



**Fig 1:** Variations in the rainy days.

minor imperfections. These measures encompass sound farming techniques, such as implementing appropriate harvesting and storage practices.

#### Weather variations on characteristics of coffee bean

During the growing season (at the 3<sup>rd</sup>, 6<sup>th</sup>, 9<sup>th</sup> and 12<sup>th</sup> month after fruit setting periods), plentiful rainfall was found in RN province (1,649.46 to 2,750.82 mm), although rainy days are high during 6<sup>th</sup> and 9<sup>th</sup> month (22 to 30 days) (Fig 1). The average Tmin was highest in CP province (23.20 to 24.87°C) after decreasing from the 6<sup>th</sup> to 12<sup>th</sup> month. Meanwhile, Tmax was similar highest in SR and KB provinces (30.37 to 36.10°C) during the growing season. All locations, RH was slightly increased during the 9<sup>th</sup> to 12<sup>th</sup> month, with an average RH of 77.33 to 84.93%, 80.87 to 88.33% and 80.47 to 86.00%, respectively.

KB-PY coffee beans exhibited the highest L\* with a value of 58.92, which was statistically distinct from the other coffee beans. In terms of a\* testing, SR-PN displayed the

highest value at 7.42, which was significantly different from RN-KR coffee beans that had the lowest value at 3.86. Regarding the b\* value, coffee beans from RN-KR recorded the lowest value at 20.31, while coffee beans from other locations fell within a similar range of 22.08-22.90 (Table 3).

When considering the physical properties (geometric mean diameter, sphericity, volume and surface area) of coffee bean, CP-TS exhibited the highest values among all parameters (0.628 cm, 75.317%, 0.266 cm<sup>3</sup> and 1.247 cm<sup>2</sup>, respectively), whereas KB-PY showed the lowest values (0.591 cm, 72.543%, 0.177 cm<sup>3</sup> and 1.107 cm<sup>2</sup>, respectively). In terms of bulk density, SR-PN had the highest value at 3.090 g/cm<sup>3</sup>, which was statistically significantly different from RN-KR coffee beans (2.963 g/cm<sup>3</sup>) (Table 4).

Table 5 shows the overall regression analysis. The results showed that total rainfall, RH, Tmax and Tmin during the 9-month period after fruit setting had a positive effect on physical properties. Importantly, the relationship

**Table 4:** Physical properties of Robusta green coffee beans in Southern Thailand.

| Coffee beans | Physical properties          |                |                           |                                 |                                   |
|--------------|------------------------------|----------------|---------------------------|---------------------------------|-----------------------------------|
|              | Geometric mean diameter (cm) | Sphericity (%) | Volume (cm <sup>3</sup> ) | Surface area (cm <sup>2</sup> ) | Bulk density (g/cm <sup>3</sup> ) |
| RN-KR        | 0.621±0.057ab                | 73.514±5.235b  | 0.204±0.110b              | 1.222±0.215ab                   | 2.963±0.001d                      |
| CP-TS        | 0.628±0.048a                 | 75.317±5.909a  | 0.266±0.224a              | 1.247±0.198a                    | 2.995±0.002c                      |
| SR-PN        | 0.060±0.070bc                | 73.681±5.016b  | 0.208±0.095b              | 1.169±0.269bc                   | 3.090±0.003a                      |
| KB-PY        | 0.591±4.197c                 | 72.543±4.197b  | 0.177±0.064b              | 1.107±0.209c                    | 3.030±0.004b                      |

Mean±S.D. in the same column with different letters are significantly different at P≤0.05 by DMRT.

**Table 5:** The relationship between weather variations (9<sup>th</sup> month after fruit setting) and physical properties of Robusta green coffee beans.

| Weather  | Physical properties | Equations            | Coefficient of determination (r <sup>2</sup> ) |
|----------|---------------------|----------------------|------------------------------------------------|
| Rainfall | GMD                 | y= 0.00005x + 0.6072 | 0.059                                          |
|          | Sphericity          | y= -0.0009x + 74.106 | 0.081                                          |
|          | Volume              | y= 0.005x + 0.2257   | 0.090                                          |
|          | Surface area        | y= 0.0005x + 1.1708  | 0.056                                          |
|          | Bulk density        | y= -0.0001x + 3.0615 | 0.554                                          |
| RH       | GMD                 | y= -0.0023x + 0.811  | 0.273                                          |
|          | Sphericity          | y= -0.2695x + 96.733 | 0.759                                          |
|          | Volume              | y= -0.0087x + 0.9574 | 0.749                                          |
|          | Surface area        | y= -0.0089x + 1.9441 | 0.281                                          |
|          | Bulk density        | y= -0.004x + 3.3577  | 0.074                                          |
| Tmax     | GMD                 | y= -0.0004x + 0.6238 | 0.001                                          |
|          | Sphericity          | y= 0.3703x + 62.171  | 0.154                                          |
|          | Volume              | y= 0.0114x - 0.1425  | 0.137                                          |
|          | Surface area        | y= -0.0004x + 1.1978 | 0.001                                          |
|          | Bulk density        | y= 0.0360x + 1.8918  | 0.656                                          |
| Tmin     | GMD                 | y= 0.0313x - 0.1366  | 0.954*                                         |
|          | Sphericity          | y= 1.9731x + 26.639  | 0.794                                          |
|          | Volume              | y= 0.0642x - 1.3188  | 0.791                                          |
|          | Surface area        | y= 0.1163x - 1.5912  | 0.938*                                         |
|          | Bulk density        | y= -0.065x + 4.5709  | 0.388                                          |

\* =Significantly different at P≤0.05.



between Tmin with GMD ( $r^2 = 0.954$ ) and Tmin with surface area ( $r^2 = 0.938$ ) was found to be statistically significant ( $p \leq 0.05$ ).

Studies have also shown that environmental conditions can affect the characteristics of Robusta green beans. The environmental factor of rainfall can affect the bulk density of coffee beans, as excessive rainfall during the ripening stage can lead to increased moisture content and lower bulk density (Kath *et al.*, 2021). This can result in lower quality coffee beans and potential difficulties in the roasting process. The air temperature is an important factor that affects the physical properties of coffee beans. An increase in air temperature can result in a decrease in coffee bean volume, diameter, surface area and sphericity. This reduction in physical properties can have negative impacts on coffee quality and processing (Halagarda and Obrok, 2023). Moreover, high temperatures can cause defects such as discoloration, surface blemishes and cracked beans, reducing the quality of the beans (Kath *et al.*, 2021; Gonzalez-Sanchez *et al.*, 2024).

The physical properties of the coffee beans can also affect the quality of the coffee. This was related to the geometric mean diameter (GMD) and volume of the beans. The annual total rainfall in the KB-PY region is only 1,474 mm, which is significantly lower than the annual total rainfall in the coffee-growing regions of RN-KR and CP-TS, which receive an average of 3,828.7-1,764.5 mm of rain per year (Fig 1). The difference in rainfall is likely a contributing factor to the size and quality of the coffee beans in the KB-PY region (Table 4). Excessive rainfall can lead to a decrease in the body and sweetness of a roasted bean. Additionally, orchard management practices, such as pruning, fertilization (Leite *et al.*, 2015; Hareesh and Jayarama, 2024) and irrigation (Lina *et al.*, 2022), have been found to affect the flavor and yield of unprocessed coffee beans, resulting in increased concentrations of aromatic compounds, higher levels of caffeine and improved yields (Hagos *et al.*, 2018; Hameed *et al.*, 2018).

## CONCLUSION

The physical characteristics and defect prevalence of coffee beans demonstrably vary according to the elevation of the growing region and the prevailing weather conditions during the cherry development, encompassing factors such as rainfall, temperature and relative humidity. These environmental elements exert a significant influence on coffee bean characteristics and, consequently, the overall quality of the final coffee product. Notably, this dependence on environmental conditions presents a unique criterion for coffee quality assessment, potentially adding value to the offerings of individual coffee farmers.

## ACKNOWLEDGEMENT

This research was supported by National Science, Research and Innovation Fund (NSRF) and Prince of

Songkla University (Fundamental Fund (FF): Grant no. NAT6505025M and NAT6505025a).

## Conflict of interest

All authors declare that there is no conflict of interest.

## REFERENCES

- Ababu, D.G. and Getahun, A.M. (2021). Time series analysis of price of coffee in case of Mettu Town, Ilu Ababor Zone, Oromia Regional State, Ethiopia. *Asian Journal of Dairy and Food Research*. 40(3): 279-284. <https://doi.org/10.18805/ajdfr.DR-204>.
- Ahmed, S., Brinkley, S., Smith, E., Sela, A., Theisen, M., Thibodeau, C., Warne, T., Anderson, E., van Dusen, N., Giuliano, P., Ionescu, K.E. and Cash, S.B. (2021). Climate change and coffee quality: systematic review on the effects of environmental and management variation on secondary metabolites and sensory attributes of *Coffea arabica* and *Coffea canephora*. *Frontiers in Plant Science* 12. <https://doi.org/10.3389/fpls.2021.708013>.
- Bunn, C., Läderach, P., Rivera, O.O. and Kirschke, D. (2015). A bitter cup: Climate change profile of global production of Arabica and Robusta coffee. *Climatic Change*. 129: 89-101. <https://doi.org/10.1007/s10584-014-1306-x>.
- Bunya-atichart, K., Nokkoul, R., Rattanadet, N. and Boonmee, W. (2018). Pre and post-harvest managements and some qualities of Robusta coffee fruits in the Chumphon area. *Songklanakarin Journal of Plant Science*. 5: 42-48.
- Cassamo, C.T., Manguzeze, A.V.J., Leitão, A.E., Pais, I.P., Moreira, R., Campa, C. *et al.* (2022). Shade and altitude implications on the physical and chemical attributes of green coffee beans from Gorongosa Mountain, Mozambique. *Agronomy*. 12(10): 2540. <https://doi.org/10.3390/agronomy12102540>.
- Chiarawipa, R. and Sirikantayakul, C. (2015). Phenotypic plasticity in Robusta coffee trees in a mixed orchard system. *Agricultural Science Journal*. 46(3): 433-436.
- Chiarawipa, R., Suteekanjanonthai, P. and Somboonsuke, B. (2021). Adaptive ecophysiological characteristics of leaves and root distribution of Robusta coffee saplings in relation to rubber ages under an intercropping system. *Journal of Agricultural Science and Technology*. 23(2): 387-402. <http://jast.modares.ac.ir/article-23-37029-en.html>.
- Coffee Quality Institute. (2018). Fine Robusta Standards and Protocols. Available: <https://www.scith.coffee/wp-content/uploads/2021/03/Fine-Robusta-Standards-Protocols.pdf>.
- DaMatta, F.M., Avila, R.T., Cardoso, A.A., Martins, S.C. and Ramalho, J.C. (2018). Physiological and agronomic performance of the coffee crop in the context of climate change and global warming: A review. *Journal of Agricultural and Food Chemistry*. 66: 5264-5274. <https://doi.org/10.1021/acs.jafc.7b04537>.
- Gonzalez-Sanchez, B., Sandoval-Gonzalez, O., Flores-Cuautle, J.d.J.A., Landeta-Escamilla, O., Portillo-Rodriguez, O. and Aguila-Rodriguez, G. (2024). A study of the physical characteristics and defects of green coffee beans that influence the sensory notes using machine learning models. *Processes* 12, <https://doi.org/10.3390/pr12010018>.

- Hagos, M., Redi-Abshiro, M., Chandravanshi, B.S., Ele, E., Mohammed, A.M. and Mamo, H. (2018). Correlation between caffeine contents of green coffee beans and altitudes of the coffee plants grown in southwest Ethiopia. *Bulletin of the Chemical Society of Ethiopia*. 32: 13-25. <https://dx.doi.org/10.4314/bcse.v32i1.2>.
- Halagarda, M. and Obrok, P. (2023). Influence of post-harvest processing on functional properties of coffee (*Coffea arabica* L.). *Molecules*. 28. <https://doi.org/10.3390/molecules28217386>.
- Hameed, A., Hussain, S.A. and Suleria, H.A.R. (2018). "Coffee Bean-Related" Agroecological Factors Affecting the Coffee. *Co-Evolution of Secondary Metabolites, Reference Series in Phytochemistry*. [Merillon, J.M. and Ramawat, K. (Eds.)] Springer, Cham.
- Hareesh, S.B. and Jayarama (2024). Nutritional composition of green Robusta coffee (*Coffea canephora*) beans under organic and integrated nutrition with varying shade and irrigation management practices in Western Ghats of India. *Asian Journal of Dairy and Food Research*. 43(1): 17-24. <https://doi.org/10.18805/ajdfr.DR-1679>.
- Kath, J., Byrareddy, V., Mushtaq, S., Craparo, A. and Porcel, M. (2021). Temperature and rainfall impact on Robusta coffee bean characteristics. *Climate Risk Management*. 32: <https://doi.org/10.1016/j.crm.2021.100281>.
- Kattalee, P. and Inprasit, C. (2012). Physical Property of Robusta Green Coffee Beans Before and After Roasting. The 9<sup>th</sup> National Symposium. Kasetsart University, Kamphaeng Saen Campus, Thailand. pp 97-104.
- Legesse, A. (2022). Climate change effect on coffee yield and quality: A review. *Food Science and Quality Management*. 113. <https://doi.org/10.7176/FSQM/113-01>.
- Leite, H.G., De Oliveira, L.S. and De Lima, E.V. (2015). Effects of fertilization on yield, size and quality of coffee beans. *African Journal of Agricultural Research*. 10(25): 2498-2505.
- Lina, S., Sugeng, P. and Budi, P. (2022). Influence of various soil and water conservation methods on the moisture balance at coffee plant root zone. *Indian Journal of Agricultural Research*. 56(3): 325-330. <https://doi.org/10.18805/IJARE.AF-703>.
- Mohsenin, N.N. (1971). *Physical Properties of Plant and Animal Materials* (v.1): Physical Characteristics and Mechanical Properties (1<sup>st</sup> Ed.). Routledge. <https://doi.org/10.4324/9781003062325>.
- Office of Agricultural Economics. (2022). *Agricultural Product Production Information: Coffee*. Available online: <https://www.oae.go.th/assets/portals/1/fileups/prcaidata/files/coffee%2065.pdf>.
- Rimbing J., Engka R., Dumalang, S. and Rorong, F. (2023). Response of the coffee berry borer (*Hypothenemus hampei*) to attractant traps in Robusta coffee plantations. *The Indian Journal of Agricultural Sciences*. 93(5): 544-548. <https://doi.org/10.56093/ijas.v93i5.133017>.
- Specialty Coffee Association. (2018). *Green coffee standards*. Coffee Standards, Specialty Coffee Association. California. pp 5-7.
- Tolessa, K., D'heer, J., Duchateau, L. and Boeckx, P. (2017). Influence of growing altitude, shade and harvest period on quality and biochemical composition of Ethiopian specialty coffee. *Journal of the Science of Food and Agriculture*. 97: 2849-2857. <https://doi.org/10.1002/jsfa.8114>.
- Wintgens, J.N. (2012). *Harvesting and Green Coffee Processing*. Coffee: Growing, Processing, Sustainable Production. pp 604-715. Wiley-VCH Verlag GmbH and Co. KGaA, Weinheim.
- Worku, M., de Meulenaer, B., Duchateau, L. and Boeckx, P. (2018). Effect of altitude on biochemical composition and quality of green arabica coffee beans can be affected by shade and postharvest processing method. *Food Research International*. 105: 278-285. <https://doi.org/10.1016/j.foodres.2017.11.016>.