



The Potential Impacts of Selecting Viable Oocytes on Further Embryonic Development in Mammals: A Review

A.A. Mohammed¹, S. Al-Suwaiegh¹, I. AlGherair¹, A. Mohammed², A. Mohammed²

10.18805/IJAR.BF-1789

ABSTRACT

Selecting viable fully grown germinal and matured oocytes were positively effect on further embryonic development and live birth. The use of feed and hormonal supplementation and biotechnologies to enhance oocyte competence was evaluated. In addition, the oocytes were selected based on ovarian follicular stage, follicles and oocytes morphology, follicular fluid characters and gene expression. Changes were obtained in oocyte maturation, fertilization and further embryo cleavage and development over selecting viable oocytes. The larger follicle diameter, the translucent follicle and the higher blood supply the higher developmental competence of resulting oocytes. The higher surrounding cumulus cells and homogeneity of cytoplasm the higher developmental competence of oocytes. In addition, oocytes obtained during luteal stage compared to those obtained during follicular phase were highly or comparably developed to embryos. Furthermore, follicular fluid characters and gene expression might predict the embryonic developmental competence of oocytes in addition to their viable staining with brilliant cresyl blue. In conclusion, the characters of ovarian follicles, oocytes, follicular fluid and gene expression have significant implications on further oocyte developmental competence in different mammalian species.

Key words: Embryos development, Follicles, Follicular fluid, Gene expression, Oocytes morphology.

Significant strides of progresses is achieved in selecting the higher developmental competence oocytes through the characters of ovarian follicles, oocytes morphology, follicular fluid and gene expression (Mohammed *et al.*, 2005; Mlyczyńska *et al.*, 2023). It has been known that several factors affect the ovarian follicle diameters and the contents of oocytes and follicular fluid parameters (Contreras-Solís *et al.*, 2021; Castro-Modesto *et al.*, 2022; Mohammed *et al.*, 2020, 2024a,b,c,d; Cañón-Beltrán *et al.*, 2024).

Oocyte developmental competence is gradually acquired during the processes of follicle growth and maturation (Al Zeidi *et al.*, 2022a,b; Mohammed *et al.*, 2022; Aljubran *et al.*, 2023). Modification of follicle environment through the use of feed and hormonal supplementation and biotechnologies to enhance oocyte competence was confirmed in several studies (Gordon 2003; Costa *et al.*, 2023; Mohammed *et al.*, 2024a,b,c). Several studies concerning the nutrition and feed additives, hormonal supplementation indicated that they were pivotal factors in determining the ovarian follicle development and oocyte quality that resulted in varied reproductive performances in different animal species and human (Mohammed 2018, 2019; Mohammed *et al.*, 2022; 2024a,b,c,d; Ali *et al.*, 2021; Al-Mufarji *et al.*, 2022a,b; Mohammed and Al-Suwaiegh 2023; Al-Mafurji *et al.*, 2023). Bragança *et al.* (2020) found that progesterone administered to sheep did not affect the pattern of ovarian first wave follicle populations and the resulting oocyte quality during ovarian stimulation with FSH. Primordial follicle activation and survival during *in vitro* culture of bovine ovarian tissue to control oxidative stress upon melatonin treatment was confirmed through different mechanisms (Silva *et al.*, 2024). Lucia Dos Santos Silva

¹Department of Animal and Fish Production, College of Agriculture and Food Sciences, King Faisal University, P.O. Box 402, Al-Ahsa 31982, KSA.

²College of Human Medicine, Assiut University, Egypt.

Corresponding Author: A.A. Mohammed, Department of Animal and Fish Production, College of Agriculture and Food Sciences, King Faisal University, P.O. Box 402, Al-Ahsa 31982, KSA.
Email: aamohammed@kfu.edu.sa

How to cite this article: Mohammed, A.A., Al-Suwaiegh, S., AlGherair, I., Mohammed, A. and Mohammed, A. (2024). The Potential Impacts of Selecting Viable Oocytes on Further Embryonic Development in Mammals: A Review. Indian Journal of Animal Research. 58(9): 1435-1443. doi: 10.18805/IJAR.BF-1789.

Submitted: 27-03-2024 **Accepted:** 30-04-2024 **Online:** 18-05-2024

et al. (2023) reviewed the role of antioxidants in prevention follicular damage caused by oxidative stress in the *in vitro* culture of preantral follicles. Therefore, our objectives were to highlight key aspects of nutrition, feed supplements and hormones on follicle and oocyte development in addition to selecting viable oocytes according to follicular and luteal cycle stages, morphology of follicles and oocytes, follicular fluid characters and gene expression (Fig 1-3).

The current study was carried out according to the procedure approved by Deanship of Scientific Research, King Faisal University, Saudi Arabi from September to March 2024. The earlier reports concerning the effects of nutrition, feed supplements and hormones in addition to follicular and luteal stages on oocytes developmental competence were utilized. In addition, the research papers concerning follicle and oocyte morphology, follicular fluid and gene

expression in addition to assisted reproductive techniques were reviewed and discussed for selecting viable oocytes, increasing outcomes of reproduction and treatment of infertility. Therefore, our targets are to highlight key aspects of nutrition, feed supplements and hormones on follicle and oocyte development followed by selecting viable oocytes based on follicular and luteal estrous stages, morphology of follicles and oocytes, follicular fluid characters and gene expression.

Most reproduction management technologies and interventions are centered on the ovarian follicle development and the contents of oocytes and follicular fluid.

Nutrition and supplements

A plethora of molecules modified by the nutritional intake as amino acids, pyruvate, fatty acids, inorganic ions, second messengers, cytokines, growth factors effect on ovarian follicle development, oocyte quality, maturation and further embryonic development and live birth in different mammalian species (Mohammed 2018; Mohammed *et al.*, 2020, 2024a,c; Hye *et al.*, 2020; Thuy Van *et al.*, 2020; Fabozzi *et al.*, 2021; Costa *et al.*, 2023; Garza *et al.*, 2023; Silva *et al.*, 2023; Liu *et al.*, 2024).

Protein played a crucial role in both follicle growth and development and the quality of resulting oocytes (Dupont *et al.*, 2014; Gao 2020; Rubio *et al.*, 2021; Machado *et al.*, 2020). Amino acids served as building blocks for protein synthesis that were essential for follicles' growth and development, granulosa cell proliferation and oocyte quality. Protein in the oocyte was used for synthesis of RNA and to store materials for sustaining future embryo development until maternal zygotic transition or genome activation, which occurred in most of mammalian species between the two- to the eight-cell stages (Li *et al.*, 2013). In addition, certain proteins function as antioxidants, protecting follicles and oocytes from oxidative stress, which damage DNA and impair their development. On the other hand, unbalanced protein diet resulted in alterations of structure, function and biogenesis with increased levels of reactive oxygen species (Schutt *et al.*, 2019).

Carbohydrates were essential sources of energy for the body function including the reproductive system (Dupont *et al.*, 2014; Fabozzi *et al.*, 2021). Ovarian follicular growth and development and oocytes require a readily available supply of energy through carbohydrates. Report of Mohammed *et al.* (2021) indicated accelerated and higher development of ovarian follicle with propylene glycol supplementation. On the other hand, the potential negative effects of high carbohydrate intake over increased oxidative stress and hormonal imbalances could be damaging to DNA and cellular components within the follicles and oocytes.

There are molecular mechanisms of functional fatty acids affecting follicular development, granulosa cell steroidogenesis and oocyte maturation (Zeng *et al.*, 2023). There was a strong relationship between fat supplementation and follicle size and oocyte quality (Khatir *et al.*, 2007; Yang

et al., 2016; Labrecque *et al.*, 2016). The unsaturated fatty acids yielded positive effects on reproductive performance but the saturated fatty acids were linked to negative outcomes as decreased follicle number and quality (Zeng *et al.*, 2023). On the other hand, the negative effects of high-fat diets included increased inflammation and oxidative stress, hormonal imbalances and decreased blood flow (Dupont *et al.*, 2014). Collectively, abnormal dietary consumption of carbohydrates, fatty acids, proteins, vitamins and minerals, had detrimental effects on ovarian functions, affecting not only the oocyte quality but also the quality of embryos and implantation of a healthy embryo (Noli *et al.*, 2020).

Feed supplements

Effects of nutritive and non-nutritive feed supplements on reproductive performances in mammals has been discussed (Mohammed *et al.*, 2024a,b). The nutritive and non-nutritive feed supplements change rumen microorganism, feed utilization, body weight gain, blood and plasma profiles (Mohammed *et al.*, 2024a,b,c). Such improvements over feed supplement/s lead to improvement in ovarian follicle development and the resulting oocyte quality, embryos and newborns in addition to amelioration of reproductive dysfunctions (Al Mafurji *et al.*, 2022a,b).

The essential macronutrient and antioxidant micronutrient improved the reproductive performances through significant increase in concentrations of reproductive hormones, numbers and sizes of ovarian follicles in addition to increase in oocyte and embryo quality (Senosy *et al.*, 2017, 2018, 2019; Mohammed 2018; Ali *et al.*, 2021). In addition, green algae as a source of beta-carotene was supplemented to support oocyte and embryo quality in goats (Silva *et al.*, 2023). The microalgae *Chlorella* did not enhance the quality of oocytes whereas it improved the quality of embryos and stimulated their mitochondrial function. Previous reports on *Dunallela salina* resulted in improvement in levels of reproductive hormones, oocytes and embryo development and (Senosy *et al.*, 2017; Mohammed, 2018; Ali *et al.*, 2021). Therefore, the continuous research in nutritive and non-nutritive feed supplements was necessitated for both animals and humans. Therefore, the future prospective studies to find beneficial new feed additives was an emerging approach for improving ovarian structural development and the resulting quality of oocytes and embryos (Mohammed *et al.*, 2024a,b,c,d).

Hormonal supplementation

Hormonal supplementation had various effects on ovarian follicle development, oocyte quality and embryo development, depending on the specific hormone, dosage and body health condition in mammalian species (Gordon, 2003; Mohammed and Attaai, 2011). The important hormones related to ovarian follicular growth and maturation were follicle stimulating hormone, luteinizing hormone (LH), estrogen and progesterone. Follicle stimulating hormone (FSH) was used routinely in protocols

of assisted reproductive techniques but excess FSH could lead to overstimulation and complications. Supplementation of LH and estrogen in specific contexts could support ovarian follicle and oocyte maturation but required careful management due to potential adverse effects. It was well-known that the physiological effects of progesterone during the pre-implantation stages of mammalian embryos was mediated by progesterone receptors and their gene expression. Although progesterone could not improve *in vitro* maturation rates, culture media supplemented with progesterone significantly improved mouse embryo development (Andersen, 1990; Mendoza *et al.*, 2002; Gordon,

2003; Salehnia and Zavareh, 2013; Flores-Herrera *et al.*, 2008). Growth hormone injection improved follicles and oocytes development in prepubertal lambs (Liu *et al.*, 2023). Collectively, hormonal supplementation could be double-edged to either give a positive or negative effect depending on several factors.

Follicular and luteal stage of the cycle

Ovarian follicles develop through two processes called *initial* and *cyclic* recruitments (McGee and Hsueh, 2000), which describe recruitment of dormant primordial follicles continuously into the growing follicle and antral follicles,




Fig 1: Effects of nutrition on ovarian follicle development and oocyte quality.




Fig 2: Factors responsible for selection of viable oocytes.




Fig 3: Morphological, molecular and metabolic assessment of oocyte quality.

respectively. After puberty, multiple antral ovarian follicular waves (2, 3 and 4 waves) were reported in several studies (Gordon 2003; Baerwald and Pierson, 2020) during ruminant estrous or human menstrual cycles. It was estimated for bovine early antral follicles to take about 40 days to progress to preovulatory follicles.

The estrous cycle was divided to follicular and luteal phases (Gordon, 2003). Ovarian follicular waves or follicular and luteal stages were factors affecting the quality of oocytes, oocyte maturation and resulting embryo development (Mohammed *et al.*, 2012, 2022). Some studies suggested that the estrous stage with its hormonal fluctuations can potentially influence oocyte quality in different mammalian species (Gordon, 2003; Majumdar *et al.*, 2023; Dastjerdi *et al.*, 2024). Some studies suggest higher oocyte quality in early follicular phase due to more favorable hormonal environment. Some other studies suggest negative impact of the corpus luteum on oocyte quality, particularly on smaller follicles.

Evidence concerning the importance of progesterone support during ovum pick-up and *in vitro* embryo production (OPU-IVP) was ascribed in studies related to transvaginal follicular aspiration in cattle and improvement of *in vitro* embryo production (Simmons *et al.*, 2023). For treatment of fertility disorders, comparison of abnormalities during follicular phase *in vitro* maturation of oocytes compared with luteal phase *in vitro* maturation in women was investigated (Hatirnaz *et al.*, 2023). The results indicated differences in maturation and fertilization rates, quality of embryos or pregnancy outcomes between follicular phase and luteal phase in women with oocyte maturation abnormalities. Additionally, Suñol *et al.* (2023) compared follicular phase stimulation versus luteal phase stimulation in suboptimal women responders. The oocyte yield did not increase in follicular-phase stimulation compared to luteal-phase stimulation. De Wit *et al.* (2000) also found that the stage of oestrous cycle had no effect on the distribution of the cumulus-oocyte complexes (COC) over the grades of oocytes qualities or on the developmental capacity of COC. In human, stimulation of

luteal phase in the same menstrual cycle was an effective strategy to rescue poor responders with follicular phase stimulation (Majumdar *et al.*, 2023; Dastjerdi *et al.*, 2024).

Morphology of follicles and quality of oocytes

The ovarian follicles are categorized to different groups including primordial follicles to preantral follicles and antral ovulatory follicles (Gordon 2003) (Fig 4-5). Healthy antral follicles larger than 3 mm contain competent oocytes that are competent to develop into preimplantation and post implantation embryos (Mohammed *et al.*, 2005). The sizes of ovarian follicles were different among mammalian species. They are categorized into small (<3 mm), medium (3-5 mm) and large follicles (>5 mm) in small ruminant species (Senosy *et al.*, 2017; Al Mufarji *et al.*, 2023). Sizes of ovulatory follicles naturally resulted in multiple ovulations and influenced on the development of corpus luteum and fertility in cattle (Echternkamp *et al.*, 2009).

The effect of follicle size on oocyte quality was a complex issue and generally a positive correlation was noted between follicle size and oocyte quality. Large ovulatory follicles (≥ 16 mm) are more likely to yield mature oocytes that are capable of normal fertilization and development into high-quality embryos compared to small follicles (<13 mm) which produced immature oocytes with lower fertilization and cleavage rates in large ruminant species and humans as well (Kahraman *et al.*, 2017). However, the relationship between follicle size and oocyte quality was not always clear and there could be exceptions where the age, follicular waves and health conditions, could also play a role in oocyte quality. Therefore, more research was needed to fully understand the complex interplay between follicle size and oocyte quality. Studies have shown that oocytes from larger follicles have higher fertilization rates and better competence for embryonic development (Mohammed unpublished report).

Follicular fluid characters

The correlation between follicular fluid characteristics and oocyte quality was complex and interesting field of research, with promising potential for improving assisted reproductive




Fig 4: Mouse ovary containing preantral and antral follicles.




Fig 5: Translucent ovarian follicles in sheep ovaries.

technologies (Mohammed *et al.*, 2005, 2020). Follicular fluid (FF) surrounds the oocyte within the ovarian follicle and provides a crucial microenvironment for growth and development of the oocyte. Analyzing the chemical composition of FF offered insights into the ovarian follicle health and viability of the resulting oocyte (Mohammed *et al.*, 2005; 2019a,b, 2020). Several studies had identified positive correlations between specific molecules in follicular fluid and oocyte quality such as insulin-like growth factors (IGFs) and their binding proteins (IGFBPs), which were often associated with higher maturation and fertilization rates of oocytes (Mohammed *et al.*, 2019a). In addition, the higher levels of total antioxidant capacity (TAC) in follicular fluid seemed to indicate the presence of high quality oocytes in mature follicles. Furthermore, metabolites and certain cytokines have also shown promising connections to oocyte quality (Mohammed *et al.* 2019a,b; 2020).

Follicular fluid composition changed during the estrous cycle due to follicular growth and development and nutritional level as well (Mohammed *et al.*, 2012). Other components still need further studies to define the changes at cellular level (Mohammed *et al.*, 2019b). Finally, the research on the correlation between follicular fluid characteristics and oocyte quality was promising but still evolving due to differences in experimental design, animal species and the analytical methods used (Mohammed *et al.*, 2020).

Gene expression

Complex relationship was noted between gene expression and oocyte quality. Gene expression in oocytes was

influenced by intrinsic and extrinsic factors (Jentoft *et al.*, 2023). The intrinsic factors include genetic encoding, age of female and epigenetic modifications. The extrinsic factors include follicular environment, maternal health, environmental factors and *in vitro* culture conditions during assisted reproductive techniques.

Functional molecules were produced through gene expression at different stages of oocyte development that played crucial roles in various processes (Maside *et al.*, 2021). Gene expression regulated various aspects of oocytes including zona pellucida formation, spindle apparatus development, enzymes and molecules involved in energy production and utilization, molecules involved in the communication of oocyte with the surrounding cells (Wassarman and Litscher, 2021). For example, the zona pellucida genes encode sperm receptors, which were required for fertilization (Evsikov *et al.*, 2009). The nutritional factors such as protein or lipids helped in regulation of gene expression in developing follicles and oocytes, influencing various processes crucial for successful development and maturation (Schutt *et al.*, 2019; Sharma *et al.*, 2020; Alves *et al.*, 2021).

CONCLUSION

Nutritive and non-nutritive feed additives/supplements and hormonal injection are positively or negatively effect on ovarian follicle development and the resulting oocytes. Selecting viable oocytes was necessitated for assisted reproductive technology, which could be according to follicular and luteal phases of the estrous cycle, morphology of follicles and oocytes, follicular fluid

characters and gene expression. Therefore, the future prospective studies to find newer methods for selection of viable oocytes was a continuous approach for improving reproductive performances.

ACKNOWLEDGEMENT

The authors want to thank and acknowledge Deanship of Scientific Research, King Faisal University, Saudi Arabia for funding and support (GrantA055).

Conflict of interest

There is no conflict of interest for authors to declare.

REFERENCES

- Al Mufarji, A., Al-Suwaiegh, S., Mohammed, A.A. (2023). Influence of organic *Moringa oleifera* leaves supplemented during gestation and lactation periods: Modulation of Production efficiency, blood and metabolic parameters of ewes and lambs in subtropics. *Advances in Animals and Veterinary Sciences*. 7(1): 385-393.
- Al Zeidi R., Al Masruri H., Al Mufarji A., Al-Hassan Mohammed A.A. (2022a). Role of cumulus cells and follicular fluid on oocyte maturation and developmental competence of embryos: Intact and reconstructed oocytes. *Advances in Animals and Veterinary Sciences*. 10(6): 1219-1226.
- Al Zeidi, R., Al Masruri, H., Al Mufarji, A., Mohammed, A.A., Mohammed, H. (2022b). Changes in nucleus and cytoplasm during oocyte maturation: Involvement in embryo production. *Advances in Animals and Veterinary*. 10(9): 2081-2089.
- Ali, M.A., Alshaheen, T., Senosy, W., Kassab, A., Mohammed, A.A. (2021). Effects of feeding green microalgae and *Nigella sativa* on productive performance and metabolic profile of boer goats during peripartum period in subtropics. *Fresen. Environ. Bull.* 30: 8203-8212.
- Aljubran, S., Al-Suwaiegh, S., Alyousef, Y., Alhajri, S., Alghareeb, M., Mohammed, A.A. (2023). Roles of assisted reproductive techniques in mammals: Developmental competence of oocytes and embryos. *Advances in Animals and Veterinary Sciences*. 11(2): 252-263.
- Al-Mufarji, A., Mohammed, A.A., Al-Masruri, H., Al-Zeidi, R. (2022a). Effects of dietary microalgae supplementation on mammals' production and health. *Advances in Animal and Veterinary Sciences*. 10(8): 1718-1724.
- Al-Mufarji, A., Mohammed, A.A., Al-Zeidi, R., Al-Masruri, H., Mohammed, A. (2022b). Effects of *Moringa oleifera* on follicular development, blood and metabolic profiles of subtropical ewes during peripartum. *Advances in Animal and Veterinary Sciences*. 10(8): 1706-1712.
- Alves, J.P.M., CCLFernandes, C.C.L., CEMCalderón, C.E.M., Rafael Rossetto, R., Marcelo Bertolini, M., Davide Rondina, D. (2021). Short-term supplementation of diets rich in lipids or glycogen precursors can affect intra-follicular environment, oocyte mitochondrial gene expression and embryo development following parthenogenesis in goat. *Small Ruminant Research*. 194: 106279. <https://doi.org/10.1016/j.smallrumres.2020.106279>.
- Andersen, C.Y. (1990). Levels of steroid-binding proteins and steroids in human preovulatory follicle fluid and serum as predictors of success in *in vitro* fertilization-embryo transfer treatment. *Journal of Clinical Endocrinology and Metabolism*. 71(5): 1375-1381.
- Baerwald, A., Pierson, R. (2020). Ovarian follicular waves during the menstrual cycle: physiologic insights into novel approaches for ovarian stimulation. *Fertility. Sterility*. 114 (3): 443-457.
- Bragança, G.M., Souza-Fabjan, J.M.G., Ribeiro, L.S., Brair, V.L. and Brandão, F.Z. (2020). Exogenous progesterone does not affect first-wave follicle populations and oocyte quality during ovarian stimulation with FSH in sheep. *Domestic Animal Endocrinology*. 72: 106369. <https://doi.org/10.1016/j.domaniend.2019.06.001>.
- Cañón-Beltrán, K., García-García, R.M., Cajas, Y.N., Fierro, N., Lorenzo, P.L., Arias-Álvarez, M. (2024). Improvement of oocyte competence and *in vitro* oocyte maturation with EGF and IGF-I in Guinea pig model. *Theriogenology*. 214: 206-214.
- Castro-Modesto, T., Mamani, P., Pella, R., Bravo, Z., Villarreal-Ugarte, S., Cancho, C., Aguila, L., Ratto, M.H., Valdivia, M. (2022). Effect of follicle size on *in vitro* maturation in alpaca oocytes (*Vicugna pacos*) and the first ICSI in alpaca species. *Small Ruminant Research*. 213: 106680. <https://doi.org/10.1016/j.smallrumres.2022.106680>.
- Contreras-Solis, I., Catalá, M., Soto-Heras, S., Roura, M., Paramio, M.T., Izquierdo, D. (2021). Effect of follicle size on hormonal status of follicular fluid, oocyte ATP content and *in vitro* embryo production in prepubertal sheep. *Domestic Animal Endocrinology*. 75: 106582. doi: 10.1016/j.domaniend.2020.106582.
- Costa, C.B., Fair, T., Seneda, M.M. (2023). Review: Environment of the ovulatory follicle: Modifications and use of biotechnologies to enhance oocyte competence and increase fertility in cattle. *Animal*. 17(1): 100866. doi: 10.1016/j.animal.2023.100866.
- Dastjerdi, M.V., Ansari-pour, S., Ataei, M. (2024). Comparison of luteal phase stimulation with follicular phase stimulation in poor ovarian response: A single-blinded randomized controlled trial. *Contraception and Reproductive Medicine*. 9: 1186. <https://doi.org/10.1186/s40834-024-00265-z>.
- de Wit, A.A.C., Wurth, Y.A., Th, Y.A., Kruip, A.M., Th, A.M. (2000). Effect of ovarian phase and follicle quality on morphology and developmental capacity of the bovine cumulus-oocyte complex. *Journal of Animal Science*. 78: 1277-1283.
- Dupont, J., Scaramuzzi, R.J., Reverchon, M. (2014). The effect of nutrition and metabolic status on the development of follicles, oocytes and embryos in ruminants. *Animal*. 8(7): 1031-1044.
- Echternkamp, S.E., Cushman, R.A., Allan, M.F. (2009). Allan size of ovulatory follicles in cattle expressing multiple ovulations naturally and its influence on corpus luteum development and fertility. *Journal of Animal Science*. 87: 3556-3568.
- Evsikov, A.V., Marín de Evsikova, C. (2009). Gene expression during the oocyte-to-embryo transition in mammals. *Molecular Reproduction Development*. 76(9): 805-818.

- Fabozzi, G., Iussig, B., Cimadomo, D., Vaiarelli, A., Maggiulli, R., Ubaldi, N., Ubaldi, F.M., Rienzi, L. (2021). The impact of unbalanced maternal nutritional intakes on oocyte mitochondrial activity: Implications for reproductive function. *Antioxidants (Basel)*. 10(1): 91. doi: 10.3390/antiox10010091.
- Flores-Herrera, H., Diaz-Cervantes, P., De la Mora, G., Zaga-Clavellina, V., Uribe-Salas, F., Castro, I. (2008). A possible role of progesterone receptor in mouse oocyte *in vitro* fertilization regulated by norethisterone and its reduced metabolite. *Contraception*. 78(6): 507-512.
- Gao, H. (2020). Amino acids in reproductive nutrition and health. *Advances in Experimental Medicine and Biology*. 1265: 111-131.
- Garza, V., West, S.M., Cardoso, R.C. (2023). Review: Gestational and postnatal nutritional effects on the neuroendocrine control of puberty and subsequent reproductive performance in heifers. *Animal*. 17(1): 100782. <https://doi.org/10.1016/j.animal.2023.100782>.
- Gordon, I. (2003). The Bovine Oestrous Cycle and Associated Events. In: *Laboratory Production of Cattle Embryos (2nd)* CABI Publishing UK. 42-78.
- Hatirnaz, S., Ebru Hatirnaz, E., Sebati Sinan Ürkmez, S.S., Samettin Çelik, S., YCÜrkmez, Y.C., Mingju Cao, M., SLTan, S.L., MH. Dahan, M.H. (2023). The comparison of luteal phase *in vitro* maturation with follicular phase *in vitro* maturation in women with oocyte maturation abnormalities. *Reproductive BioMedicine Online*. 48(4): 103648. doi: 10.1016/j.rbmo.2023.103648.
- Hye, N., Klein-Jöbstl, D., Blessing, A., Burmeister, J., Hamann, N., Aurich, C., Drillich, M. (2020). Effect of two postpartum intramuscular treatments with β -carotene (Carofertin®) on the blood concentration of β -carotene and on the reproductive performance parameters of dairy cows. *Theriogenology*. 148: 1-7.
- Jentoft, I.M.A., Baüerlein, F.J.B., Welp, L.M., Cooper, B.H., Petrovic, A., So, A. *et al.* (2023). Mammalian oocytes store proteins for the early embryo on cytoplasmic lattices. *Cell*. 186: 5308-5327.
- Kahraman, S., Cetinkaya, C.P., Cetinkaya, M., Yelke, H., Colakoglu, Y.K., Aygun, M., Montag, M. (2017). The effect of follicle size and homogeneity of follicular development on the morphokinetics of human embryos. *Journal of Assisted Reproduction and Genetics*. 34(7): 895-903.
- Khatir, H., Anouassi, A., Tibary, A. (2007). Effect of follicular size on *in vitro* developmental competence of oocytes and viability of embryos after transfer in the dromedary (*Camelus dromedarius*). *Animal Reproduction Science*. 99(3-4): 413-20.
- Labrecque, R., Fournier, E., Sirard, M.A. (2016). Transcriptome analysis of bovine oocytes from distinct follicle sizes: Insights from correlation network analysis. *Molecular Reproductive Development*. 83(6): 558-569.
- Li, L., Lu, X., Dean, J. (2013). The maternal to zygotic transition in mammals. *Aspects of Molecular Medicine*. 34(5): 919-938.
- Liu, K., Zhang, L., Qi, Q., Li, J., Yan, F., Hou, J. (2023). Growth hormone treatment improves the development of follicles and oocytes in prepubertal lambs. *Journal of Ovarian Research*. 16(1): 132. doi: 10.1186/s13048-023-01209-y.
- Liu, M., Chen, R., Wang, T., Ding, Y., Zhang, Y., Huang, G., Huang, J., Qu, Q., Weijie Lv, Shining Guo (2024). Dietary Chinese herbal mixture supplementation improves production performance by regulating reproductive hormones, antioxidant capacity, immunity and intestinal health of broiler breeders. *Poultry Science*. 103(1): 103201. <https://doi.org/10.1016/j.psj.2023.103201>.
- Lucia Dos Santos Silva, R., de Sousa Barberino, R., Tavares de Matos, M.H. (2023). Impact of antioxidant supplementation during *in vitro* culture of ovarian preantral follicles: A review. *Theriogenology*. 207: 110-122.
- Machado, A.F., Guimarães, S.E.F., Guimarães, J.D., Santos, G.M., Silva, A.L., Silva, Y.F.R.S., Lollobrigida Netto, D.S., Correa, P.V.F., Marcondes, M.I. (2020). Effect of protein supplement level on the productive and reproductive parameters of replacement heifers managed in intensive grazing systems. *Plos One*. 15(10): e0239786. <https://doi.org/10.1371/journal.pone.0239786>.
- Majumdar, A., Majumdar, G., Tiwari, N., Singh, A., Gupta, S.M., Satwik, R. (2023). Luteal phase stimulation in the same cycle is an effective strategy to rescue poseidon poor responders with no embryos after the first follicular stimulation. *Journal of Human Reproductive Sciences*. 16(3): 218-226.
- Maside, C., Sánchez-Ajofrín, I., Medina-Chávez, D., Alves, B., Garde, J.J., Soler, A.J. (2021). Oocyte morphometric assessment and gene expression profiling of oocytes and cumulus cells as biomarkers of oocyte competence in sheep. *Animals*. 11(10): 2818. doi: 10.3390/ani11102818.
- McGee, E.A., Hsueh, A.J.W. (2000). Initial and cyclic recruitment of ovarian follicles. *Endocr. Rev*. 21(2): 200-214.
- Mendoza, C., Ruiz-Requena, E., Ortega, E., Cremades, N., Martinez, F., Bernabeu, R. (2002). Follicular fluid markers of oocyte developmental potential. *Human Reproduction*. 17(4): 1017-1022.
- Mlyczyńska, E., Kurowska, P., Rytelewska, E., Zaobina, E., Pich, K., Kieżun, M., Dobrzyń, K., Kisieleska, K., Kopij, G., Smolińska, N., Kamiński, T., Rak, A. (2023). Expression of visfatin in the ovarian follicles of prepubertal and mature gilts and *in vitro* effect of gonadotropins, insulin, steroids and prostaglandins on visfatin levels. *Theriogenology*. 211: 28-39.
- Mohammed, A.A., Al Mufarji, A., Alawaid, S. (2022). Developmental potential of ovarian follicles in mammals, involvement in assisted reproductive techniques. *Pakistan Journal Zoology*. 1-11.
- Mohammed, A.A., Attaai, A.H. (2011). Effects of dietary urea on timing of embryo cleavages and blood components in Mice. *Veterinary World*. 4(8): 360-363.
- Mohammed, A.A. (2018). Development of oocytes and preimplantation embryos of mice fed diet supplemented with *Dunaliella salina*. *Advances in Animal and Veterinary Sciences*. 6 (1): 33-39.
- Mohammed, A.A. (2019). *Nigella sativa* oil improves physiological parameters, oocyte quality after ovarian transplantation and reproductive performance of female mice. *Pakistan Journal of Zoology*. 51(6): 2225-2231.

- Mohammed, A.A., Abd El-Hafiz, G.A., Ziyadah, H.M.S. (2012). Effect of dietary urea on ovarian structures in Saidi ewes during follicular and luteal phases. *Egyptian Journal of Animal Production*. 49(1): 29-35.
- Mohammed, A.A., AlGherair I., Al-Suwaiegh S., Al-Khamis S., Alessa, F., Al-Madni, A. and Al-Ghamdi, A. (2024a). Effects of nutritive and non-nutritive feed supplements on feed utilization, growth and reproductive performances in mammals. *Indian Journal Animal Research*. 58(4): 615-621. doi: 10.18805/IJAR.BF-1744.
- Mohammed, A.A., Al-Hizab, F., Al-Suwaiegh, S., Alshaheen, T., Kassab, A., Hamdon, H., Senosy, W. (2021). Effects of propylene glycol on ovarian Restoration, reproductive performance, metabolic status and milk production of farafra ewes in subtropics. *Fresenius Environmental Bulletin*. 30(7): 8192-8202.
- Mohammed, A.A., Al-Saiady, M., El-Waziry, A., Alshaheen, T. (2024d). Effects of dietary omega-3 fatty acids on reproductive performance and biochemical parameters of lactating cows in arid subtropics. *Pakistan Journal of Zoology*. (Accepted).
- Mohammed, A.A., Al-Shaheen, T., Al-Suwaiegh, S. (2019a). Changes of follicular fluid composition during estrous cycle: The effects on oocyte maturation and embryo development *in vitro*. *Indian Journal of Animal Research*. 54(7): 797-804. doi: 10.18805/ijar.B-1030.
- Mohammed, A.A., Al-Shaheen, T. and Al-Suwaiegh, S. (2020). Effects of *Myo-inositol* on physiological and reproductive traits through blood parameters, oocyte quality and embryo transfer in mice. *Indian Journal of Animal Research*. doi: 10.18805/ijar.B-1300.
- Mohammed, A.A., Al-Suwaiegh, S., AlGherair, I., Al-Khamis, S., Alessa, F., Al-Awaid, S., Alhujaili, W.F., Mohammed, A., Mohammed, A. (2024b). The potential impacts of antioxidant micronutrients on productive and reproductive performances of mammalian species during stressful conditions in mammals. *Indian Journal Animal Research*. 58(6): 1031-1038. doi: 10.18805/IJAR.BF-1773.
- Mohammed, A.A., Al-Suwaiegh, S., AlGherair, I., Al-Khamis, S., Al-Awaid, S., Al-Sornokh, H., Alhujaili, W.F., Mohammed, A., Mohammed, A. (2024c). Morphological characteristics of ovarian tissues and follicular fluid metabolites of female lambs and ewes in subtropics. *Indian Journal Animal Research*. doi: 10.18805/IJAR.BF-1756.
- Mohammed, A.A., Al-Suwaiegh, S., Al-Shaheen, T. (2019b). Effects of follicular fluid components on oocyte maturation and embryo development *in vivo* and *in vitro*. *Advances Animal Veterinary Sciences*. 7(5): 346-355.
- Mohammed, A.A., Karasiewicz, J., Papis, K., Modlinski, J.A. (2005). Oocyte maturation in the presence of randomly pooled follicular fluid increases bovine blastocyst yield *in vitro*. *Journal Animal Feed Sciences*. 14: 501-512.
- Mohammed, A.A., Al-Suwaiegh, S. (2023). Impacts of *Nigella sativa* inclusion during gestation and lactation on ovarian follicle development, as well as the blood and metabolic profiles of ardi goats in subtropics. *Agriculture*. 13: 674. <https://doi.org/10.3390/agriculture13030674>.
- Noli, S.A., Ricci E., Cipriani S., Ferrari S., Castiglioni M., La Vecchia I., Somigliana E., Parazzini F. (2020). Dietary carbohydrate intake, dietary glycemic load and outcomes of *in vitro* fertilization: Findings from an observational Italian cohort study. *Nutrients*. 12(6): 1568. doi: 10.3390/nu12061568.
- Rubio, I., White, F.J., Spicer, L.J., Wettemann, R.P. (2021). Postpartum nutrition affects the insulin-like growth factor system in dominant follicles and plasma of anestrus beef cows. *Animal Reproduction Science*. 229: 106760. <https://doi.org/10.1016/j.anireprosci.2021.106760>.
- Salehnia, M., Zavareh, S. (2013). The effects of progesterone on oocyte maturation and embryo development. *International Journal of Fertility and Sterility*. 7(2): 74-81.
- Schutt, A.K., Blesson, C.S., Hsu, J.W., Valdes, C.T., Gibbons, W.E., Jahoor, F., Yallampalli, C. (2019). Preovulatory exposure to a protein-restricted diet disrupts amino acid kinetics and alters mitochondrial structure and function in the rat oocyte and is partially rescued by folic acid. *Reproductive Biology and Endocrinology*. 17(12). <https://doi.org/10.1186/s12958-019-0458-y>.
- Senosy W., Kassab, A.Y., Ezzat, A.A., Mohammed, A.A. (2019). Effect of copper supplementation on ovarian function and blood profiles of native goats in grazing copper-deficient alfalfa desert oasis. *Assiut Veterinary Medical Journal*. 65(160): 133-141.
- Senosy, W., Kassab, A.Y., Mohammed, A.A. (2017). Effects of feeding green microalgae on ovarian activity, reproductive hormones and metabolic parameters of boer goats in arid subtropics. *Theriogenology*. 96: 16-22.
- Senosy, W., Kassab, A.Y., Hamdon, H.A., Mohammed, A.A. (2018). Influence of organic phosphorus on reproductive performance and metabolic profiles of anoestrous Farafra ewes in subtropics at the end of breeding season. *Reproduction Domestic Animal*. 53: 904-913.
- Sharma, A., Baddela, V.S., Roettgen, V., Vernunft, A., Viergutz, T., Dannenberger, D., Hammon, H.M., Schoen, J. and Vanselow, J. (2020) Effects of dietary fatty acids on bovine oocyte competence and granulosa cells. *Frontiers in Endocrinology*. 11: 87. doi: 10.3389/fendo.2020.00087.
- Silva, B.R., Costa, F.C., De Lima Neto, M.F., Filho, F.F.C., de Assis, E.I.T., Aguiar, F.L.N. *et al.* (2024). Melatonin acts through different mechanisms to control oxidative stress and primordial follicle activation and survival during *in vitro* culture of bovine ovarian tissue. *Domestic Animal Endocrinology*. 86: 106824.
- Silva, M.R.L., Alves, J.P.M., Fernandes, C.C.L., Cavalcanti, C.M., Conde, A.J.H., *et al.* (2023). Use of green microalgae *Chlorella* as a nutritional supplement to support oocyte and embryo production in goats. *Animal Reproduction Science*. 256: 107296. doi: 10.1016/j.anireprosci.2023.107296.
- Simmons, R., Des, A.R., Guven-Ates, T.G., Kwong, W.Y., Randi, R.L.F., Sinclair, K.D. (2023). Enhanced progesterone support during stimulated cycles of transvaginal follicular aspiration improves bovine *in vitro* embryo production. *Theriogenology*. 199: 77-85.

- Suñol J., Castillo, J.C., Ortiz, J.A., Ten, J., Fuentes, A., Belén, M.B., María, M.M., *et al.* (2023). Conventional follicular phase ovarian stimulation versus luteal phase stimulation in suboptimal responders: A randomized controlled trial. *F and S Reports*. 2666-3341.
- Thuy Van, N.T., My, L.B.A., Van Thuan, N., Bui, H.T. (2020). Improve the developmental competence of porcine oocytes from small antral follicles by pre-maturation culture method. *Theriogenology*. 149: 139-148.
- Wassarman, P.M., Litscher, E.S. (2021). Zona pellucida genes and proteins: Essential players in mammalian oogenesis and fertility. *Genes*. 12(8): 1266. doi: 10.3390/genes12081266.
- Yang, M., Hall, J., Fan, Z.Q., Regouski, M., Meng, Q.G., Rutigliano, H.M. (2016). Oocytes from small and large follicles exhibit similar development competence following goat cloning despite their differences in meiotic and cytoplasmic maturation. *Theriogenology*. 86(9): 2302-2311.
- Zeng, X., Li, S., Liu, L., Cai, S., Ye, Q., Xue, B., Wang, X., Zhang, S., Chen, F., Cai, C., Wang, F., Zeng, X. (2023). Role of functional fatty acids in modulation of reproductive potential in livestock. *Journal of Reproductive Biology and Endocrinology*. 14(1): 24. <https://doi.org/10.1186/s40104-022-00818-9>.