



Exploring the Effects of Cold Plasma on Preserving and Processing of Foods: A Review

Sheela Punniyamorthy¹, Kanchana Shanmugavel², Chandra Prabha Shanmugavel¹,
Umamaheswari Thiagamoorthy³, Kokilavani Sakthivel⁴

10.18805/ajdfr.DR-2164

ABSTRACT

In the current scenario, global food productions need to be incredibly enormous to feed everyone. Though food supply plays a tremendous role in reaching out to the people, meanwhile advancement in food processing technology is a predominant factor in increasing food availability. In the food industry, there was a huge number of technologies for food processing that made food more convenient to eat and might be stable for extended storage. Among the food processing technologies, cold plasma may be one of the novel non-thermal technology that functions at atmospheric pressure or low-pressure processing technology. Cold plasma is an eco-friendly technique involved in food preservation, mainly destructing microbes in food thereby extending its shelf life. Cold plasma processes the food without degrading the physical, chemical, biological, nutritional and sensory qualities of food products. This review elaborates on cold plasma utilization and its quality changes in various foods.

Key words: Cold plasma, Food quality, Processing and preservation, Reactors.

Cold plasma is a novel evolving technology; which attracted scientists globally. It had been initially formed for perfecting the printing and binding properties of polymers. Within the last couple of decades, its application has become a strong tool for novel non-thermal processing methods (Chizoba *et al.*, 2017). Cold plasma consists of both charged and neutral particles including free radicals and helps to destroy or inactivate the microbes (Brany *et al.*, 2020). Microbial toxins and spores are predominant contaminants of foodstuffs. In the process of preparing food, these spores or microbes or microbial toxins are diminished or inactivated by the process of preservation or processing methods.

Most of the researchers examined plasma technology which will be used as a broad spectrum in the field of food preservation, biomedical and commercial applications (Sen *et al.*, 2018). Inward the food industry, the new trend is cold plasma technology which is highly profitable and safer. The custom of cold plasma in the food industry has been demonstrated for wastewater treatment (Sarangapani *et al.*, 2016), inactivating enzymes (Misra *et al.*, 2016), removing toxins (Misra *et al.*, 2015) and food decontamination (Misra *et al.*, 2011). Many researchers investigated, the application of cold plasma on fruits and vegetables using minimum levels successfully decontaminated harmful microorganisms like *Listeria monocytogenes*, *Escherichia coli* and *Salmonella typhimurium* (Ziuzina *et al.*, 2014; Jeyasena *et al.*, 2015; Albertos *et al.*, 2017). Thus, sterilization using cold plasma offers minimal cost with the least changes in food. Moreover, in the agriculture and food industries, there is an increasing need for innovative sustainable technologies due to the growing population, which increases the need for food, water and energy resources.

Cold plasma is a novel non-thermal processing technology which has exhibited noteworthy prospects within

¹SRM College of Agricultural Science, Baburayanpettai, Chengalpattu-603 201, Tamil Nadu, India.

²Community Science College and Research Institute, Madurai-625 104, Tamil Nadu, India.

³Anbil Dharmalingam Agricultural College and Research Institute, Trichy-620 009, Tamil Nadu, India.

⁴ICAR-Krishi Vigyan Kendra, Perambalur-621 115, Tamil Nadu, India.

Corresponding Author: Sheela Punniyamorthy, SRM College of Agricultural Science, Baburayanpettai, Chengalpattu-603 201, Tamil Nadu, India. Email: sheelap@srmist.edu.in

How to cite this article: Punniyamorthy, S., Shanmugavel, K., Shanmugavel, C.P., Thiagamoorthy, U. and Sakthivel, K. (2024). Exploring the Effects of Cold Plasma on Preserving and Processing of Foods: A Review. Asian Journal of Dairy and Food Research. 43(4): 624-630. doi: 10.18805/ajdfr.DR-2164.

Submitted: 04-09-2023

Accepted: 14-02-2024

Online: 11-03-2024

the food processing industries to produce safer foods. This technology helps to inactivate harmful microorganisms and extend the shelf life of foods (Pankaj and Keener, 2018). The purpose of this review is to provide an overview of cold plasma technology, its use in the food processing sector and the quality exposure of cold plasma-treated food products.

Plasma chemistry and generation

Plasma is an ionized gas that becomes the fourth state of matter (liquids, solids, gases). Plasma is attained once adequate energy is applied to a gas. Application of energy to a gas will generate an electrical field which will increase the speed of free electrons within the gas. The accelerated electron undergoes contact with neutral gas atoms, leading to incitement or electrolysis. This ionization releases the

surplus amount of free electrons faster causing an avalanche effect of highly reactive chemical species that are capable of inactivating enormous amounts of microorganisms. Cold atmospheric plasma is generated using direct current (DC) or electricity power supply and Alternating current (AC) power supplies. The frequency varies in a wide range from AC power of low (kHz) frequencies through Radio frequencies (MHz) to Microwave (GHz) frequencies which finally release plasma discharges. Different reactors mentioned in Table 1 and Fig 1.

Application of cold plasma in food preservation and processing

Fruits and vegetable sector

Cold plasma technology offers a wider benefit in the food processing industry, decontamination, surface modification and improvised products with desired properties (Warne, 2021). According to Wang *et al.* (2012), *Salmonella* sp. were inactivated up to 90% in carrot slices and 60% in cucumber slices and reported to be less effective in pear slices by about 40% by cold plasma treatment. After one second of treatment, the rate of inactivation on cucumber and pear slices reached above 80%; however, this effect was time

and dose-dependent. Kovacevic *et al.* (2016), studied the effect of cold plasma on anthocyanin and colour in pomegranate juice and chokeberry juice. The results showed an optimistic effect on anthocyanin content and slight changes in pomegranate juice colour; whereas increased level of hydroxycinnamic acids in chokeberry juice. There were increased levels of colour, citric acid and ascorbic acid in prebiotic orange juice treated by the effect of cold plasma (Almeida *et al.*, 2015); these scenarios revealed that cold plasma technology protects the functional compounds (polyphenols). The plasma application on fresh-cut melon probably increases the shelf life and keeps quality (Tappi *et al.*, 2016). The plasma treatment improved the colour and reduced dark area formation in freshly cut kiwifruit. The kiwi fruit-treated samples and control were observed to record same antioxidant content and antioxidant activity (Ramazzina *et al.*, 2015). Misra *et al.* (2014) attempted to increase the cherry tomatoes shelf life by applying cold plasma and concluded that plasma technology could be an effective method of decontamination and retaining the quality of cherry tomatoes (Misra *et al.*, 2014).

Microbial inactivation of aerobic mesophilic bacteria, molds and yeast was observed in treated strawberries where

Table 1: Different reactors used in cold plasma.

Different reactors	Operating principles	References
Dielectric barrier discharge	Dielectric barrier discharge (DBD) is made with two electrodes in-between separated by an insulating dielectric barrier. DBD materials encompass polypropylene, glass and metals like copper, aluminium and brass. It can suppress sparks by regulating the current and operating in a wide range of pressures (104-106 Pa).	Albertos <i>et al.</i> (2017)
Gliding arc	Gliding discharges are low-temperature plasma which is operated by multi-electrode reactors using air pressure. The plasma is produced by two different aluminum electrodes, which comprise a plasma generator along with two copper electrodes. Gas input is given at the top of the nozzle. High voltage up to 9kV or higher is used for plasma generation.	Moreau <i>et al.</i> (2007)
Corona discharge	Corona discharge is a form of electrical discharge produced by the action of fluid ionization with the help of air surrounding a conductor that's electrically charged. The plasma is a self-maintained discharge that occurs at the tip of a pointed electrode where an inconsistent electric field is present.	Scholtz <i>et al.</i> (2015)
Radiofrequency discharge	A radio frequency field that is provided externally creates radio frequency plasma in an extremely fast flow of gas. The plasma is created at the tip of the needle electrode and it then expands to the grounded ring electrode outside the ceramic nozzle. The ceramic nozzle is connected with the above two RF voltage (often 13.56 MHz) electrodes.	Hertwig <i>et al.</i> (2015)
Microwave discharge	Microwaves' electrical discharges are produced with the help of electromagnetic waves at a frequency range of greater than 300 MHz. The plasma is formed by a magnetron cooling system with open, closed and resonance of three structures.	Won <i>et al.</i> (2017)
Plasma jet	A powered electrode creates the plasma, which then flows to another surface. There are four different types of plasma jets viz., DBD jets, single electrode jets, dielectric-free electrode jets and DBD-like jets.	Lu <i>et al.</i> (2012)

Note: RF: radio frequency; AC: Alternating current; DC: Direct current; DBD: Dielectric barrier discharge.

the microbial load got reduced by 2 log₁₀ by treating for 5 minutes in atmospheric cold plasma (Misra *et al.*, 2016). Xu *et al.* (2017) conducted a study on the application of high-voltage atmospheric cold plasma technology in orange juice and reported a positive effect against *Salmonella enterica* serovar *typhimurium* (*S. enterica*). A similar study was conducted by Pankaj and Keener (2018) described that plasma-treated juices had a significant reduction of *S. cerevisiae* and deprived of any change in pH and acidity. Kim *et al.* (2017) studied that cold plasma against spores of *A. brasiliensis*, *B. cereus* and *E. coli* on onion powder for 40 minutes which was known to effectively reduce the number of spores. Min *et al.* (2016) reported that DBD atmospheric cold plasma effectively inactivated *Salmonella* sp. without any change in color or firmness. Cold plasma could be used to inhibit food-borne pathogens and increase the shelf life of fresh lettuce storage life (Song *et al.*, 2015). As a result, cold plasma technology is still widely used in the processing and preservation of fruits and vegetables.

Meat and poultry sector

Microorganism plays an important role in meat decay. Due to their highly perishable nature meat and poultry products are easily attacked by microbial populations and lead to deterioration. Depending upon the way cold plasma therapy, the effects on the microbiology of meat and meat product

differs. Ulbin-Figlewicz *et al.* (2013) investigated the use of cold plasma treatment to inactivate microorganisms and effect on their colour using nitrogen, argon and helium plasma on meat surfaces. The researchers found that helium and argon plasma treatment for 10 minutes effectively reduced the total number of microorganisms (3 log cfu/cm²). The mold and yeast were observed to be 6 log cfu/cm² and 3 log cfu/cm² respectively. Yeast and molds were slightly low in nitrogen plasma and reduction of 1 log cfu/cm² after 10 minutes of exposure. The atmospheric pressure plasma promotes a current innovation in the meat industry for curing process without nitrite additives. At this identical time, the sample received an improved score in organoleptic score and overall acceptability (Lee *et al.*, 2018). Ulbin-Figlewicz and Jarmoluk (2015) proved the consequences of low-pressure plasma on the stability of raw pork samples. After 14 days of meat samples storage, ferric reducing ability reduced from 1.93 to 1.40 mmol Trolox Eq.kg. In numerous studies with *E. coli*, *L. monocytogenes* and *Salmonella* sp., the plasma treatment reduced the counts of pathogenic bacteria and did not influence the textural properties of tested food (Jeyasena *et al.*, 2015; Albertos *et al.*, 2017).

Yong *et al.* (2017) found an alternate method for pork manufacturing, using cold plasma along with sodium




Fig 1: Pictorial representation of types of discharge.

nitrite, there was an increase in quality parameters such as texture, colour and in case of microbial quality, pathogens like *Staphylococcus aureus* and *Bacillus cereus* were significantly reduced. The application of cold plasma using modified air condition (65% O₂: 30% CO₂:5% N₂) study revealed that bacterial isolates were completely inactivated at 120 seconds in ambient air condition, while modified atmospheric condition is needed for complete inactivation of *S. typhimurium* (90 seconds) and *P. fluorescens* (180 seconds), whereas minimum time of 30 seconds for killing *C. jejuni* (Rothrock *et al.*, 2017). Microbial population in the atmosphere cold plasma-treated meat exposed below level was compared with control (Wang *et al.*, 2018) and Zhuang *et al.* (2019) reported that DBD ACP possesses positive consequences on decontamination of meat surface against *Campylobacter*, *Salmonella* and improving meat quality. Finally, numerous researchers reported that there is no significant loss of texture and colour of the meat products when treated with cold plasma.

Grains and flour sectors

According to Misra *et al.* (2015), the use of cold plasma technology improves the viscosity moduli and elastic properties of wheat flour. Plasma technology increased the water uptake and simultaneously it reduced the hardness and chewiness in textural parameters (Thirumdas *et al.*, 2016). The study by Sarangapani *et al.* (2015) reported that low-pressure plasma technology increased the water absorption of rice and reduced the cooking period by up to eight minutes and at the same time, the surface energy was increased. Plasma treatment reduced the amylose content within the amylose to amylopectin ratio and sequentially increased the gelatinization temperature and crystallinity of flour. It also enhanced physicochemical properties like proximate composition and textural properties of black gram with reduced hardness with the range of 22.50 to 23.36 N by cold plasma treatment (Sarangapani *et al.*, 2016). Wu *et al.* (2018) found that corona electrical discharge applied to raw banana starch analysis doesn't showed any significant changes within the amylose content and resistant starch but it reduced the area of diffraction peak and viscosity.

According to Lee *et al.* (2016), after 5 minutes of treatment, cold plasma modestly lowered the pH of brown rice. In germinated brown rice, Chen *et al.* (2016) discovered an increased amount of gamma aminobutyric acid from 19 to 28 mg/100 g. Application of low-temperature plasma technology helped to improve functional properties like flour hydration, antioxidant, hydration properties and gelatinization temperature on basmati rice. The researcher also observed increased water holding capacity and water binding capacity with the rise in plasma and time of treatment (Thirumdas *et al.*, 2016). Chen *et al.* (2012) reported an application of cold plasma technology to decrease the brown rice cooking time, expansion ratio and elongation ratio. Indica brown rice showed a noteworthy amount of decrease in viscosity and breakdown. The application of

low-pressure plasma offered a promising method to decrease the time of cooking on brown rice.

Milk and dairy sector

Milk decay occurs when the pH of the milk got reduced, the pH of milk is lowered as a result of the lactose in the milk being converted to lactic acid. According to Kim *et al.* (2015), utilizing DBD plasma, aerobic bacterial count got decreased along with pH level. Korachi *et al.* (2015), studied the application of cold plasma to milk indicating biochemical alterations in the form of elevated levels of free fatty acids, protein and volatile compounds. Following a 10-minute flexible thin layer DBD plasma treatment, the amount of *E. coli*, *S. typhimurium* and *L. monocytogenes* on slices of cheddar cheese got significantly decreased. The results proved that DBD plasma could probably be an efficient technique to sterilize food products and also this condition should be evaluated for food industry application.

The bacterial count in the entire batch of milk was reduced from 7.78 to 3.63 Log CFU/ml after 20 minutes of plasma treatment (Gurol *et al.*, 2012). To treat the *E. coli*, *L. monocytogenes* and *S. typhimurium* bacteria present in the cheese slices, Yong *et al.* (2015) proposed using atmospheric pressure DBD plasma. The results showed 2.67, 3.10 and 1.65 decimal reductions after plasma treatment. Therefore, developed plasma treatments have a positive effect on improving the protection of cheese products and increase the shelf life of the products. Cold plasma technology has a positive role as an alternative to the chocolate industry. According to Coutinho *et al.* (2019), the physicochemical properties, fatty acid composition, pH and volatile component profiles of chocolate milk drinks revealed that processing had a negative impact on the amount of bioactive and fatty acids present.

Effects of cold plasma in food processing

The behavior of food during the cold plasma treatment and the relationship between the food molecules are of great interest to researchers. Additionally, cold plasma clearly defines the relationship between each food properties physical and chemical indexes and can have a significant impact on how food is processed and preserved.

Cold plasma is effective in water treatment. However, to verify the effects of cold plasma certain analyses must be conducted on the combination of water and beverage components. Hou *et al.* (2019), conducted a study on the effect of cold plasma on blueberry juice. The results emphasized significant increase in phenolic content in juice, which could better preserve the natural colour of blueberry juice.

Misra *et al.* (2018), studied DBD cold plasma on xanthan gum to increase the viscosity and emulsion stabilizing properties. Treated xanthan gum possessed more benefits for salad dressing and instant dry soup formulations with high viscosity at lowering shear rates.

Proteins are predominantly used as a thickening agent, development of dairy products, texturizing agent and

gel formation. Cold plasma penetrates whey protein and isolates structural modification along with foaming quality, functionality and emulsifying properties (Segat *et al.*, 2015). Atmospheric plasma treatment improved oil binding and reduced water binding capacities which enhance the food texture and functionality (Bubler *et al.*, 2015). It provoked the properties within crystallinity and gelatinization of starch. This phenomenon extended surface energy and hydrophilicity and the splitting of the amylose side chain. Plasma treatment persuades similar variations in the functional properties of starch viscosity and the rheological properties of weal wheat flour (Misra *et al.*, 2015).

Pankaj *et al.* (2017), reported that cold plasma treatment increased the glass transition temperature, surface oxygenation and surface roughness of starch films. DBD cold plasma treatment increased the solubility of peanut protein isolates (Ji *et al.*, 2018). Low-pressure plasma increased the hydrophobic properties of whey protein (Terpilowski *et al.*, 2017). Cold plasma treatment positively degrades the mycotoxins (Shi *et al.*, 2017), lowering the pesticide residues (Sarangapani *et al.*, 2016).

CONCLUSION

A non-thermal food processing technique is well known as cold plasma innovations. The uses are enthused with responsive gases to inactivate contaminating microorganisms and antagonistically extend the timeframe of food products. Cold plasma technology can be applied to a wide range of food as they are an effective and flexible antimicrobial process. Since, consumer preference shifted towards healthy and tasty foods that are easily available, ready to eat and easily stored. Cold plasma technology plays a challenging role in food industries for the distribution and mass production of foods without affecting their texture, flavor and colour and it is technically complex and expensive.

Future studies

More research is needed in the field of the beverage industry specifically on sports drinks, naturally fortified drinks *etc.* The effect of cold plasma on such beverages on its organic molecules and its bioactive compounds needs to be extensively studied. One most important research on the conversion of ocean water into drinking water also can be attempted by using cold plasma.

Conflict of interest

All authors declare that they have no conflicts of interest.

REFERENCES

- Albertos, I., Martin-Diana, A., Cullen, P., Tiwari, B., Ojha, S., Bourke, P., Álvarez, C and Rico, D. (2017). Effects of dielectric barrier discharge (DBD) generated plasma on microbial reduction and quality parameters of fresh mackerel (*Scomber scombrus*) fillets. *Innovative Food Science and Emerging Technologies*. 44: 117-122.
- Almeida, F.D.L., Cavalcante, R.S., Cullen, P.J., Frias, J.M., Bourke, P., Fernandes, F.A.N. and Rodrigues, S. (2015). Effects of atmospheric cold plasma and ozone on prebiotic orange juice. *Innovative Food Science and Emerging Technologies*. 32: 127-135.
- Brany, D., Dvorska, D., Halasova, E. and Skovierova, H. (2020). Cold Atmospheric Plasma: A powerful tool for modern medicine. *International Journal of Molecular Sciences*. 21(8): 2932. doi: 10.3390/ijms21082932.
- Bubler, S., Herppich, W.B., Neugart, S., Schreiner, M., Ehlbeck, J., Rohn, S. and Schlüter, O. (2015). Impact of cold atmospheric pressure plasma on physiology and flavonol glycoside profile of peas (*Pisum sativum* 'Salamanca'). *Food Research International*. 76: 132-141. <https://doi.org/10.1016/j.foodres.2015.03.045>.
- Chen, H.H., Chang, H.C., Chen, Y.K., Hung, C.L., Lin, S.Y. and Chen, Y.S. (2016). An improved process for high nutrition of germinated brown rice production: Low-pressure plasma. *Food Chemistry*. 191: 120-127.
- Chen, H.H., Chen, Y.K and Hung C.C., (2012). Evaluation of physicochemical properties of plasma treated brown rice. *Food Chemistry*. 135(1): 74-79.
- Chizoba, F.G., Sun, D.W., Han, Z. and Cheng, J.H. (2017). Microwave-assisted food processing technologies for enhancing product quality and process efficiency: A review of recent developments. *Trends in Food Science and Technology*. 67: 58-69.
- Coutinho, M.N., Marcello R.S., Tatiana C.P., Monica Q.F., Jeremias M., *et al.* (2019). Chocolate milk drink processed by cold plasma technology: physical characteristics, thermal behavior and microstructure. *LWT*. 102: 324-329.
- Gurol, C., Ekinci, F.Y., Aslan, N. and Korachi, M. (2012). Low temperature plasma for decontamination of *E. coli* in milk. *International Journal of Food Microbiology*. 157: 1-5.
- Hertwig, C., Kai, R., Jorg, E., Knorr, D. and O. Schluter. (2015). Decontamination of whole black pepper using different cold atmospheric pressure plasma applications. *Food Control*. 55: 221-229.
- Hou, Y., Wang, R., Gan, Z., Shao, T., Zhang, X., He, M. (2019). Effect of cold plasma on blueberry juice quality. *Food Chemistry*. 290: 79-86. doi: 10.1016/j.foodchem. 2019. 03.12.
- Jayasena, D.D., Kim, H.J., Yong, H.I., Park, S., Kim, K., Choe, W. and Jo, C. (2015). Flexible thin-layer dielectric barrier discharge plasma treatment of pork butt and beef loin: Effects on pathogen inactivation and meat-quality attributes. *Food Microbiology*. 46: 51-57.
- Ji, H., Dong, S., Han, F., Li, Y., Chen, G., Li, L. (2018). Effects of dielectric barrier discharge (DBD) cold plasma treatment on physicochemical and functional properties of peanut protein. *Food Bioprocess Tech*. 11: 344-54. doi: 10.1007/s11947-017-2015-z.
- Kim, H.J., Yong, H.I., Park, S., Kim, K., Choe, W., Jo, C. (2015). Microbial safety and quality attributes of milk following treatment with atmospheric pressure encapsulated dielectric barrier discharge plasma. *Food Control*. 47: 451-56.

- Kim, J.E., Oh, Y.J., Won, M.Y., Lee, K.S and Min, S.C. (2017). Microbial decontamination of onion powder using microwave-powered cold plasma treatments. *Food Microbiology*. 62: 112-123.
- Korachi, M., Ozen, F., Aslan, N., Vannini, L., Guerzoni, M.E., *et al.* (2015). Biochemical changes to milk following treatment by a novel, cold atmospheric plasma system. *International Dairy Journal*. 42: 64-69.
- Kovacevic, D.B., Kljusuric, J.G., Putnik, P., Vukusic, T., Herceg, Z. and Dragovic-Uzelac, V. (2016). Stability of polyphenols in chokeberry juice treated with gas phase plasma. *Food Chemistry*. 212: 323-331.
- Lee, J., Jo, K., Lim, Y., Jeon, H.J., Choe, J.H., Jo, C. and Jung, S. (2018). The use of atmospheric pressure plasma as a curing process for canned ground ham. *Food Chemistry*. 240: 430-436.
- Lee, K.H., Kim, H.J., Woo, K.S., Jo, C., Kim, J.K., Kim, S.H., Park, H.Y., Oh, S.K. and Kim, W.H. (2016). Evaluation of cold plasma treatments for improved microbial and physicochemical qualities of brown rice. *LWT- Food Science and Technology*. 73: 442-447. doi: 10.1016/j.lwt.2016.06.055.
- Lu, X., Laroussi, M. and Puech, V. (2012). On atmospheric-pressure non-equilibrium plasma jets and plasma bullets. *Plasma Sources Science and Technology*. 21(3): 034005. doi: 10.1088/0963-0252/21/3/034005.
- Min, C.S., Si, H.R., Brendam, A.N., Joseph, E.S., Glenn, B. and Alison, L. (2016). Dielectric barrier discharge atmospheric cold plasma inhibits *Escherichia coli* O157:H7, *Salmonella*, *Listeria monocytogenes* and *Tulane virus* in romaine lettuce. *International Journal of Food Microbiology*. 237(21): 114-120.
- Misra, N., Kaur, S., Tiwari, B.K., Kaur, A., Singh, N. and Cullen, P. (2015). Atmospheric pressure cold plasma (ACP) treatment of wheat flour. *Food Hydrocolloids*. 44: 115-121.
- Misra, N., Patil, S., Moiseev, T., Bourke, P., Mosnier, J., Keener, K. and Cullen, P. (2014). In-package atmospheric pressure cold plasma treatment of cherry tomatoes. *Journal of Food Engineering*. 125: 131-138.
- Misra, N.N., Kaur, S., Tiwari, B.K., Kaur, A., Singh, N. and Cullen, P.J. (2015). Atmospheric pressure cold plasma (ACP) treatment of wheat flour. *Food Hydrocolloids*. 44: 115-121.
- Misra, N.N., Pankaj, S.K., Segat, A and Ishikawa, K. (2016). Cold plasma interactions with enzymes in foods and model systems. *Trends Food Science and Technology*. 55: 39-47.
- Misra, N.N., Tiwari, B.K., Raghavarao, K.S.M.S. and Cullen, P.J. (2011). Nonthermal plasma inactivation of food-borne pathogens. *Food Engineering Reviews*. 3(3- 4): 159-170.
- Misra, N.N., Yong, H.I., Phalak, R. and Jo, C. (2018). Atmospheric pressure cold plasma improves viscosifying and emulsion stabilizing properties of xanthan gum. *Food Hydrocolloids*. 82: 29-33.
- Moreau, M., Feuilletoy, M.G., Veron, W., Meylheuc, T., Chevalier, S., Brisset, J.L. and Orange, N. (2007). Gliding arc discharge in the potato pathogen *Erwinia carotovora* subsp. *atroseptica*: Mechanism of lethal action and effect on membrane-associated molecules. *Applied and Environmental Microbiology*. 73(18): 5904-5910. doi: 10.1128/AEM.00662-07.
- Pankaj, S.K. and Keener, K.M. (2018). Cold plasma processing of fruit Juices. *Fruit Juices - extraction, composition, quality and analysis*. Academic Press. 26: 529-537.
- Pankaj, S.K., Wan, Z.J., León, E., Mosher, C., Colonna, W., Keener, K.M. (2017). High-voltage atmospheric cold plasma treatment of different types of starch films. *Starch*. 69: 170009. doi: 10.1002/star.201700009.
- Ramazina, I., Berardinelli, A., Federica, R. (2015). Effect of cold plasma treatment on physic-chemical parametrs and antioxidant activity of minimally processed kiwifruit. *Postharvest Biology and Technology*. 107: 55-65.
- Rothrock, M.J., Zhuang, H., Lawrence, K.C., Bowker, B.C., Gamble, G.R. and Hiett, K.L. (2017). Inpackage inactivation of pathogenic and spoilage bacteria associated with poultry using dielectric barrier discharge-cold plasma treatments. *Current Microbiology*. 74(2): 149-158.
- Sarangapani, C., Devi, Y., Thirumdas, R., Annapure, U.S, Deshmukh, R.R. (2015). Effect of low-pressure plasma on physico-chemical properties of parboiled rice. *Food Science and Technology*. 63: 452-460.
- Sarangapani, C., Thirumdas, R., Devi, Y., Trimukhe, A., Deshmukh, R. and Annapure, U.S. (2016). Effect of low-pressure plasma on physicochemical and functional properties of parboiled rice flour. *Food Science and Technology*. 69: 482-489.
- Scholtz, V., Pazlarova, J., Souskova, H., Khun, J. and Julak, J. (2015). Nonthermal plasma-a tool for decontamination and disinfection. *Biotechnology Advances*. 33(6): 1108-1119. doi: 10.1016/j.biotechadv.2015.01.002.
- Segat, A., Misra, N.N., Cullen, P.J. and Nadia, I. (2015). Atmospheric pressure cold plasma (ACP) treatment of whey protein isolates model solution. *Innovative Food Science and Emerging Technologies*. 29: 247-254.
- Sen, Y., Bakir, H., Basarslan, M.S. and Yegin, M. (2018). Medical application of plasma technology. *Journal of Engineering Research and Applied Sciences*. 7(1): 731-738.
- Shi, H., Ilelej, K., Stroshine, R.L., Keener, K., Jensen, J.L. (2017). Reduction of aflatoxin in corn by high voltage atmospheric cold plasma. *Food Bioprocess Tech*. 10: 1042-1052. doi: 10.1007/s11947-017-1873-8.
- Song, A.Y., Oh, Y.J., Kim, J.E., Song, K.B., Oh, D.H. and Min, S.C. (2015). Cold plasma treatment for microbial safety and preservation of fresh lettuce. *Food Science and Biotechnology*. 24: 1717-1724.
- Tappi, S., Gozzi, G., Vannini, L., Berardinelli, A., Romani, S., Ragni, L. and Rocculi, P. (2016). Cold plasma treatment for fresh-cut melon stabilization. *Innovative Food Science and Emerging Technologies*. 33: 225-233. doi: 10.1016/j.ifset.2015.12.022.
- Terpilowski, K., Tomczynska-Mleko, M., Nishinari, K., Mleko, S. (2017). Surface properties of ion-induced whey protein gels deposited on cold plasma treated support. *Food Hydrocolloid*. 71: 17-25. doi: 10.1016/j.foodhyd.2017.04.032.
- Thirumdas, R., Deshmukh, R.R. and Annapure, U.S. (2016). Effect of low temperature plasma on the functional properties of basmati rice flour. *Journal Food Science Technology*. 53(6): 2742-2751.

- Ulbin-Figlewicz, Jarmoluk, A. and Marycz, K. (2015). Antimicrobial activity of low-pressure plasma treatment against selected foodborne bacteria and meat microbiota. *Annals of Microbiology*. 65: 1537-1546.
- Ulbin-Figlewicz, N., Brychcy, E. and Jarmoluk, A. (2013). Effect of low-pressure cold plasma on surface microflora of meat and quality attributes. *Journal of Food Science and Technology*. 52: 1228-1232.
- Wang, J.M., Zhuang, H., Lawrence, K and Zhang, J.H. (2018). Disinfection of chicken fillets in packages with atmospheric cold plasma: Effects of treatment voltage and time. *Journal of Applied Microbiology*. 124(5): 1212-1219.
- Wang, R., Nian, W.F., Wu, H.Y and Feng, H. (2012). Atmospheric-pressure cold plasma treatment of contaminated fresh fruit and vegetable slices: Inactivation and physiochemical properties evaluation. *The European Physical Journal*. 66(10): 276-278.
- Warne, R.G., Philip, M.W., Hue, Q.P., Nam, N.T., Volker, H., Ian, D.F. (2021). Impact of cold plasma on the biomolecules and organoleptic properties of foods: A review. *Journal of Food Science*. 86(9): 3762-3777.
- Won, M.Y., Lee, S.J., Min, S.C. (2017). Mandarin preservation by microwave-powered cold plasma treatment. *Innov. Food Sci. Emerg. Technol.* 39: 25-32.
- Wu, T.Y., Nan, N.S. and Chi, F.C. (2018). Application of corona electrical discharge plasma on modifying the physicochemical properties of banana starch indigenous to Taiwan. *Analysis*. 26(1): 244-251.
- Xu, L., Garner, A.L., Tao, B. and Keener, K.M. (2017). Microbial inactivation and quality changes in orange juice treated by high voltage atmospheric cold plasma. *Food Bioprocess Technology*. 10: 1-14.
- Yong, H.I., Kim, H.J., Park, S., Alahakoon, A.U., Kim, K., Choe, W. and Jo, C. (2015). Evaluation of pathogen inactivation on sliced cheese induced by encapsulated atmospheric pressure dielectric barrier discharge plasma. *Food Microbiology*. 46: 46-50.
- Yong, H.I., Lee, S.H., Kim, S.Y., Park, S., Park, J., Choe, W. and Jo, C. (2017). Color development, physiochemical properties and microbiological safety of pork jerky processed with atmospheric pressure plasma. *Innovative Food Science and Emerging Technologies*. 53: 78-84.
- Zhuang, H., Rothrock, M., Hiatt, K., Lawrence, K., Gamble, G., Bowker, B. and Keener, K. (2019). In-package antimicrobial treatment of chicken breast meat with high voltage dielectric barrier discharge-electric voltage effect. *Journal of Applied Poultry Research*. 28(4): 801-807.
- Ziuzina, D., Patil, S., Cullen, P.J., Keener, K.M. and Bourke, P. (2014). Atmospheric cold plasma inactivation of *Escherichia coli*, *Salmonella enterica* serovar *Typhimurium* and *Listeria monocytogenes* inoculated on fresh produce. *Food Microbiology*. 42: 109-116.