



Anti-proliferation Activity of Some Plant Extracts against MCF-7 and Hdfn Cell Lines

H.A. Emaduldeen¹, K.J. Alaa², J.S. Dhifaf³, J.M. Dhuha⁴,
M.T.A. Mohammed¹, M.A. Moatasem⁵, H.M.Z. Haniza⁶

10.18805/ajdfr.DRF-395

ABSTRACT

Background: In continuation of the efforts made to find an effective cancer treatment that is relatively safe and with fewer side effects, the current study was developed. Several plants and natural products extracted that are used in folk medicine were used in this study.

Methods: The plant extracts were divided into two groups to evaluate their antiproliferation activity against MCF-7 and Hdfn cell lines by using an MTT assay.

Result: Group A [*Saussurea costus*, *Linum usitatissimum*, *Teucrium polium*, *Artemisia annua*, *Salvia officinalis* and *Berytephusa cunicularis* (crab)] gave the best results by decreasing the viability of MCF-7 cancer cell line to 42.98% using 400 µg/ml from the extract, well show no high effect on normal cells. Also, the IC₅₀ was calculated for both groups, also group A showed the lower IC₅₀ with 65.22 µg/ml. In conclusion, plant extracts and natural products can be effective anticancer drugs with high safety, further studies will establish to confirm the results.

Key words: Antiproliferation activity, Cancer, Medical plants, Natural products.

INTRODUCTION

Cancer continues to be a leading cause of mortality globally, despite the development of numerous treatment approaches as mentioned by Zhang *et al.* (2017); Mishra and Hoque (2021) and Haleem *et al.* (2024). According to Choudhuri *et al.* (2018), cancer is generally defined as a medical disorder marked by uncontrollably growing and reproducing cells as a result of gene accumulating mutations. Conventional ways of treating cancer typically involve radiotherapy, chemotherapy and surgery. However, these treatments' efficiency is diminished by cell resistance to them according to Zhang *et al.* (2017) and Moatasem *et al.* (2023). The number of cancer-related deaths is rising, thus it's important to find effective treatment strategies with fewer side effects and cell toxicity. Natural substances have a variety of health benefits, including the ability to treat cancer. Natural products are made from abundant natural resources, such as plants and are naturally occurring chemicals with biological activity as found by Emaduldeen *et al.* (2022).

According to Hasanpourghadi *et al.* (2017), a large number of medicinal medications in use today are produced from natural substance resources. Natural ingredients have played a significant role in the development of anticancer medications Kalaiselvi *et al.* (2022) and Abutaha *et al.* (2024). Many commonly used anti-cancer medications come from natural sources, including bleomycin from marine sources, actinomycin D and C from bacteria, fencristine, etoposide and paclitaxel from plants. For the foreseeable future, several of these substances which are still the cornerstones of cancer treatment will be essential according to Wall and Wani (1995). In order to continue efforts to find new

¹Environmental Research Center, University of Technology-Iraq, Baghdad, Iraq.

²Department of Research and Development, Ministry of Higher Education and Scientific Research, Baghdad, Iraq.

³Department of Medical Laboratories, College of Applied Medical Sciences, Al Muthanna University, Al Muthanna, Iraq.

⁴Department of Studies and Planning, University of Technology-Iraq, Baghdad, Iraq.

⁵Department of Complementary Medicine, European Board, Iraq.

⁶Department of Biology, Faculty of Science and Mathematics, Sultan Idris Education University, Perak 35900 Malaysia.

Corresponding Author: M.T.A. Mohammed, Environmental Research Center, University of Technology-Iraq, Baghdad, Iraq. Email: mohammed.m.ta@uotechnology.edu.iq

How to cite this article: Emaduldeen, H.A., Alaa, K.J., Dhifaf, J.S., Dhuha, J.M., Mohammed, M.T.A., Moatasem, M.A. and Haniza, H.M.Z. (2026). Anti-proliferation Activity of Some Plant Extracts against MCF-7 and Hdfn Cell Lines. *Asian Journal of Dairy and Food Research*. **45(4)**: 657-661. doi: 10.18805/ajdfr.DRF-395.

Submitted: 28-03-2024 **Accepted:** 14-01-2025 **Online:** 08-03-2025

treatments, our current study includes an influence combination of medical plant extracts known to contain anti-cancer compounds and some natural products. Previous research has proven these plant extracts affect against different types of cancers, these plants are *Saussurea costus* that was proven to have anti-proliferation against breast, liver, colon prostate, lung and gastric cancers as reported by Shati *et al.* (2020); Tian *et al.* (2017); Hung *et al.* (2010) and Ko *et al.* (2005). Also, *Teucrium polium* has shown anticancer activity against glioblastoma, colon,

melanoma, lung and breast cancers according to Nematollahi-Mahani *et al.* (2012); Menichini, *et al.* (2009); and Nematollahi-Mahani *et al.* (2007). Moreover, *Linum usitatissimum* inhibited the growth of prostate, breast, colon and skin cancers reported by Zhou *et al.* (2020); Calado *et al.* (2018); DeLuca *et al.* (2018) and Sharma *et al.* (2014). Furthermore, *Artemisia annua* shown to be an anticancer against Breast, colon and lung cancers these were founded by Jung *et al.* (2021); Ko *et al.* (2020) and Rassias and Weathers (2019). Also, *Zingiber officinale* in previous studies was found to work against leukemia, prostate, breast, skin, ovarian, lung, pancreatic and colorectal cancers as mentioned by Wei *et al.* (2005), Salehi *et al.* (2019) and Lee *et al.* (2008). *Curcuma longa* has anticancer against prostate, colorectal, head and neck, breast, brain, glioblastoma and pancreatic cancers these results proved by Dorai *et al.* (2001); Mudduluru *et al.* (2011) and Chakravarti *et al.* (2006). *Allium sativum* has also been shown to anticancer on pancreas, lung, breast, prostate, colon, stomach, cervical and liver cancers these mentioned by Gore *et al.* (2021) and Chhabria *et al.* (2015). Additionally, the crab extract was also used in the current study which previously showed antiproliferation against breast and lung cancers reported by Tsukada *et al.* (1990) and Rezakhani *et al.* (2014). Also, Propolis poses anticancer activity against leukemia, breast, pancreas, cervical and oral cancers according to Aso *et al.* (2004), Vatansever *et al.* (2010) and Lapidot *et al.* (2002). For these reasons and others these plants were chosen to evaluate their antiproliferation effects on the cancer cell lines as groups not individual.

MATERIALS AND METHODS

Plants extraction

Groups A and B of plants were used in the current investigation. *Saussurea costus*, *Teucrium polium*, *Artemisia annua*, *Salvia officinalis*, *Linum usitatissimum* and *Berytephusa cunicularis* (crab) are among the plants in Group A. *Salvia officinalis*, *Artemisia annua*, *Teucrium polium*, *Linum usitatissimum*, *Curcuma longa*, *Ceratonia silique*, *Zingiber officinale* and *Propolis* were among the plants in group B. All plants and natural materials were purchased in dry, pure circumstances from Baghdad, Iraq and local markets. Each crude plant weighed fifty grams, which were combined with distilled water in a 1:4 ratio (crud plant: water) and allowed to boil for fifteen minutes. According to Handa *et al.* (2008), the concentrated extracts were filtered and allowed to evaporate completely at room temperature. Following that, the dried extract was weighed and seven working concentrations were made.

This study evaluated the anticancer activity of the plant mixture extract using primary dermal fibroblast normal (HdFn) and human breast adenocarcinoma cancer (MCF-7) cell lines. The cells were grown in accordance with Freshney (2015) in 75 cm² tissue culture flasks with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin (100 U/mL penicillin and 100 µg/mL streptomycin) in a

humidified 5% CO₂ environment at 37°C in Eagle's Minimum Essential Medium EMEM (ATCC, USA).

Cytotoxicity of plants Group A and B on MCF-7 and HdFn cell lines was assessed using the MTT assay, in which the cell lines from groups A and B were treated for 48 hours at the following concentrations: 400, 200, 100, 50, 25, 12.5 and 6.25 µg/ml. After that, they were incubated with 10 µL of MTT at 37°C for two hours. Following the incubation period, each well was filled with 100 µL of DMSO after the media within the cells was aspirated. At 570 nm, the absorbance was measured with a microtiter plate reader.

Statistical analysis

One-way analysis of variance (ANOVA) was used to evaluate all the data, which were reported as the mean plus or minus three replicates' standard deviation. In addition, the half-maximal inhibitory concentration (IC₅₀) values were computed using the regression equation and the best-fit approach. P values were also computed for every group.

RESULTS AND DISCUSSION

In the current study, the anticancer effects of a number of plants that people use as traditional medicine were assessed collectively against the MCF-7 cancer cell line by comparing it with the effect against the normal HdFn cell line at 48 hours. These plants were gathered in two groups A and B. Group A showed a high antiproliferation effect against MCF-7 cells by using high concentrations (400, 200 and 100 µg/ml) from the extract and the highest was 400 µg/ml where the viability dropped to 42.978% (Table 1). While the control cell has 76.50% viability at the same concentration as shown in Table (1). Also, IC₅₀ for group A was calculated and found to be 65.22 µg/ml against the MCF-7 cell line and 338.6 µg/ml on the HdFn cell line. These data showed a promising result where group A extracts were effective on cancer cell lines well it safe on normal cells. The P value was (<0.0001).

Furthermore, group B showed an antiproliferation effect against MCF-7 cells by using a high concentration of 400 µg/ml from the extract where the viability decreased to 62.886%. While the control cell has 74.26% viability at the same concentration as illustrated in Table (2). Also, IC₅₀ for group B was calculated and found to be high concentration against the MCF-7 cell line which was 315.1 µg/ml and

Table 1: Results of Group A against MCF-7 and HdFn cell line.

Concentration µg/ml	Mean viability (%) ± SD	
	MCF-7	HdFn
400	42.98±4.46	76.50±4.84
200	51.66±2.9	87.92±3.57
100	59.68±1.54	92.21±2.7
50	73.95±1.72	96.48±1.2
25	86.72±1.30	96.95±1.1
12.5	94.71±2.54	94.2±2.97
6.25	94.83±1.9	94.25±1.62

Table 2: Results of Group B against MCF-7 and Hdfn cell line.

Concentration $\mu\text{g mL}^{-1}$	Mean viability (%) $\pm$ SD	
	MCF-7	Hdfn
400	62.89 $\pm$ 3.7	74.26 $\pm$ 2.7
200	79.44 $\pm$ 1.5	88.27 $\pm$ 2.6
100	90.39 $\pm$ 1.4	93.6 $\pm$ 2.1
50	93.82 $\pm$ 1.29	94.17 $\pm$ 1.5
25	94.02 $\pm$ 2.63	95.22 $\pm$ 0.82
12.5	95.60 $\pm$ 2.02	95.56 $\pm$ 1.38
6.25	96.29 $\pm$ 0.80	95.29 $\pm$ 1.05

804.2 $\mu\text{g/ml}$ against the Hdfn cell line. Although IC_{50} of group B on cancer cells is lower than on normal cells it is considered high. The P value was (<0.0001).

The medical plants that are used in the current study have phytochemical compounds that possess anticancer effects such as the *Saussurea costus* which contain Sesquiterpene lactones (SLs) according to Lin *et al.* (2016). Furthermore, *Teucrium polium* is indicated to contain mono and sesquiterpenes, flavonoids, phenolic, saponins terpenoids, alkaloids and terpenoids as reported by Sharifi-Rad *et al.* (2022). Also, Garros *et al.* (2018) found that the *Linum usitatissimum* used in the current study has phenolic acids, flavonoids and lignin. *Artemisia annua* is known to contain Artemisinin, coumarins, flavones, flavonols and phenolic acids, sesquiterpenes illustrated by Das (2012). Additionally, *Zingiber officinale* has Sesquiterpene and gingerols as An *et al.* (2016) mentioned. The well-studied *Curcuma longa* has many phytochemicals like tannins, alkaloids, saponins, flavonoids, terpenoids, cardiac glycosides and phenolic acids (curcumin) according to Singh and Madan, (2019). Also, *Allium sativum* is known to contain alliin, allicin, ajoenes, vinylthiins and flavonoids as Slusarenko *et al.* (2008) found. By influencing the proteins, enzymes and signaling pathways involved in the genesis or progression of malignancies, these phytochemicals-many of which are thought to be pharmaceutically active-have been shown to have potent anticancer properties according to Tariq *et al.* (2017). Phytochemicals are found in trace amounts in the plant extract so using individual plant extracts as anti-cancers requires a huge amount of the same plant, well using multiple plant extracts together provides the required concentrations of the active substance as reported by Lewis and Tollefsbol (2017). So, in the present study, the combination of these plants provided high concentrations of phytochemicals which resulted in duplicated effects against cancer cell lines. Furthermore, a combination of multi-plant extracts or natural products gained more attention in recent years to develop new anticancer drugs or enhance the activity of the exciting ones. In addition, the synergistic effect of plant extracts will show up when multi-plant extracts are used together, which refers to the enhanced or increased effect that is observed when different plant extracts are combined. Many studies have

investigated the synergistic effect of plant extracts on various properties, such as antioxidant potential, antimicrobial activity and antifungal activity. The previous study by Saba and colleagues (2017) found that the mixtures of plant extracts exposed higher levels of phenolics, flavonoids and antioxidant activity compared to individual plants which showed more activity than separate plant extracts. For example, the PHY906, Chinese herb medicine contains four plant extracts (*Scutellaria baicalensis* Georgi, *Glycyrrhiza uralensis* Fisch., *Paeonia lactiflora* Pall. and *Ziziphus jujube* Mill), is a well-known traditional medicine that used for 1800 years. This drug reported to enhance the activity of chemotherapy against gastrointestinal cancer via several strategies according to Lam *et al.* (2010), also its work combined with capecitabine to enhance the antitumor activity and decrease the toxicity of capecitabine in HCC (hepatocellular carcinoma) mentioned by Changou *et al.* (2021). There is a hypothesis that suggests the mixture of plant extracts may attack several targets such as receptors, enzymes, signal pathways, etc. and all of these result in one goal which destroying the cancer cells as Imming *et al.* (2006) reported. Furthermore, the combination of drugs or natural products may affect the bioavailability in the target site according to Butterweck *et al.* (2000). Furthermore, multidrug therapy in which two or more agents were mixed is a well-known strategy for killing cancer cells and acting the human body immunity or repair mechanisms reported by Doos *et al.* (2014).

CONCLUSION

Among the results we obtained in our current study is that the combination of plant and natural product extracts in group A gave the best result in the antiproliferation of cancer cells where it reduced the viability at most of the concentrations studied. It also did not affect the normal control cells. These finding prof the anticancer effect of group A extract as well as safety against normal cells. Further studies will be established to evaluate the antioxidant and antiapoptosis ability of the studied extracts on different cancer cell lines. Also, *in vivo* study will be studied and thus, we will try to find the exact mechanism through which these extracts gave their results.

ACKNOWLEDGEMENT

The authors express their gratitude to the following institutions and organizations for their invaluable support and scientific assistance: Agriculture College, Al Muthanna University, Environment Research Center, University of Technology- Iraq, Biology Department, Faculty of Science and Mathematics, Sultan Idris Education University.

Ethical statement

The university of technology in Iraq oversaw the study's completion.

Conflict of interest

There is no conflict of interest.

REFERENCES

- Abutaha, N.M.T. Alghamdi, R. and Al-Waddan, M. (2024). Induction of apoptosis and ROS production in liver cancer cells by saponin fraction from *Alcea rosea* L. Seeds. *Indian Journal of Animal Research*. **58(10)**: 1765-1771.
- An, K., Zhao, D., Wang, Z., Wu, J., Xu, Y., Xiao, G. (2016). Comparison of different drying methods on chinese ginger (*Zingiber officinale* Roscoe): Changes in volatiles, chemical profile, antioxidant properties and microstructure. *Food Chemistry*. **197**: 1292-1300.
- Aso, K., Kanno, S.I., Tadano, T., Satoh, S. and Ishikawa, M. (2004). Inhibitory effect of propolis on the growth of human leukemia U937. *Biological and Pharmaceutical Bulletin*. **27(5)**: 727-730.
- Butterweck, V., Jürgenliemk, G., Nahrstedt, A. and Winterhoff, H. (2000). Flavonoids from *Hypericum perforatum* show antidepressant activity in the forced swimming test. *Planta Medica*. **66(01)**: 3-6.
- Calado, A., Neves, P.M., Santos, T. and Ravasco, P. (2018). The effect of flaxseed in breast cancer: A literature review. *Frontiers in Nutrition*. **5**: 4.
- Chakravarti, N., Myers, J.N. and Aggarwal, B.B. (2006). Targeting constitutive and interleukin 6 inducible signal transducers and activators of transcription 3 pathway in head and neck squamous cell carcinoma cells by curcumin (diferuloyl methane). *International Journal of Cancer*. **119(6)**: 1268-1275.
- Changou, C.A., Shiah, H.S., Chen, L.T., Liu, S., Luh, F., Liu, S.H., Cheng, Y.C. and Yen, Y. (2021). A phase II clinical trial on the combination therapy of PHY906 plus capecitabine in hepatocellular carcinoma. *The Oncologist*. **26(3)**: e367-e373.
- Chhabria, S.V., Akbarsha, M.A., Li, A.P., Kharkar, P.S. and Desai, K.B. (2015). *In situ* alliin generation using targeted alliinase delivery for inhibition of MIA PaCa-2 cells via epigenetic changes, oxidative stress and cyclin-dependent kinase inhibitor (CDKI) expression. *Apoptosis*. **20(10)**: 1388-1409.
- Choudhuri, S., Chanderbhan, R. and Mattia, A. (2018). Chapter 20-carcinogenesis: Mechanisms and models. *Veterinary Toxicology*. **3**: 354.
- Das, S. (2012). *Artemisia annua* (Qinghao): A pharmacological review. *Int. J. Pharm. Sci. Res.* **3(12)**: 4573-4577.
- DeLuca, J.A. Garcia-Villatoro, E.L. and Allred, C.D. (2018). Flaxseed bioactive compounds and colorectal cancer prevention. *Current Oncology Reports*. **20(8)**: 1-8.
- Doos, L., Roberts, E.O., Corp, N. and Kadam, U.T. (2014). Multi-drug therapy in chronic condition multimorbidity: A systematic review. *Family Practice*. **31(6)**: 654-663.
- Dorai, T., Cao, Y.C., Dorai, B., Buttyan, R. and Katz, A.E. (2001). Therapeutic potential of curcumin in human prostate cancer. III. Curcumin inhibits proliferation, induces apoptosis and inhibits angiogenesis of LNCaP prostate cancer cells *in vivo*. *Prostate*. **47(4)**: 293-303.
- Emaduldeen, H.A., Shamran, D.J., Mahdi, D.J. and Mohammed, M.T.A. (2022). Risk factors of cancer: A review. *Science Archives*. **3(4)**: 284-288 <https://doi.org/10.47587/SA.2022.3407>.
- Freshney, R.I. (2015). *Culture of Animal Cells: A manual of Basic Technique and Specialized Applications*. John Wiley and Sons.
- Garros, L., Drouet, S., Corbin, C., Decourtil, C., Fidel, T., Lebas de Lacour, J. and Hano, C. (2018). Insight into the influence of cultivar type, cultivation year and site on the lignans and related phenolic profiles and the health-promoting antioxidant potential of flax (*Linum usitatissimum* L.) seeds. *Molecules*. **23(10)**: 2636.
- Gore, G.G., Satish, S., Ganpule, A., Srivastava, S. and Athavale, M. (2021). Garlic (*Allium sativum*) exhibits anticancer and anticancer stem cell activity on breast, prostate, colon, hepatic and cervical cancer cell lines. *Int. J. Herbal Med.* **9**: 93-99.
- Haleem, A.M., Taha, M.M. and Ayoub, A.A. (2024). Anti-tumor and antioxidant effects of *Ganoderma lucidum* extracts on oral squamous cell carcinoma and skin squamous cell carcinoma *in vitro*. *Current Issues in Pharmacy and Medical Sciences, Sciencdo*. **37(2)**: 79-84. <https://doi.org/10.2478/cipms-2024-0013>.
- Handa, S.S., Khanuja, S.P.S., Longo, G. and Rakesh, D.D. (2008). *Extraction Technologies for Medicinal and Aromatic Plants*, (1st edn), no. 66. Italy: United Nations Industrial Development Organization and the International Centre for Science and High Technology. pp. 747-752.
- Hasanpourghadi, M., Yeng Looi, C., Kumar Pandurangan, A., Sethi, G., Fen Wong, W. and Rais Mustafa, M. (2017). Phytometabolites targeting the Warburg effect in cancer cells: A mechanistic review. *Current Drug Targets*. **18(9)**: 1086-1094.
- Hung, J.Y., Hsu, Y.L., Ni, W.C., Tsai, Y.M., Yang, C.J., Kuo, P.L. and Huang, M.S. (2010). Oxidative and endoplasmic reticulum stress signaling are involved in dehydrocostuslactone-mediated apoptosis in human non-small cell lung cancer cells. *Lung Cancer*. **68(3)**: 355-365.
- Imming, P., Sinning, C. and Meyer, A. (2006). Drugs, their targets and the nature and number of drug targets. *Nature Reviews Drug Discovery*. **5(10)**: 821-834.
- Jung, E.J., Paramanathan, A., Kim, H.J., Shin, S.C., Kim, G.S., Jung, J.M., Ryu, C.H., Hong, S.C., Chung, K.H., Kim, C.W. and Lee, W.S. (2021). *Artemisia annua* L. Polyphenol-induced cell death is ROS-independently enhanced by inhibition of JNK in HCT116 colorectal cancer cells. *International Journal of Molecular Sciences*. **22(3)**: 1366.
- Kalaiselvi, L., Sriram, P., Gokulakannan, R., Parthiban, M., Kannan, T.A., Preetha, S.P. and Pazhanivel, N. (2022). Chemopreventive effect of *stoechospermum marginatum* against experimental colon carcinogenesis induced by 1,2- Dimethyl Hydrazine. *Indian Journal of Animal Research*. **56(4)**: 460-467. doi: 10.18805/IJAR.B-4495
- Ko, S.G., Kim, H.P., Jin, D.H., Bae, H.S., Kim, S.H., Park, C.H. and Lee, J.W. (2005). *Saussurea lappa* induces G2-growth arrest and apoptosis in AGS gastric cancer cells. *Cancer Letters*. **220(1)**: 11-19.
- Ko, Y.S., Jung, E.J., Go, S.I., Jeong, B.K., Kim, G.S., Jung, J.M., Hong, S.C., Kim, C.W., Kim, H.J. and Lee, W.S. (2020). Polyphenols extracted from *Artemisia annua* L. exhibit anti-cancer effects on radio-resistant MDA-MB-231 human breast cancer cells by suppressing stem cell phenotype, β -catenin and MMP-9. *Molecules (Basel, Switzerland)*. **25(8)**: 1916.

- Lam, W., Bussom, S., Guan, F., Jiang, Z., Zhang, W., Gullen, E.A. and Cheng, Y.C. (2010). The four-herb chinese medicine PHY906 reduces chemotherapy-induced gastrointestinal toxicity. *Science Translational Medicine*. **2(45)**: 45ra59.
- Lapidot, T., Walker, M.D. and Kanner, J. (2002). Antioxidant and prooxidant effects of phenolics on pancreatic β -cells *in vitro*. *Journal of Agricultural and Food Chemistry*. **50(25)**: 7220-7225.
- Lee, C.S., Kim, Y.J., Lee, M.S., Han, E.S. and Lee, S.J. (2008). 18 β -Glycyrrhetic acid induces apoptotic cell death in SiHa cells and exhibits a synergistic effect against antibiotic anti-cancer drug toxicity. *Life Sciences*. **83(13-14)**: 481-489.
- Lewis, K.A. and Tollefsbol, T.O. (2017). The influence of an epigenetics diet on the cancer epigenome. *Epigenomics*. **9(9)**: 1153-1155.
- Lin, X., Peng, Z., Fu, X., Liu, C., Xu, Y., Ji, W., Fan, J., Chen, L., Fang, L., Huang, Y. and Su, C. (2016). Volatile oil from *Saussurea lappa* exerts antitumor efficacy by inhibiting epithelial growth factor receptor tyrosine kinase-mediated signaling pathway in hepatocellular carcinoma. *Oncotarget*. **7(48)**: 79761-79773.
- Menichini, F., Conforti, F., Rigano, D., Formisano, C., Piozzi, F. and Senatore, F. (2009). Phytochemical composition, anti-inflammatory and antitumor activities of four *Teucrium* essential oils from Greece. *Food Chemistry*. **115(2)**: 679-686.
- Mishra, M. and Hoque, M. (2021). Advancement in cancer immuno therapy in veterinary medicine: A review. *Indian Journal of Animal Research*. **55(9)**: 993-998. doi: 10.18805/ijar.B-4141.
- Moatasem, M.A., Dhifaf, J.S., Emaduldeen, H.A., Mohammed, M.T.A. Ibrahim, S.A., Dhuha, J.M., Haniza, H.M.Z., Ulrazaq, D.J.A. Falah, H.I. and Rana, R.K. (2023). The synergistic anticancer effect of some plant extracts in combination against human liver cancer cell line. *Asian Journal of Dairy and Food Research*. **42(4)**: 529-534. doi: 10.18805/ajdfr.DRF-297.
- Mudduluru, G., George-William, J.N., Muppala, S., Asangani, I.A., Kumarswamy, R., Nelson, L.D. and Allgayer, H. (2011). Curcumin regulates miR-21 expression and inhibits invasion and metastasis in colorectal cancer. *Bioscience Reports*. **31(3)**: 185-197.
- Nematollahi-Mahani, S.N., Mahdinia, Z., Eftekhavaghefi, R., Mehrabani, M., Hemayatkhah-Jahromi, V. and Nabipour, F. (2012). *In vitro* inhibition of the growth of glioblastoma by *teucrium polium* crude extract and fractions. *International Journal of Phytomedicine*. **4(4)**: 582.
- Nematollahi-Mahani, S.N., Rezazadeh-Kermani, M., Mehrabani, M. and Nakhaee, N. (2007). Cytotoxic effects of *teucrium polium* on some established cell lines. *Pharmaceutical Biology*. **45(4)**: 295-298.
- Rassias, D.J. and Weathers, P.J. (2019). Dried leaf *Artemisia annua* efficacy against non-small cell lung cancer. *Phytomedicine: International Journal of Phytotherapy and Phytopharmacology*. **52**: 247-253.
- Rezakhani, L., Rashidi, Z., Mirzapour, P. and Khazaei, M. (2014). Anti proliferatory effects of crab shell extract on breast cancer cell line (MCF7). *Journal of Breast Cancer*. **17(3)**: 219-225.
- Saba, A., Nazish, J., Khalil-ur, R., Fatiqa, Z.M. and Yasin, A. (2017). Synergistic interactions of polyphenols and their effect on antiradical potential. *Pakistan Journal of Pharmaceutical Sciences*. **30(4)**: 1297-1304.
- Salehi, B., Fokou, P.V.T., Yamthe, L.R.T., Tali, B.T., Adetunji, C.O., Rahavian, A. and Sharifi-Rad, J. (2019). Phytochemicals in prostate cancer: From bioactive molecules to upcoming therapeutic agents. *Nutrients*. **11(7)**: 1483.
- Sharifi-Rad, M., Pohl, P., Epifano, F., Zengin, G., Jaradat, N. and Messaoudi, M. (2022). *Teucrium polium* (L.): Phytochemical screening and biological activities at different phenological stages. *Molecules*. **27(5)**: 1561.
- Sharma, J., Singh, R. and Goyal, P. (2014). Chemopreventive role of flaxseed oil against chemical induced skin cancer in mammals. *J. Cancer Biol. Treat.* **1(001)**. doi: 10.24966/CBT-7546/100001.
- Shati, A.A., Alkahtani, M.A., Alfaifi, M.Y., Elbehairi, S., Elsaid, F.G., Prasanna, R. and Mir, M.A. (2020). Secondary metabolites of *saussurea costus* leaf extract induce apoptosis in breast, liver and colon cancer cells by Caspase-3- dependent intrinsic pathway. *Bio. Med. Research International*. pp. 1608942.
- Singh, I. and Madan, V.K. (2019). Effect of moisture levels on various phytoconstituents of turmeric (*Curcuma longa* L.). *Journal of Pharmacognosy and Phytochemistry*. **8(1)**: 1427-1432.
- Slusarenko, A.J., Patel, A. and Portz, D. (2008). Control of plant diseases by natural products: Allicin from garlic as a case study. *European Journal of Plant Pathology*. pp. 313-322.
- Tariq, A., Sadia, S., Pan, K., Ullah, I., Mussarat, S., Sun, F. and Adnan, M. (2017). A systematic review on ethnomedicines of anti cancer plants. *Phytotherapy Research*. **31(2)**: 202-264.
- Tian, X., Song, H.S., Cho, Y.M., Park, B., Song, Y.J., Jang, S. and Kang, S.C. (2017). Anticancer effect of *Saussurea lappa* extract via dual control of apoptosis and autophagy in prostate cancer cells. *Medicine*. **96(30)**: e7606.
- Tsukada, K., Matsumoto, T., Aizawa, K., Tokoro, A., Naruse, R.S., Suzuki, S. and Suzuki, M. (1990). Antimetastatic and growth inhibitory effects of N acetylchitohexaoase in mice bearing Lewis lung carcinoma. *Japanese Journal of Cancer Research*. **81(3)**: 259-265.
- Vatansever, H.S., Sorkun, K., Gurhan, S.I.D., Ozdal-Kurt, F., Turkoz, E., Gencay, O. and Salih, B. (2010). Propolis from Turkey induces apoptosis through activating caspases in human breast carcinoma cell lines. *Acta Histochemical*. **112(6)**: 546-556.
- Wall, M.E. and Wani, M.C. (1995). Camptothecin and taxol: Discovery to clinic-thirteenth Bruce F. Cain Memorial Award Lecture. *Cancer Research*. **55(4)**: 753-760.
- Wei, Q.Y., Ma, J.P., Cai, Y.J., Yang, L. and Liu, Z.L. (2005). Cytotoxic and apoptotic activities of diarylheptanoids and gingerol-related compounds from the rhizome of Chinese ginger. *Journal of Ethnopharmacology*. **102(2)**: 177-184.
- Zhang, C.L., Huang, T., Wu, B.L., He, W.X. and Liu, D. (2017). Stem cells in cancer therapy: opportunities and challenges. *Oncotarget*. **8(43)**: 75756.
- Zhou, X., Huang, N., Chen, W., Xiaoling, T., Mahdavi, B., Raoofi, A. and Atabati, H. (2020). HPLC phenolic profile and induction of apoptosis by *Linum usitatissimum* extract in LNCaP cells by caspase 3 and Bax pathways. *Amb. Express*. **10(1)**: 1-11.