



Hepatoprotective Mechanism of Apigenin *via* Suppression of Oxidative Inflammatory Signaling and Apoptosis against Hepatotoxicity Induced by CCl₄ in Rats

Hebah Alabbad¹, Manal Alfwuaires¹, Ashraf M. Abdel-Moneim², Hany Elsaywy^{3,4},
Nourah Almulhim³, Ademola C. Famurewa^{5,6}, Azza Sedky^{1,2}

10.18805/IJAR.BF-1609

ABSTRACT

Background: Carbon tetrachloride (CCl₄) is a critical hepatotoxicant causing liver injury and fibrosis via hepatic production of reactive oxygen species (ROS). Apigenin (APG) is a natural bioactive compound and flavonoid antioxidant. We, therefore, evaluated whether APG could mitigate CCl₄-mediated hepatotoxicity.

Methods: Rats were randomly divided and administered APG and/or CCl₄ in Control group, CCl₄ group, APG + CCl₄ groups (APG: 10 and 20 mg/kg bw) and APG groups (APG: 10 and 20 mg/kg bw) 2 times per week for 7 consecutive weeks.

Result: Rats exposed to CCl₄ demonstrated marked increases in serum alanine aminotransferase (ALT), aspartate aminotransferase (AST) and monoamine oxidase (MAO) activities and decreased hepatic malondialdehyde (MDA) level compared to control. The hepatic activities of superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPx) decreased appreciably. The CCl₄ intoxication caused significant increases in inflammatory cytokines (IL-6 and TNF-α) and apoptosis markers, while the anti-inflammatory cytokines (IL-4 and IL-10) decreased with evident histopathological lesions compared to control. APG-dose-dependently prevented these hepatic alterations.

Key words: Apigenin, Carbon tetrachloride, Hepatotoxicity.

INTRODUCTION

Environmental toxicants are ubiquitous and have been implicated in the pathophysiology of chronic diseases and organ damage (Famurewa *et al.*, 2022). Carbon tetrachloride is among these environmental pollutants that inadvertently enter the human body and trigger pathologies. It is an exogenous industrial solvent with a strong affinity for the liver and it has been recognized as a hepatotoxicant (Zhou *et al.*, 2020). In spite of its hepatotoxic and nephrotoxic stress in humans and experimental animals, it is still being used in dry cleaning, fumigation of grains, insecticide and filling fire extinguishers (Que *et al.*, 2022). The hepatic metabolism of CCl₄ *via* the action of cytochrome P450-dependent monooxygenases results in the production of its hepatic metabolites, trichloromethyl (CCl₃) and trichloromethyl peroxy (•OCCl₃) reactive oxygen species (ROS) (El-Hadary and Hassanien, 2016). The metabolites are potent free radicals that trigger subsequent reactive oxygen species (ROS) generation. Hepatic diseases, including hepatic cirrhosis and fibrosis have been associated with ROS effect (El-Hadary and Hassanien, 2016). The prevailing mechanism of CCl₄-induced hepatotoxic damage is consistent with the potential of its oxidative metabolites to cause hepatic antioxidant impairment leading to oxidative stress and pro-inflammatory activation (Yang *et al.*, 2018). Oxidative stress is capable of initiating membrane lipid peroxidation and, finally cell death (Zhou *et al.*, 2020). The trichloromethyl radical can react with sulfhydryl groups of glutathione

¹Department of Biological Sciences, College of Science, King Faisal University, Al-Ahsa, 31982, Saudi Arabia.

²Department of Zoology, Faculty of Science, Alexandria University, Alexandria, Egypt.

³Department of Chemistry, College of Science, King Faisal University, PO Box 400 Al-Ahsa, 31982, Saudi Arabia.

⁴Department of Chemistry, Faculty of Science, Tanta University, Tanta, Egypt.

⁵Department of Medical Biochemistry, Faculty of Basic Medical Sciences, College of Medical Sciences, Alex Ekwueme Federal University, Ndufu-Alike, Ikwo, Ebonyi State, Nigeria.

⁶Department of Pharmacology, Manipal College of Pharmaceutical Sciences, Manipal Academy of Higher Education, Manipal-576 104, Karnataka, India.

Corresponding Author: Hany Elsaywy, Department of Chemistry, College of Science, King Faisal University, PO Box 400 Al-Ahsa, 31982, Saudi Arabia. Email: hmostafa@kfu.edu.sa

How to cite this article: Alabbad, H., Alfwuaires, M., Abdel-Moneim, A.M., Elsaywy, H., Almulhim, N., Famurewa, A.C. and Sedky, A. (2026). Hepatoprotective Mechanism of Apigenin *via* Suppression of Oxidative Inflammatory Signaling and Apoptosis against Hepatotoxicity Induced by CCl₄ in Rats. *Indian Journal of Animal Research*. 60(9): 1612-1617. doi: 10.18805/IJAR.BF-1609.

Submitted: 16-11-2022 **Accepted:** 02-01-2023 **Online:** 10-02-2023

enzymes and other protein thiols to cause deficit in glutathione metabolism and consequently reduces cell activities of superoxide dismutase and catalase (Almatroodi

et al., 2020). Existing literature has implicated the involvement of oxidative pro-inflammation and apoptosis in CCl₄-induced hepatic pathologies Yue *et al.* (2020).

A robust body of literature reports that natural antioxidants are efficacious in preventing oxidative stress-related liver pathologies (Almatroodi *et al.*, 2020). Apigenin (4,5,7-trihydroxyflavone) is a natural polyphenolic flavone widely found in fruits, herbs and vegetables (Shankar *et al.*, 2017). Systematic investigations on the health benefits of APG have shown its several pharmacological effects, including antioxidant, anti-inflammatory, antidiabetic, anti-Alzheimer's disease, anticancer, antiviral and antihypertensive (DeRango-Adem and Blay, 2021 Ahmad *et al.*, 2019 and Wu *et al.*, 2021). Therefore, the study herein was designed to explore APG's possible hepatoprotective effect and mechanism against CCl₄-induced hepatotoxicity in rats.

MATERIALS AND METHODS

Chemicals

Carbon tetrachloride (Cat. No. 56-23-5) was purchased from Loba Chemie (India), olive oil (Cat. No. EL 40-105) was purchased from AGROVIM (Greece) and apigenin (Cat. No. 520-36-5) was purchased from Matrix Scientific (Columbia, SC, USA). Reagent kits for liver function markers (ALT: E-BC-K235-S and AST: E-BC-K236-M). The kits for SOD (SOD: Cat. No. SD 2521), CAT (CAT: Cat. No. CA 2517) and GPx (GPx: Cat. No. GP 2524) activities and MDA (MDA: Cat. No. MD 2529) level were obtained from BioDiagnostics, Giza, Egypt. The MAO kit (EMAO-100) was purchased from BioAssay Systems, CA, USA. The kits for cytokines were procured from OriGene Technologies Inc., Rockville, MD and MyBioSource, Inc., San Diego, USA, while ELISA kits for apoptosis markers were obtained from PEVIVA, USA.

Experimental animals

Thirty male rats (weighing 200-220 g) were used which were obtained from the Faculty of Science at King Faisal University, Kingdom of Saudi Arabia. The experimental design used in this study was approved by the Department of Chemistry Research and Ethics Committee, College of Science, King Faisal University, Kingdom of Saudi Arabia, with reference number KFUE-REC/2020-09-02. The rats were housed in a laboratory animal room under standard management conditions of a temperature of 20-25°C and the exposure time to light per day was 12 hours.

Experimental design

Following 14 days of acclimatization, rats were randomly divided into six groups (n=5/group).

Group I (Control): Rats received an intraperitoneal (i.p.) injection of olive oil (3 ml/kg b.w.).

Group II (CCl₄): Rats received CCl₄ (30% in olive oil) (3 ml/kg b.w., i.p) [20].

Group III (APG + CCl₄): Rats received APG (10 mg/kg, orally) + CCl₄ (3 ml/kg b.w., i.p).

Group IV (APG + CCl₄): Rats received APG (20 mg/kg, orally) + CCl₄ (3 ml/kg bw, ip).

Group V (APG): Rats received APG (10 mg/kg b.w, orally).

Group VI (APG): Rats received APG (20 mg/kg b.w, orally).

The treatment was performed twice per week for seven consecutive weeks. The doses of CCl₄ and APG doses were chosen according to Liu *et al.* (2018) and Anusha *et al.* (2017), respectively.

Blood samples and liver tissues were collected at the end of the experimental period from all experimental animals. The serum was separated and liver samples were taken for the biochemical analysis and histological examination.

Biochemical analysis

Determination of liver function indices

Liver enzymes, including AST and ALT, were quantitatively estimated in serum using commercial kits, following the manufacturer's instructions. Monoamine oxidase (MAO) was analyzed using a commercial kit.

Determination of oxidative stress markers

Liver antioxidant CAT, SOD, GPx and TBARS as MDA levels were measured using standard assay kits, following the procedures of the kits' manufacturer.

Determination of inflammatory markers

The inflammatory cytokines interleukin-6 (IL-6), interleukin-4 (IL-4) and interleukin-10 (IL-10) levels were measured in serum using rat ELISA kits. The tumor necrosis factor- α (TNF- α) was analyzed using ELISA kit. The analyses were done according to the manufacturers' instructions.

Apoptotic markers

According to the protocols described in the ELISA kits.

Histopathological analysis

Liver samples from each group were fixed in 4% formaldehyde for 24 h, dehydrated in ascending ethanol series, embedded in paraffin, sectioning (4 μ m thick) and stained with hematoxylin and eosin dye (H and E) and examined under light microscope. The histopathological alterations in tissue sections were scored and an average value was determined as follows: normal histostructure (0), mild (1), moderate (2) and severe (3) following extensive alterations according to their histopathological findings (Bancroft and Gamble 2002).

Statistical analysis

The SPSS software was used for data analysis and results presented as mean $\pm$ SEM. The one-way analysis of variance (ANOVA) was used to determine statistical differences among groups, followed by a post hoc "LSD test: the least significant difference." A p<0.05 was considered statistically significant.

RESULTS AND DISCUSSION

Effect of APG on liver function indices

ALT and AST are enzymes mainly confined within the hepatocytes, although they are also found in the heart, kidney, blood cells and pancreas (Aja *et al.*, 2020) while monoamine oxidase is a mitochondrial enzyme with high

activity in the brain, GIT and hepatic tissue (Jaka *et al.*, 2021). CCl₄ significantly increased the serum activities of ALT, AST and MAO compared to normal control. However, the administration of APG prevented these toxic effects in a dose-dependent manner reduction (Table 1). Similar result was obtained by Liu *et al.* (2018) and Almatroodi *et al.* (2020). Elevations in activities of ALT and AST is due to cellular leakage and loss of functional integrity of hepatic cell membrane, whereas elevated ALP activity is a marker of hepatic-cholestatic damage while increased hepatic MAO activity reveals mitochondrial injury (Abou Seif, 2016). Furthermore, hepatic necrosis which was observed in our histopathological analysis has been implicated as a contributory factor to ALT and AST release into the blood circulation. Contrarily, concomitant administration of APG (10 and 20 mg/kg body weight) to rats inhibited CCl₄-mediated hepatic damage dose-dependently in this study. This result agree with that study of Yue *et al.* (2020).

Effect of APG on oxidative stress markers

It is known that SOD, CAT and GPx are cellular antioxidant enzymes that scavenge ROS and thus promote antioxidant mechanism. CCl₄ significantly reduced the activities of SOD, CAT and GPx and increased MDA level.

However, the administration of APG (at 10 and 20 mg/kg body weight) significantly increased the activities of these enzymes and decreased MDA level compared to CCl₄ group. APG exerted significant dose-dependent reduction on MDA alone (Table 2). These findings can be confirm by previous reports (Ubhenin *et al.*, 2016). The significant depression in the hepatic activities of SOD, CAT and GPx implies the overwhelming oxidative imbalance exerted by the CCl₄ metabolites leading to antioxidant imbalance and/or oxidative stress in the rat liver exposed to CCl₄.

The chief mechanism underlying CCl₄ hepatotoxicity is oxidative stress arising from the hepatic metabolites of CCl₄ (trichloromethyl and trichloromethyl peroxy radicals). These hepatic metabolites, are ROS generators and consumers of antioxidant balance (El-Hadary and Hassanien, 2016). Intriguingly, trichloromethyl can react with sulfhydryl groups present in glutathione, GPx and protein thiols to form an oxidative complex, exerting deleterious effects on SOD and CAT (El-Hadary and Hassanien, 2016).

Interestingly, the APG administration scavenged the ROS and enhanced hepatic antioxidant homeostasis. This was evident through prominently elevated hepatic activities of SOD, CAT and GPx and decreased level of MDA compared to CCl₄ group, in consonance with earlier studies (Raskovic *et al.*, 2017 and Ubhenin *et al.*, 2016). Consequently, appreciably indicating ability of APG to inhibit oxidative stress.

It was noteworthy to observe that the two doses of APG failed to demonstrate dose-dependent increases in SOD, CAT and GPx activities but only in MDA level (Table 2). A robust body of literature indicates APG antioxidant property (DeRango-Adem and Blay, 2021). It is a natural polyphenolic flavonoid flavone with ROS-scavenging activity and other pharmacological properties (Salehi *et al.*, 2014).

By implication, therefore, APG demonstrates a hepatoprotective effect against CCl₄ oxidative stress via its antioxidant property. The free hydroxyl groups present on the A/B rings of APG are responsible for the antioxidant effects of this flavone (Singh *et al.*, 2014).

Effect of APG on inflammatory markers in CCl₄-intoxicated rats

In CCl₄ group, the levels of IL-6 and TNF-α significantly increased while the levels of anti-inflammatory markers, IL-4 and IL-10 is significantly reduced compared to normal control. This result indicates that CCl₄ provokes pro-inflammation and depresses anti-inflammation (Yang *et al.*, 2018). The oxidative stress observed herein might have enhanced the induction of cytokine expression. Oxidative stress status may trigger the nuclear translocation of nuclear factor-kappa B (NF-κB) to stimulate the expression of cytokine proteins Edeogu *et al.* (2020), which has been reported to occur during CCl₄ hepatotoxicity (Tsai *et al.*, 2017).

However, APG administration prominently abrogated the effect of CCl₄ on these cytokines. Interestingly, the two

Table 1: Effect of APG on liver function indices (U/L) in CCl₄-intoxicated rats.

	ALT	AST	MAO
Control	37.8±1.1	55.6±1.1	21.6±0.8
CCl ₄	86.8±1.8 ^a	245.6±1.5 ^a	59.8±0.9 ^a
APG 10 + CCl ₄	62.4±0.5 ^b	195±10.1 ^b	39.6±1.3 ^b
APG 20 + CCl ₄	46.8±1.2 ^{bc}	134.6±1.4 ^{bc}	27.8±0.9 ^{bc}
APG 10	37.1±0.5	56.6±0.9	21.2±1.1
APG 20	37.2±0.9	54.6±1.3	21.4±0.7

Data were displayed as mean±SEM (n = 5 rats/group). CCl₄: Carbon tetrachloride; APG: Apigenin (where 10 and 20 refer to the dose in mg/kg bw); ^ap<0.05: Significant when compared to control group in the same column. ^bp<0.05: Significant when compared to CCl₄ group in the same column. ^cp<0.05: Significant when compared to APG 10 + CCl₄ group in the same column.

Table 2: Effect of APG on liver oxidative stress markers in CCl₄-intoxicated rats.

	CAT (U/mg protein)	SOD (mU/mg protein)	GP _x (mU/mg protein)	MDA (nmol/gm tissue)
Control	34.6±1.5	55.4±1.7	55.6±1.1	5.0±0.2
CCl ₄	17.6±1.3 ^a	19.8±0.9 ^a	25.6±1.1 ^a	25.4±1.2 ^a
APG 10+CCl ₄	27.4±1.0 ^b	31.8±1.4 ^b	35.4±1.2 ^b	18.4±1.2 ^b
APG 20+CCl ₄	31.4±0.6 ^b	36.2±1.2 ^b	43.8±1.8 ^b	13.4±0.9 ^{bc}
APG 10	35.5±1.3	54.2±1.5	55.4±1.2	5.6±0.1
APG 20	33.4±1.3	53.6±0.8	54.8±1.4	5.4±0.1

Data are displayed as mean±SEM (n = 5 rats/group). CCl₄: Carbon tetrachloride; APG: Apigenin (where 10 and 20 refer to the dose in mg/kg bw); ^ap<0.05: Significant when compared to control group in the same column. ^bp<0.05: Significant when compared to CCl₄ group in the same column. ^cp<0.05: Significant when compared to APG 10 + CCl₄ group in the same column.

Table 3: Effect of APG on inflammatory markers (pg/ml) in CCl₄ intoxicated rats.

	Pro-inflammatory		Anti-inflammatory	
	IL-6	TNF- α	IL-4	IL-10
Control	37.2 $\pm$ 1.2	51.4 $\pm$ 1.7	38.4 $\pm$ 1.4	52.4 $\pm$ 1.2
CCl ₄	364.6 $\pm$ 1.3 ^a	392 $\pm$ 1.0 ^a	26.4 $\pm$ 1.1 ^a	24.4 $\pm$ 1.2 ^a
APG 10 + CCl ₄	222.4 $\pm$ 5.2 ^b	220.8 $\pm$ 5.2 ^b	30.2 $\pm$ 2.6 ^b	35.7 $\pm$ 0.9 ^b
APG 20 + CCl ₄	134.4 $\pm$ 1.1 ^{bc}	115 $\pm$ 1.5 ^{bc}	34.6 $\pm$ 0.9 ^b	43.4 $\pm$ 1.2 ^b
APG 10	36.6 $\pm$ 0.9	52.2 $\pm$ 1.9	37.2 $\pm$ 0.6	51.4 $\pm$ 1.0
APG 20	36 $\pm$ 1.5	51.2 $\pm$ 1.2	39.8 $\pm$ 0.9	52.2 $\pm$ 1.3

Data were displayed as mean $\pm$ SEM (n = 5 rats/group). CCl₄: Carbon tetrachloride; APG: Apigenin (where 10 and 20 refer to the dose in mg/kg bw); ^ap<0.05: Significant when compared to control group in the same column. ^bp<0.05: Significant when compared to CCl₄ group in the same column. ^cp<0.05: Significant when compared to APG 10 + CCl₄ group in the same column.

doses of APG expressed a dose-dependent effect on IL-6 and TNF- α only (Table 3). Accumulating number of studies demonstrate that APG can suppresses inflammatory cascades (Salehi *et al.*, 2019).

Effect of APG on apoptosis markers in CCl₄-intoxicated rats

Serum levels of M30 and M65 are indicator of apoptosis of cells undergoing necrosis and cell death de Haas *et al.* (2008). The release of TNF- α promotes cell apoptosis leading to hepatocyte cell death (Li *et al.*, 2020). The CCl₄ significantly increased the level of M65 and M30 in comparison to the normal control (Fig 1). The induction of apoptosis by CCl₄ in this study corroborate the findings of previous studies (Li *et al.*, 2020).

On the contrary, the APG doses significantly reduced the levels of M65 and M30 compared to CCl₄ group (Fig 1).




Fig 1: Effect of APG on apoptosis markers in CCl₄-intoxicated rats. CCl₄: Carbon tetrachloride; APG: Apigenin (where 10 and 20 refer to the dose in mg/kg bw); *p<0.05: significant when compared to control group. #p<0.05: significant when compared to CCl₄ group. @p<0.05: significant when compared to APG 10 + CCl₄ group.




Fig 2: Photomicrograph representation of the effect of APG on liver histology of CCl₄-exposed rats (H and E stain). Control group (1a); CCl₄ group (2a), APG + CCl₄ group (3a and 4a) and APG group (5a and 6a). Control showed normal hepatocytes (H), blood sinusoids (S) and central vein (CV). The liver from CCl₄ group showed infiltration of inflammatory cells (star), severe hepatic necrosis (arrow) and cytoplasmic degeneration. The APG + CCl₄ groups revealed ameliorated structures showing mildly congested central vein (CV), cytoplasmic degeneration (thick arrow), Kupffer cells (K) and Kupffer cellular infiltration (star). APG group showed normal structures consistent with normal hepatocytes (H), Kupffer cells (K), central vein (CV) and blood sinusoid (S). Values are expressed as mean $\pm$ SEM (n=5).

*Significant when compared to control (p<0.05); #significant when compared to CCl₄ group.

Interestingly, the two doses of APG revealed a dose-dependent effect on M65 and M30, respectively.

Also, there was dose-dependent antiapoptotic effects of APG as mentioned in previous studies (Mohamed *et al.*, 2020 and Zhong *et al.*, 2017).

Histopathological analysis

Fig 2 showed that, the control group (1a), the liver architecture appears normal with hepatocytes, blood sinusoids and central vein. On the contrary, the liver histological analysis from CCl₄ group revealed inflammatory cells infiltration (star), cytoplasmic vacuolation and degeneration and severe hepatic necrosis (arrow) (2a). The oxidative milieu created by CCl₄ may be the cause of these histopathologic lesions. The observed infiltration of inflammatory cells in our histopathological analysis could also account for the increased IL-6 and TNF- α levels in this study (Yeh *et al.*, 2013).

The administration of APG in the APG + CCl₄ groups ameliorated the CCl₄-induced alterations to mild lesions (3a and 4a). The APG only did not alter the liver structures (5a and 6a).

CONCLUSION

The present study demonstrated the hepatotoxic effect of CCl₄ and emphasized that APG possesses a mechanistic hepatoprotective effect against CCl₄ induced hepatotoxicity via abrogation of oxidative stress, pro-inflammation and apoptosis. Chiefly, these beneficial effects can be attributed to antioxidant and anti-inflammatory activities of APG.

Funding information

Deanship of Scientific Research, King Faisal University, Saudi Arabia, with Grant Number: GRANT 2214.

ACKNOWLEDGEMENT

The authors acknowledge the Deanship of Scientific Research, Vice Presidency for Graduate Studies and Scientific Research at King Faisal University, Saudi Arabia for financial support under the Student Researcher Funding track [GRANT 2214].

Conflict of interest

The authors declare no conflict of interests.

REFERENCES

- Abou Seif, H.S. (2016). Physiological changes due to hepatotoxicity and the protective role of some medicinal plants. *Benisuef University Journal of Basic and Applied Sciences*. **5(2)**: 134-146.
- Ahmad, A., Kumari, P. and Ahmad, M. (2019). Apigenin attenuates edifenphos-induced toxicity by modulating ROS-mediated oxidative stress, mitochondrial dysfunction and caspase signal pathway in rat liver and kidney. *Pesticide Biochemistry and Physiology*. **159**: 163-172.
- Aja, P.M., Ekpono, E.U., Awoke, J.N., Famurewa, A.C., Izekwe, F.I., Okoro, E.J. Okorie, C.F. *et al.* (2020). Hesperidin ameliorates hepatic dysfunction and dyslipidemia in male Wistar rats exposed to cadmium chloride. *Toxicology Reports*. **7**: 1331-1338.
- Almatroodi, S.A., Anwar, S., Almatroudi, A., Khan, A.A., Alrumaihi, F., Alsahli, M.A. and Rahmani, A.H. (2020). Hepatoprotective effects of garlic extract against carbon tetrachloride (CCl₄)-induced liver injury via modulation of antioxidant, anti-inflammatory activities and hepatocyte architecture. *Applied Sciences*. **10(18)**: 6200. <https://doi.org/10.3390/app10186200>.
- Anusha, C., Sumathi, T. and Joseph, L.D. (2017). Protective role of apigenin on rotenone induced rat model of Parkinson's disease: Suppression of neuroinflammation and oxidative stress mediated apoptosis. *Chemicobiological Interactions*. **269**: 67-79.
- Bancroft, D. and Gamble, M. (2002). *The Theory and Practice of Histology Technique*. 5th ed. Churchill Living Stone.
- de Haas, E.C., di Pietro, A., Simpson, K.L., Meijer, C., Suurmeijer, A.J., Lancashire, L.J., Cummings, J., de Jong, S., de Vries, E.G., Dive, C. and Gietema, J.A. (2008). Clinical evaluation of M30 and M65 ELISA cell death assays as circulating biomarkers in a drug-sensitive tumor, testicular cancer. *Neoplasia*. **10(10)**: 1041-1048.
- DeRango-Adem, E.F. and Blay, J. (2021). Does oral apigenin have real potential for a therapeutic effect in the context of human gastrointestinal and other cancers? *Frontiers in Pharmacology*. **12**: 681477.
- Edeogu, C.O., Kalu, M.E., Famurewa, A.C., Asogwa, N.T., Onyeji, G.N. and Ikpeomo, K.O. (2020). Nephroprotective effect of *Moringa oleifera* seed oil on gentamicin-induced nephrotoxicity in rats: Biochemical evaluation of antioxidant, anti-inflammatory and antiapoptotic pathways. *Journal of the American College of Nutrition*. **39(4)**: 307-315.
- El-Hadary, A.E. and Ramadan, H.M.F. (2016). Hepatoprotective effect of cold-pressed *Syzygium aromaticum* oil against carbon tetrachloride (CCl₄)-induced hepatotoxicity in rats. *Pharmaceutical Biology*. **54(8)**: 1364-1372.
- El-Kashef, D.H. and Zaghloul, R.A. (2022). Ameliorative effect of montelukast against carbon tetrachloride-induced hepatotoxicity: Targeting NLRP3 inflammasome pathway. *Life Sciences*. **304**: 120707.
- Famurewa, A.C., Renu, K., Eladi, M.A., Chakraborty, R., Myakala, H., El-Sherbiny, M., Elsherbini, D.M.A. *et al.* (2022). Hesperidin and hesperetin against heavy metal toxicity: Insight on the molecular mechanism of mitigation. *Biomedicine and Pharmacotherapy*. **149**: 112914.
- Jaka, O., Iturria, I., van der Toorn, M., Hurtado de Mendoza, J., Latino, D., Alzualde, A., Peitsch, M. C., Hoeng, J., Koshibu, K. (2021). Effects of natural monoamine oxidase inhibitors on anxiety-like behavior in zebrafish. *Frontiers in Pharmacology*. **13(12)**: 669370.
- Li, N., Li, B., Zhang, J., Liu, X., Liu, J., Li, K., Pan, T., Wang, S. and Diao, Y. (2020). Protective effect of phenolic acids from chebulae fructus immaturus on carbon tetrachloride induced acute liver injury via suppressing oxidative stress, inflammation and apoptosis in mouse. *Natural Product Research*. **34(22)**: 3249-3252.
- Liu, Y., Liu, Q., Hesketh, J., Huang, D., Gan, F., Hao, S., Tang, S., Guo, Y. and Huang, K. (2018). Protective effects of selenium-glutathione-enriched probiotics on CCl₄-induced liver fibrosis. *The Journal of Nutritional Biochemistry*. **58**: 138-149.

- Mohamed, W.R., Kotb, A.S., Abd El-Raouf, O.M. and Mohammad, F.E. (2020). Apigenin alleviated acetaminophen-induced hepatotoxicity in low protein-fed rats: Targeting oxidative stress, STAT3 and apoptosis signals. *Journal of Biochemical and Molecular Toxicology*. **34(5)**: e22472.
- Que, R., Cao, M., Dai, Y., Zhou, Y., Chen, Y. and Lin, L. (2022). Decursin ameliorates carbon-tetrachloride-induced liver fibrosis by facilitating ferroptosis of hepatic stellate cells. *Biochemistry and Cell Biology*. **100(5)**: 378-386.
- Rašković, A., Gigov, S., Ćapo, I., Paut Kusturica, M., Milijašević, B., Kojić-Damjanov, S. and Martić, N. (2017). Antioxidative and protective actions of apigenin in a paracetamol-induced hepatotoxicity rat model. *European Journal of Drug Metabolism and Pharmacokinetics*. **42(5)**: 849-856.
- Salehi, B., Venditti, A., Sharifi-Rad, M., Kręgiel, D., Sharifi-Rad, J., Durazzo, A., Lucarini, M. et al. (2019). The therapeutic potential of apigenin. *International Journal of Molecular Sciences*. **20(6)**: 1305. doi: 10.3390/ijms20061305.
- Shankar, E., Goel, A., Gupta, K. and Gupta, S. (2017). Plant flavone apigenin: An emerging anticancer agent. *Current Pharmacology Reports*. **3(6)**: 423-446.
- Singh, M., Kaur, M. and Silakari, O. (2014). Flavones: An important scaffold for medicinal chemistry. *European Journal of Medicinal Chemistry*. **84**: 206-239.
- Tsai, J.C., Chiu, C.S., Chen, Y.C., Lee, M.S., Hao, X.Y., Hsieh, M.T., Kao, C.P. and Peng, W.H. (2017). Hepatoprotective effect of *Coreopsis tinctoria* flowers against carbon tetrachloride-induced liver damage in mice. *BMC Complementary and Alternative Medicine*. **17(1)**: 139. doi: 10.1186/s12906-017-1604-8.
- Ubhenin, A., Igbe, I., Adamude, F. and Falodun, A. (2016). Hepatoprotective effects of ethanol extract of *Caesalpinia bonduca* against carbon tetrachloride induced hepatotoxicity in Albino Rats. *Journal of Applied Sciences and Environmental Management*. **20(2)**: 396-401.
- Wu, Q., Li, W., Zhao, J., Sun, W., Yang, Q., Chen, C., Xia, P., Zhu, J., Zhou, Y., Huang, G., Yong, C., Zheng, M., Zhou, E. and Gao, K. (2021). Apigenin ameliorates doxorubicin-induced renal injury via inhibition of oxidative stress and inflammation. *Biomedicine and Pharmacotherapy*. **137**: 111308.
- Yang, C., Li, L., Ma, Z., Zhong, Y., Pang, W., Xiong, M. and Fang, S., Li, Y. (2018). Hepatoprotective effect of methyl ferulic acid against carbon tetrachloride-induced acute liver injury in rats. *Experimental and Therapeutic Medicine*. **15(3)**: 2228-2238.
- Yeh, Y.H., Hsieh, Y.L. and Lee, Y.T. (2013). Effects of yam peel extract against carbon tetrachloride-induced hepatotoxicity in rats. *Journal of Agricultural and Food Chemistry*. **61(30)**: 7387-7396.
- Yue, S., Xue, N., Li, H., Huang, B., Chen, Z. and Wang, X. (2020). Hepatoprotective effect of apigenin against liver injury via the non-canonical NF-κB pathway *in vivo* and *in vitro*. *Inflammation*. **43(5)**: 1634-1648.
- Zhong, Y., Jin, C., Gan, J., Wang, X., Shi, Z., Xia, X. and Peng, X. (2017). Apigenin attenuates patulin-induced apoptosis in HEK293 cells by modulating ROS-mediated mitochondrial dysfunction and caspase signal pathway. *Toxicon*. **137**: 106-113.
- Zhou, J., Zhang, Y., Li, S., Zhou, Q., Lu, Y., Shi, J., Liu, J., Wu, Q. and Zhou, S. (2020). *Dendrobium nobile* Lindl. alkaloids-mediated protection against CCl₄-induced liver mitochondrial oxidative damage is dependent on the activation of Nrf2 signaling pathway. *Biomedicine and Pharmacotherapy*. **129**: 110351.