



# Impact of Abiotic Factors and Dates of Sowing on *Helicoverpa armigera* (Hübner) Parasitoid *Campoletis chloridae* (Uchida) under Chickpea Cropping System in Tarai Region of Uttarakhand

S.D. Divija, Meena Agnihotri<sup>1</sup>

10.18805/LR-4500

## ABSTRACT

**Background:** The solitary endo-larval parasitoid *Campoletis chloridae* Uchida (Hymenoptera: Ichneumonidae) is an important biocontrol agent of *H. armigera* in chickpea ecosystem. Abiotic factors are likely to affect the insect host and activity of its parasitoid. Therefore, the present study was conducted to understand the impact of weather factors and different dates of sowing on per cent parasitization of *Helicoverpa armigera* (Hübner) by *Campoletis chloridae* (Uchida) in chickpea ecosystem.

**Methods:** In this field-laboratory investigation were carried out at Pantnagar during the cropping season of 2017-18. Twenty-five larvae of *H. armigera* (first and second instars) were collected randomly at weekly basis from untreated chickpea crop. In the laboratory, the collected larvae were reared till parasitoid emergence and observed for per cent parasitization.

**Result:** The observation recorded revealed that mean seasonal parasitism by *Campoletis* on larvae of *H. armigera* was found to be 21%, 43.5% and 34.5% in early (20<sup>th</sup> October), normal (11<sup>th</sup> November) and late sown crop (1<sup>st</sup> December), respectively. Various abiotic parameters like maximum and minimum temperature, maximum and minimum relative humidity registered high level of significant association with per cent parasitization in early and normal sown crop compared to the low level of association in late sown crop. The multiple regression analysis showed that the combined effect of all the meteorological parameters were responsible for 90%, 57.9% and 30.5% variation in per cent parasitization in early, normal and late sown crop, respectively.

**Key words:** *Campoletis chloridae*, *Helicoverpa armigera*, Parasitization, Weather.

## INTRODUCTION

Chickpea, *Cicer arietinum* is an important *rabi* pulse grown all over the world. India is the single largest producer of chickpea accounting for 65 per cent of the total production (FAOSTAT, 2017). Insect pests are the major limiting factors in chickpea production. It is known to be attacked by about 60 insect-pests (Reed *et al.*, 1987), which causes 15% yield loss annually. Among these Chickpea pod borer (CPB), *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) is the most notorious insect pest. Yield losses due pod borer damage in chickpea may range from 70 to 95% (Prakash *et al.*, 2007). The increasing concern for environmental problems that are created by the use of toxic pesticide, development of high-level resistance in *Helicoverpa* to the conventional insecticides (Kranthi *et al.*, 2002) has promoted a worldwide interest in the adoption of ecological basis of pest management. Biological control is one important tool of Integrated Pest Management that utilizes the contribution of natural enemies to reduce pest population. The ichneumonid, endo-larval parasitoid *Campoletis chloridae* (Uchida) recorded to be the most effective natural enemy of, *Helicoverpa armigera* on chickpea. *Campoletis chloridae* (Uchida) was found to parasitize the early instars of *Helicoverpa armigera*. The parasitoid deposit eggs singly into first or second instar larvae and comes out from the host during third or fourth instar to spin a cocoon and pupation takes place on the plant inside the spun cocoon. In chickpea ecosystem this endo-larval parasitoid reported

Department of Entomology, College of Agriculture, G.B. Pant University of Agriculture and Technology, Pantnagar-263 145, Uttarakhand, India.

<sup>1</sup>University of Agricultural Sciences, Bengaluru-560 065, Karnataka, India.

**Corresponding Author:** S.D. Divija, Department of Entomology, College of Agriculture, G.B. Pant University of Agriculture and Technology, Pantnagar-263 145, Uttarakhand, India.

Email: sd03ento@gmail.com

**How to cite this article:** Divija, S.D. and Agnihotri, M. (2021). Impact of Abiotic Factors and Dates of Sowing on *Helicoverpa armigera* (Hübner) Parasitoid *Campoletis chloridae* (Uchida) under Chickpea Cropping System in Tarai Region of Uttarakhand. Legume Research. DOI: 10.18805/LR-4500.

**Submitted:** 03-09-2020 **Accepted:** 04-01-2021 **Online:** 26-03-2021

to cause 0.98 to 68.50 per cent parasitization of *H. armigera* (Kaur *et al.*, 2000). Date of sowing has a great impact on the activity of pod borer as well as *Campoletis* due to environmental variation. Therefore, the present study was conducted to observe the effect of chick pea sowing time on natural parasitization of gram pod borer by *Campoletis*.

## MATERIALS AND METHODS

The present study was carried out in the Norman E. Borlaug crop research centre (NEB-CRC), G.B.P.U.A and T,

Pantnagar to study the impact of abiotic factors and the different dates of sowing on Percent parasitization of *Helicoverpa armigera* by *Campoletis chloridae* in chickpea ecosystem. The chickpea variety PG 186 was sown on three dates (20<sup>th</sup> October, 11<sup>th</sup> November and 1<sup>st</sup> December) during 2017-18 in an area of 10m<sup>2</sup>. Twenty-five larvae of *H. armigera* (first and second instars) were collected randomly at weekly basis from untreated chickpea crop. The collected Larvae were brought to the laboratory in plastic vials and reared individually in laboratory till the parasitoid emergence. Observations were recorded on number of adult parasitoid emerging from *H. armigera* larvae and per cent parasitization was computed. Simple correlation coefficient was derived between the per cent parasitization during a standard week and average maximum temperature, minimum temperature, maximum relative humidity and minimum relative humidity. To find out the individual as well as combined impact of the abiotic factors on the per cent parasitization was worked out using regression analysis. The statistical analysis was done using SPSS statistical tool.

## RESULTS AND DISCUSSION

### Per cent parasitization of *H. armigera* in three different dates sown chickpea

In the present study the effect of different dates of sowing on seasonal parasitization of *Campoletis chloridae*, a larval parasitoid on *Helicoverpa* was studied in chickpea ecosystem during 2017-18 and the results have been presented in Table 1. The first appearance of parasitoid was observed on 51<sup>st</sup> standard meteorological week (S.W.) in all the three different dates sown crop. It was observed that


mean seasonal parasitism by *Campoletis* on larvae of *H. armigera* was found to be 21, 43.5 and 34.5 per cent in early (20<sup>th</sup> October), normal (11<sup>th</sup> November) and late sown crop (1<sup>st</sup> December), respectively. Higher parasitism of 80 per cent was occurred on 3<sup>rd</sup> S.W. in normal sown crop as compared to the 48 per cent (2<sup>nd</sup> S.W.) and 72 per cent (6<sup>th</sup> S.W.) in early and late sown crop, respectively (Fig 1). The present results were in agreement with Bisane *et al.* (2008) who also observed the incidence of *C. chloridae* from 51<sup>st</sup> S.W. (17-23 December) until 3<sup>rd</sup> S.W. (15-21 January) and it reported to cause 8.11 to 11.54 per cent parasitization of *Helicoverpa*. Ojha *et al.* (2017) recorded the maximum natural parasitization of *H. armigera* by *C. chloridae* as high as 51.67 and 56.67 per cent during 2010-11 and 2011-12, respectively.

### Influence of abiotic factors on percent parasitization in *H. armigera*

The relationships between per cent parasitization and various abiotic factors were worked out through correlation and regression analysis. The perusal of the results reveals that the abiotic parameters mostly registered high level of significant association with the per cent parasitization data in early and normal sown crop compared to the low level of non-significant association in late sown crop (Table 2). In early and normal sown crop, it is observed that per cent parasitization by *Campoletis* had a highly significant negative correlation with maximum and minimum temperature compared to non-significant association in late sown crop. Whereas highly significant positive correlation was recorded between maximum and minimum relative humidity in early sown crop compared normal sown crop which recorded

**Table 1:** Effect of different sowing dates on per cent parasitization during 2017-18.

Date of observation	Early sown (20.10.17)			Normal sown (11.11.17)		Late sown (1.12.17)	
	No. of <i>H. armigera</i> larvae collected	No. of larvae parasitized	Per cent Parasitization (%)	No. of larvae parasitized	Per cent Parasitization (%)	No. of larvae parasitized	Per cent Parasitization (%)
51 <sup>st</sup> S.W	25	6	24	6	24	1	4
52 <sup>nd</sup> S.W	25	5	20	12	48	3	12
1 <sup>st</sup> S.W	25	9	36	10	40	5	20
2 <sup>nd</sup> S.W	25	12	48	11	44	7	28
3 <sup>rd</sup> S.W	25	9	36	20	80	11	44
4 <sup>th</sup> S.W	25	7	28	11	44	15	60
5 <sup>th</sup> S.W	25	8	32	15	60	17	68
6 <sup>th</sup> S.W	25	6	24	19	76	18	72
7 <sup>th</sup> S.W	25	7	28	18	72	15	60
8 <sup>th</sup> S.W	25	4	16	14	56	16	64
9 <sup>th</sup> S.W	25	6	24	12	48	7	28
10 <sup>th</sup> S.W	25	3	12	9	36	8	32
11 <sup>th</sup> S.W	25	1	4	8	32	6	24
12 <sup>th</sup> S.W	25	0	0	3	12	2	8
13 <sup>th</sup> S.W	25	1	4	4	16	3	12
14 <sup>th</sup> S.W	25	0	0	2	8	4	16
Mean			21		43.5		34.5



**Fig 1:** Effect of different sowing dates on per cent parasitization during 2017-18.

**Table 2:** Correlation coefficient (r) between per cent parasitization and abiotic factors.

Variable		Temperature °C		R. Humidity %	
		Max.	Min.	Max.	Min.
Per cent parasitization	Early sown	-.940**	-.855**	.831**	.797**
	Normal sown	-.517*	-.708**	.667**	0.278 <sup>NS</sup>
	Late sown	-0.294 <sup>NS</sup>	-0.401 <sup>NS</sup>	0.386 <sup>NS</sup>	0.067 <sup>NS</sup>

\*\* Significant at 1%, \* Significant at 5%, NS = Non-significant.

**Table 3:** Multiple regression equations of per cent parasitization and abiotic factors.

Date of sowing	Regression equation	Coefficient of Determination (R <sup>2</sup> )	'F' estimate
20 <sup>th</sup> October	Y = 100.457-2.856 (TMAX)+0.366 (TMIN)+0.079 (RHM)-0.349 (RHE)	0.9	0.001
11 <sup>th</sup> November	Y = -26.509 +1.256 (TMAX)-4.734 (TMIN)+1.067 (RHM)-0.218 (RHE)	0.579	0.035
01 <sup>st</sup> December	Y = 344.626-5.584 (TMAX)+0.065 (TMIN)-0.567 (RHM)-2.145 (RHE)	0.305	0.363

highly significant positive correlation in maximum relative humidity with per cent parasitization. Non-significant positive correlation was observed between minimum relative humidity in normal sown crop, maximum and minimum relative humidity in late sown crop with per cent parasitization. The result was in cogent evidence with Pillai *et al.* (2016) who observed the significant negative correlation between weather parameters and the population of the larval parasitoid with maximum temperature and minimum temperature. Whereas positive correlation was recorded with maximum relative humidity and minimum relative humidity during 2010-11 and 2011-12, respectively.

The multiple regression analysis of the per cent parasitization data with the abiotic factors for the three different dates sown crop reveals that the combined effect of all the meteorological parameters was statistically significant in early and normal sown crop as compared to non-significant up to 95% probability level in late sown crop as evident from the estimated values of 'F' (Table 3). Together all the weather factors were responsible for 90%, 57.9% and 30.5% variation in per cent parasitization in early, normal and late sown crop, respectively.

## CONCLUSION

The abiotic factors and different dates of sowing were found to have high level of influence on the per cent parasitization by *C. chloridae* on *H. armigera* during the experimental period in tarai region. Maximum and minimum temperature had significant negative correlation with per cent parasitisation in early and normal sown crop compare to non significant negative correlation in late sown crop. Maximum and minimum humidity had significant positive correlation with per cent parasitisation in early sown crop.

## ACKNOWLEDGEMENT

The authors are grateful to Norman E. Borlaug crop research centre (NEB-CRC), G.B. Pant University of Agriculture and Technology for providing required facilities.

## REFERENCES

- Bisane, K.D., Nehare, S.K. and Khande, D.M. (2008). Extent of parasitization of *Helicoverpa armigera* (Hübner) by larval and pupal parasitoids on chickpea. *Journal of Entomological Research*. 32(4): 295-301.

- FAOSTAT-2017-18 (<http://www.fao.org/faostat/en/>).
- Kaur, S., Brar, K.S., Sekhon, B.S., Joshi, N., Shenhmar, M. and Singh, J. (2000). Role played by *Campoletis chloridae* Uchida in natural mortality of *Helicoverpa armigera* (Hübner) on chickpea in Punjab. *Journal of Biological Control*. 14 (1): 51-54.
- Kranthi, K.R., Jadhav, D.R., Kranthi, S., Wanjari, R.R., Ali, S.S. and Russell, D.A. (2002). Insecticide resistance in five major insect pests of cotton in India. *Crop Protection*. 21: 449-460.
- Ojha, P.K., Kumari, R. and Chaudhary, R.S., (2017). Impact of abiotic and biotic factors on population dynamics of *Helicoverpa armigera* Hubner (Noctuidae: Lepidoptera) in chickpea. *Journal of Entomology and Zoology Studies*. 5(1): 636-642.
- Pillai, A.K., Selvaraj, S. and Agnihotri, M. (2016). Seasonal abundance of *Campoletis chloridae* Uchida (Hymenoptera: Ichneumonidae), A larval parasitoid of *Helicoverpa armigera* (Hubner) in chickpea. *Legume Research-An International Journal*. 39(4): 643-647.
- Prakash, M.R., Ram, U. and Tariq, A. (2007). Evaluation of chickpea (*Cicer arietinum* L.) germplasm for the resistance to gram pod borer, *Helicoverpa armigera* Hubner (Lepidoptera: Noctuidae). *Journal of Entomological Research*. 31: 215-218.
- Reed, W., Cardona, C., Sithanatham, S. and Lateef, S.S. (1987). The chickpea insect pests and their control. In: Saxena, M.C. and Singh, K.B. (eds). *The Chickpea*. CAB International, Wallington, UK. pp. 283-318.