

nk/c {sk "keyh tuin eaxlsi"kykgrqpkjsdh miy0/krk ij ,d v/; ;u jkth*] jfolnz dckj] jktchj fl g ,oa jfolnz fl g

I jnkj cYyHk Hkkbz iVsy dfrk ,oaisk ksd fo"fo |ky;] ejB &250 110] mUkj cnskj Hkkjra
i klr%vDne] 2020

Lohdr% tuo]h 2021

I kjlk

mRrj insk ds "keyh tuin eai "kiky I kekftd fodkl eegRoikwzHkfedk fuHk jgk gA ; gkanwk: i "kykses xk; ikyu dk cgr cMk ; kxnu gA i "kiky dks dfrk fo"fo |ky; I jnkj cYyHk Hkkbz iVsy jkjk l e; & l e; ij tkudkj nh tkrh gA ftl eafHkx n/k mRiknu c<k l dA tuin dsrhu CykHkksdk l odf; k x; kA ftl es90 fdl kuka dks l cl svPNh f[kykbz rFk izl/k ds vk/kkj ij puk x; kA v/; ; u {k= es l cl svf/kd vxkyk ds 96-66% tksdkVdj f[kyk; k tkrk gA vxkyk dsl kFk&l kFk tS scj l he ; k l j l kadh dcl vDVne] l sekpZeghusrd f[kyk; k x; k tcf d dN gjspkjs tS sfd& tokgj] dchj cjl he vls ykfc; k 27-77] 2-33] 2-22 vls 2-22% ifr"kr ik; k x; kA T; knkrj ; gk fdl ku gjspkjs eavxkyk rFk Tokj dk iz kx djrsgA ftl es n/k mRiknu fnu ifrfnu c<rk tk jgk gA
'kn dfr%ued] xgw dk Hkk k] pkjk ds idkj] A

Bhartiya Krishi Anushandhan Patrika, 35(4): 267-269

Feed and fodder availability for cattle in shamli district of Uttar Pradesh

Rajeev*, Ravinder Kumar, Rajbir Singh and Ravinder Singh

Department of Animal Husbandry, College of Veterinary and Animal Science,

Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut-250 110, Uttar Pradesh, India.

Received: October 2020

Accepted: January 2021

ABSTRACT

A study was conducted on 90 respondents of cattle keepers from three blocks of Shamli district of Uttar Pradesh. During the study it was found that sugarcane tops were the main fodder with mixed with some green grasses or without grasses from October to May months. The mostly green fodder is available in their respective seasons. However, about 96.66% fodders are fed through chaffing. The sugarcane tops fed with mix of barseem or mustard cake or others concentrates in the months of January to May. Whereas, wheat straw is used by the 31.11% respondents with the mixed other green fodders or concentrate followed by Jawar, Karbi, Barseem and Lobia as 27.77, 3.33, 2.22 and 2.22%, respectively, different fodder was available in study area of Shamli district. In the study it was observed that cultivated green fodder in their land or either in rent field. The most cultivated green fodder in different season Sugarcane tops, Berseem, Lobia, and Jawer major fodder crops of the area.

Key words: Common salt, Concentrate, Type of feeds.

iLrkouk

mRrj insk es dfrk ds {k= eai "kiky dh cgr vge Hkfedk gA n/k mRiknu dsfy; s i "kiky vls dfrk dsfy; scNMk ikyu ij xtfk {k=ks es Ms jh vls xk; izl/k ds rjhdks ij v/; ; u fd; k x; kA "keyh tuin eaT; knkrj fdl ku xk; ikyus dk dk; Z djrsgA ; gk fdl ku dsikl de l sde 2&2 xk; gS yxHkx fdl ku 65% fdl ku xk; ksd ikyu djrsgA dN fdl ku xjhc

gsftl dsikl viuh dkbz xk; ugh gsvls fdl ku fdl h n/ jsfdl ku l sfjdk; s; k eghusdsfgl kc l sxk; yrs gS rFk mudk ikyu djrsgA dN fdl kuksdksde Kku gkus dsdkj.k og vPNh rjg l si "kiky ugh dj ukrsgA ftl l smUgsu d l ku dk l keuk djuk i Mrk gA bl l eL; k dks nj djus dsfy; s dfrk fo"fo |ky; us xkp&xkp es fjl pZNk=ks dksHkst k vls fdl kuksdh l eL; kvk dksckj ses tkudkj yH rFk fQj mu l eL; kvk dksnj fd; k ftl es fdl ku dksi "kiky l svf/kd l svf/kd vkenuh gks l dA

*Corresponding author's E-mail: drkashyap1952@gmail.com
I jnkj cYyHk Hkkbz iVsy dfrk ,oaisk ksd fo"fo |ky;] ejB &250 110] mUkj cnskj Hkkjra

Department of Animal Husbandry, College of Veterinary and Animal Science, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut-250 110, Uttar Pradesh, India.

rkfydk 1%

fo"ksrk, a	iR; k; h dh l Å; k	lk'r"kr
mez		
1 toku	07	07-77
2 e/;	41	45-55
3 cks	42	46-66
f"kk		
1 vui <	16	17-77
2 i <sfy[ksyk	04	04-44
3 fy[kus i <usokys	07	07-77
4 ikbejh f"kk	08	8-88
5 tfu; j gkbLdny	19	21-11
6 b.Vj fefM, V	31	34-44
7 Lukrd rFkk ml l s Åij	05	05-55
pkjs dh miyC/krk		
1 xlusdk Åijh Hkx	30	34-33
2 cktjk@dohz	03	03-33
3 xgwdk Hkx k	28	31-11
4 cjl he	02	02-22
5 Tokj	25	27-77
6 ykfc; k	02	02-22
;ks	90	

l lexb , oa ijhkk k fof/k

; g v/; ; u mRrj inš'k dsnf'Vdsk l svR; Ur
egroi wlgÅ "kkeyh tuin dsi "kikydk }jkk nkk: xk; s
l sl Ecfu/kr fofHku l eL; kvkadsckjsesdf'k fo"fo | ky;
ejB ds "kks'k Nk=ks }jkk l ošij vk/kfjr gÅ "kks'k v/; ; u
df'k fo"fo | ky; ds i "kikydu fofHkx }jkk dj; k x; k
gÅ bl v/; ; u es "kkeyh tuin dksppuk x; k FkA "kkeyh
tuin ds rhu Cykkl tš sfd "kkeyh] dšjuk rFkk vl;
Cykkl gÅ iR; d Cykkl l srhu&rhu xkpsdksppuk x; k FkA
tš s "kkeyh Cykkl l sykcl eqMš/dyk rFkk [kj] dšjuk
Cykkl l sdi škk dyk] d l škk [kpvš fVvšyh vl; Cykkl
l sFkeVh] ; q Q pšvj vkš jakuk xkpsdksv/; ; u dsfy; s
ppuk x; k FkA iR; d xkp l s10&10 fdl kukadksppuk x; k
Fk rFkk dy 90 fdl kuksdksv/; ; u dsfy; spuk x; k Fk
ftl ds ikl de&de 2 ; k 3 xk; sgk bu fdl kuks l s
tkudkj buVj0; wueus l s iklr dh x; hA uueus es tš s
fdl ku dh mej f"kk] HkfeJkr] ifjokj dk vkdkj] fl pkbz
dsL=kr] gjspkjsrFkk >qM dk vkdkj ; s l Hk tkudkj
uueusdsvk/kj ij yh x; hA iklr tkudkj dk eV; kadu
djrsgq sml sifr"kr }jkk fd; k x; k FkA

ifj.kk , oafoopuk

fdl kuks dh fo"ksrk; a

mez & bl v/; ; u es; g ik; k fd rhu mez l enj esckV
fy; k x; k 50 o'kz l sde 46-46% 36 l s50 o'kz es45-55%
rFkk 50 o'kz es7-77% FkA bl es56 o'kz esde dk vkš r
l cl svf/kd FkA rkfydk ¼½

f"kk & bl v/; ; u ea; g ik; k fd 34-44% fdl ku ek/
; fed 4-4% vui < FkA 17-77] 8-88] 7-7 rFkk 5-5 fdl ku/
e/; e] vui <] fy[k l drsrFkk Lukrd FkA

ifjokj dsl nL; & bl v/; ; u es; g ik; k fd 209
dy ifjokj dsl nL; FkA ftl eal s53-58% iq 'k 18 o'kz
l s Åij ds FksrFkk 31&70% fdl ku 6 l s18 dschp vkš
14-43% fdl ku 6 o'kz ds FkA rkfydk¼½

i"kkadk idkj & bl v/; ; u ea; g ik; k fd 28-80%
xk; snvk nusokyh FkA 21-10% ; u] 13-59% X; kfou]
10-6 [kyh xk; srFkk 3-14% l qM FkA

fl pkbz dsL=kr & "kkeyh tuin ds85-55% fdl kuks
dsikl Vošy dsL=ku gÅ dšy 14-44% fdl kuksdsikl
viuh dkbzL=kr ugh gÅ

pkjsdsfy; s[kr & [kjhQ ekš e eanyguh Ql y 29%

rFkk 24-04% vukt okyh FkhA tcfð jfo ekð e 24-04% ifr"kr nyguh rFkk 21-12 vukt okyh QI y FkhA

pkjsdh mi yC/krk % "kkeyh tuin esT; knkrj i "kikyð fdl ku vius [kr eadðk QI y dh [krh djrs gA ftl es xluuk mRiknu ; gk ds {ks= es l cl svf/kd gkrk gA ; gk fdl ku 88-88% vius [krks es xluuk mRiknu djrs gA fdl kuks }kjk xluuks dk uhps dk Hkkx fey dks nrs gS rFkk Åij dk Hkkx i "kq dks dkV dj f[kykrsgA 89% fdl ku ; g xluus ds Åijh Hkkx dks dkV dj f[kykrsgA xluus ds l kFk fdl ku gA ; g døy 5 ifr"kr fdl ku Tkogj ds /kii es l qk d j ml sviuh Nrks ij j [k yrk gA ; gk gjk pkjk ipj ek=k es mxk; k tkrk gA ftl es fdl kuks dks Mcy & Mcy vlenuh gkrh gA xluus dk Åijh Hkkx 34-33% dkV dj f[kykrsgA bl tuin esyxllx dy pkjses 35% xluus dk Åijh Hkkx 3-33] 31-11] 2-22] 27-27 vlg 2-22 ifr"kr ckt j k] Hkk k] cjl he]tokgj] rFkk ykfc; k vkfn pkjs mi yC/k FkA ; gk fdl ku i "kq/ks ds fy; s vius vki vukt dk feJ.k r\$ kj djrk gA T; knkrj fdl ku vukt dks ctkjkses l s [kjh yrk gS rFkk ml s ?kj ij fil rk gA

fu'd'k

v/; ; u l sfu'd'k fudy rk gS fd ftyes nllk: xk; ksl scgrj mRiknu ds fy; sgj spkjsdh mi yC/krk gA vko"; d gS dks & dks l s pkjs pkjs fdl ekð e es mxk l drsgA bl dh tkudkj h fdl kuks dks l e; & l e; ij feyrh jguh pkfg; A dñ fdl kuks dks gjs pkjs ds ckjs es ugh i rk jgrk gA bu fdl kuks dks df'k fo"fo | ky; }kjk l e; & l e; ij tkudkj h mi yC/k djkbz tk, ftl es fdl ku nll/k mRiknu c=k l dA bl ds l kFk cjl he dks feyk nrs gA ftl es i "kq bl s l pk: : lk l s [kk l dA tuin es fdl ku l kbyst r\$ kj ugh djrk A

l nHk

Aggarwal, S.B. and Sharma, K.N. (1986). Dairy management practices of Bovines in key village and non key village area around Karnal. *Indian J. Dairy Science*. **39**(1): 113-115.

Bhagyalazmi, K., Gopalakrishna Rao, V. and Sudarshanreddy, M. (2003). Profile of the rural women micro-entrepreneurs. *Journal of Research. Agricultural University, Hyderabad*. **31**(4): 51-54.

Chauhan, D.S., Kamble, V.J. Padghan, P.V, Sawant, R.C. and Kamble, R.R (2004). Impact of farmers' status on milk production in tribal area of Kinwat Tahasil (Marathwada Region). *Indian Journal of Animal Research*. **38** (2): 137-140.

Dubey, V.K. and Kumar, Ram (1981). Identifying existing animal husbandry practices followed by the cattle Keepers. *Dairy Guide*, **3**: 2.

Garg, M. K.; Jain, L. S.; Chaudhary, J. (2005). Studies on housing, feeding and milking management practices of dairy cattle in Baran District of Rajasthan. *Indian Journal of Dairy Science*. **58** (2): 123-128.

Gill, T.K., Saini, S.K.A. (2008). Study of awareness of recommended dairy practice among farmers *International Journal of Agricultural Sciences*. **4** (1): 296-300.

Garg, M. K., Jain, L.S., Chaudhary, J.L. (2005). Studies on housing, feeding and milking management practices of dairy cattle in Baran District of Rajasthan. *Indian Journal of Dairy Science*. **58** (2): 123-128.

Gill, T.K., Saini, S.K.A. (2008). Study of awareness of recommended dairy practices among farmers. *International Journal of Agricultural Sciences*. **4**(1): 296-300.

Gour, A.K., (2002). Factors influencing adoption of some improved animal husbandry practices of dairying in Anand and Vadadara district of Gujarat state. Ph.D. thesis, Gujarat Agricultural University. S.K. Nagar.

Hessian, Z.M.A., Husain, S.M.J., Rashid, M. M., Sultana, N., Ali, M.H. (2004). Study on the present management condition of private dairy farm at Rangpur Sadar Thana in Bangladesh. *Journal of Biological Sciences*. **4**(2): 185-188.